U.S. Department
of Commerce

Volume 99
Number 4
October 2001

# Fishery Bulletin

**U.S. Department of Commerce**
Donald L. Evans
Secretary

**National Oceanic and Atmospheric Administration**
Scott B. Gudes
Acting Under Secretary for Oceans and Atmosphere

**National Marine Fisheries Service**
William T. Hogarth
Acting Assistant Administrator for Fisheries

The *Fishery Bulletin* (ISSN 0090-0656) is published quarterly by the Scientific Publications Office, National Marine Fisheries Service, NOAA, 7600 Sand Point Way NE, BIN C15700, Seattle, WA 98115-0070. Periodicals postage is paid at Seattle, WA, and at additional mailing offices. POSTMASTER: Send address changes for subscriptions to Fishery Bulletin, Superintendent of Documents, Attn.: Chief, Mail List Branch, Mail Stop SSOM, Washington, DC 20402-9373.

Although the contents of this publication have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

The Secretary of Commerce has determined that the publication of this periodical is necessary according to law for the transaction of public business of this Department. Use of funds for printing of this periodical has been approved by the Director of the Office of Management and Budget.

**For sale** by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402. Subscription price per year: $45.00 domestic and $56.25 foreign. Cost per single issue: $28.00 domestic and $35.00 foreign. **See back for order form.**

# Fishery Bulletin

**Scientific Editor**
Dr. John V. Merriner

**Editorial Assistant**
Sarah Shoffler

Center for Coastal Fisheries and Habitat Research, NOS
101 Pivers Island Road
Beaufort, NC 28516

**Managing Editor**
Sharyn Matriotti
National Marine Fisheries Service
Scientific Publications Office
7600 Sand Point Way NE, BIN C15700
Seattle, Washington 98115-0070

**Editorial Committee**

| | |
|---|---|
| Dr. Andrew E. Dizon | National Marine Fisheries Service |
| Dr. Harlyn O. Halvorson | University of Massachusetts, Boston |
| Dr. Ronald W. Hardy | University of Idaho, Hagerman |
| Dr. Richard D. Methot | National Marine Fisheries Service |
| Dr. Theodore W. Pietsch | University of Washington, Seattle |
| Dr. Joseph E. Powers | National Marine Fisheries Service |
| Dr. Harald Rosenthal | Universität Kiel, Germany |
| Dr. Fredric M. Serchuk | National Marine Fisheries Service |

***Fishery Bulletin* web site: fishbull.noaa.gov**

The *Fishery Bulletin* carries original research reports and technical notes on investigations in fishery science, engineering, and economics. It began as the Bulletin of the United States Fish Commission in 1881; it became the Bulletin of the Bureau of Fisheries in 1904 and the *Fishery Bulletin* of the Fish and Wildlife Service in 1941. Separates were issued as documents through volume 46; the last document was No. 1103. Beginning with volume 47 in 1931 and continuing through volume 62 in 1963, each separate appeared as a numbered bulletin. A new system began in 1963 with volume 63 in which papers are bound together in a single issue of the bulletin. Beginning with volume 70, number 1, January 1972, the *Fishery Bulletin* became a periodical, issued quarterly. In this form, it is available by subscription from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402. It is also available free in limited numbers to libraries, research institutions, State and Federal agencies, and in exchange for other scientific publications.

U.S. Department
of Commerce
Seattle, Washington

Volume 99
Number 4
October 2001

# Fishery Bulletin

## Contents

*Articles*

519–527 **Arendt, Michael D., Jon A. Lucy, and Thomas A. Munroe**
Seasonal occurrence and site-utilization patterns of adult tautog, *Tautoga onitis* (Labridae), at manmade and natural structures in lower Chesapeake Bay

528–544 **Gharrett, Anthony J., Andrew K. Gray, and Vladimir Brykov**
Phylogeographic analysis of mitochondrial DNA variation in Alaskan coho salmon, *Oncorhynchus kisutch*

545–553 **Hurst, Thomas P., and David O. Conover**
Diet and consumption rates of overwintering YOY striped bass, *Morone saxatilis,* in the Hudson River

554–571 **Lo, Nancy C. H., John R. Hunter, and Richard Charter**
Use of a continuous egg sampler for ichthyoplankton surveys: application to the estimation of daily egg production of Pacific sardine (*Sardinops sagax*) off California

572–587 **Loher, Timothy, David A. Armstrong, and Bradley G. Stevens**
Growth of juvenile red king crab (*Paralithodes camtschaticus*) in Bristol Bay (Alaska) elucidated from field sampling and analysis of trawl-survey data

588–600 **MacNair, Leslie S., Michael L. Domeier, and Calvin S. Y. Chun**
Age, growth, and mortality of California halibut, *Paralichthys californicus,* along southern and central California

601–616 **Miyashita, Shigeru, Yoshifumi Sawada, Tokihiko Okada, Osamu Murata, and Hidemi Kumai**
Morphological development and growth of laboratory-reared larval and juvenile *Thunnus thynnus* (Pisces: Scombridae)

The conclusions and opinions expressed in *Fishery Bulletin* are solely those of the authors and do not represent the official position of the National Marine Fisheries Service (NOAA) or any other agency or institution.

The National Marine Fisheries Service (NMFS) does not approve, recommend, or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

**617–627** **Patterson III, William F, James H. Cowan Jr, Charles A. Wilson, and Robert L. Shipp**
Age and growth of red snapper, *Lutjanus campechanus,* from an artificial reef area off Alabama in the northern Gulf of Mexico

**628–640** **Sipe, Ann M., and Mark E. Chittenden Jr.**
A comparison of calcified structures for aging summer flounder, *Paralichthys dentatus*

**641–652** **Somerton, David A., and Peter Munro**
Bridle efficiency of a survey trawl for flatfish

**653–664** **Wilson, Charles, A., and David L. Nieland**
Age and growth of red snapper, *Lutjanus campechanus,* from the northern Gulf of Mexico off Louisiana

*Notes*

**665–670** **Arendt, Michael D., John E. Olney, and Jon A. Lucy**
Stomach content analysis of cobia, *Rachycentron canadum,* from lower Chesapeake Bay

**671–678** **Blandon, Ivonne R., Rocky Ward, Tim L. King, William J. Karel, and James P. Monaghan Jr.**
Preliminary genetic population structure of southern flounder, *Paralichthys lethostigma,* along the Atlantic Coast and Gulf of Mexico

**679–684** **Hernández-López, José L., José J. Castro-Hernández, and Vicente Hernández-García**
Age determined from the daily deposition of concentric rings on common octopus (*Octopus vulgaris*) beaks

**685–690** **Jones, Thomas S., and Karl I. Ugland**
Reproduction of female spiny dogfish, *Squalus acanthias,* in the Oslofjord

**691–696** **Porter, Steven M., Annette L. Brown, and Kevin M. Bailey**
Estimating live standard length of net-caught walleye pollock (*Theragra chalcogramma*) larvae using measurements in addition to standard length

**697–701** **Takagi, Motohiro, Tetsuro Okamura, Seinen Chow, and Nobuhiko Taniguchi**
Preliminary study of albacore (*Thunnus alalunga*) stock differentiation inferred from microsatellite DNA analysis

**702** *Acknowledgment of reviewers*

**703** *Index*

**711** *Erratum*

**712** *Subscription form*

# Seasonal occurrence and site-utilization patterns of adult tautog, *Tautoga onitis* (Labridae), at manmade and natural structures in lower Chesapeake Bay*

**Michael D. Arendt**

School of Marine Science
College of William and Mary
Virginia Institute of Marine Science
Gloucester Point, Virginia 23062
Present address: Marine Resources Division, Department of Natural Resources
Marine Resources Research Institute
217 Fort Johnson Road
Charleston, South Carolina 29422-2559
E-mail address: arendtm@mrd.dnr.state.sc.

**Jon A. Lucy**

Sea Grant Marine Advisory Program
Virginia Institute of Marine Science
Gloucester Point, Virginia 23062

**Thomas A. Munroe**

National Marine Fisheries Service National Systematics Laboratory
National Museum of Natural History
Washington, D.C. 20560-0153

**Abstract**—Ultrasonic transmitters were surgically implanted into adult tautog ($n$=27, 400–514 mm TL) to document seasonal occurrence and site utilization at four sites situated within known tautog habitat near Cape Charles, Virginia, in lower Chesapeake Bay. Tagged tautog were released at the same sites where originally caught within 2 h of capture. Sites were continuously monitored with automated acoustic receivers between 9 November 1998 and 13 October 1999. Two sites consisted of natural bedform materials and two sites consisted of manmade materials. Ninety-four percent of tautog ($n$=15) released in fall 1998 remained inshore during winter at sustained water temperatures of 5–8°C, rather than moved offshore during winter as documented for tautog off New York, Rhode Island, and Massachusetts. Ninety-one percent ($n$=10) of tautog released in spring 1999 remained inshore during summer when water temperature was 27°C and in the absence of an important food item, blue mussels (*Mytilus edulis*). These findings conflict with assertions that tautog move to cooler water in summer when water temperatures reach 20°C. Tautog released at natural bedform sites were detected only at these sites throughout the study. Tautog released at manmade structures also displayed high site-utilization patterns, but several tautog periodically moved 2–10.2 km away from these sites over featureless bottom, a known deterrent to emigration for large temperate labrids in other waters. Benthic communities were similar at manmade sites and natural bedform sites, and movement away from manmade sites may have been influenced by habitat size as well as habitat structure. Understanding temporal and spatial utilization of habitats is an important first step to identifying essential fish habitat and to evaluating and protecting fishery resources within Chesapeake Bay and elsewhere.

Manuscript accepted 11 April 2001.
Fish. Bull. 99:519–527 (2001).

The labrid *Tautoga onitis* (tautog) is a highly prized game fish targeted by anglers fishing at natural and manmade structure (Briggs, 1977; Lucy and Barr, 1994). Tautog are distributed between Georgia (Parker, 1990) and Nova Scotia (Bigelow and Schroeder, 1953); peak abundance is found between Massachusetts and the Delaware Capes.[1] Slow growth rate, late age at maturity, predictable distribution, and localized population structure suggest high vulnerability to overexploitation (Hostetter and Munroe, 1993). Extended residence at accessible fishing sites may increase the potential for overexploitation; thus, residence and site-utilization patterns of tautog throughout this species' distribution range must be well understood for effective management of this resource.

Tag-recapture studies in New York, Rhode Island, and Massachusetts suggest that adult tautog spend spring and fall months inshore, may move offshore during the warmest summer months (Cooper, 1966; Briggs, 1969), and overwinter offshore (Cooper, 1966; Briggs, 1977). Tautog leave inshore waters at varying rates between July and October (Cooper, 1966) and are recaptured in coastal waters in fall (Cooper, 1966; Briggs, 1977), consistent with indirect observations on seasonal abundance (Stolgitis, 1970; Olla et al., 1974). In contrast, tag-recapture studies report limited evidence of a seasonal inshore-offshore migration for tautog in the Chesapeake Bay and coastal Virginia waters.[2] Seasonal abundance data also suggest that tautog remain inshore in Chesapeake Bay (Hostetter and Munroe, 1993) and in Delaware Bay (Eklund and Targett, 1991) during winter.

* Contribution 2391 of the Virginia Institute of Marine Science, Gloucester Point, Virginia 23062.

[1] Atlantic States Marine Fisheries Commission (ASMFC). 1996. Fishery management plan for tautog, rep. 25, 56 p. [Available from ASMFC, 1444 Eye Street NW, Washington, DC 20005.]

[2] Virginia Game Fish Tagging Program. 1995–1999. Marine Resources Commission, 968 Oriole Dr. South, Suite 102, Virginia Beach, VA 23451.

The tag-recapture method is not a suitable method for evaluating site-utilization patterns because this technique does not provide information on the location of tagged animals between times of release and recapture. Furthermore, the tag-recapture method requires that tagged animals be recaptured, and be reported as recaptured, before any information is available. Ultrasonic telemetry, however, enables continuous observations on all tagged animals, in their natural environment, without requiring that tagged animals be recaptured (Winter, 1996). Previously, only in one other study (Olla et al., 1974) was ultrasonic telemetry used to monitor adult tautog. Olla et al. (1974) tagged and "tracked" 10 adult tautog ultrasonically in Great South Bay, NY, for up to 80 h after their release. Although an important study, their sample size was small and total observations too limited (<400 h, single season) to document seasonal occurrence and site-utilization patterns for this species.

Ultrasonic telemetry was selected to address seasonal occurrence and site-utilization patterns of adult tautog in lower Chesapeake Bay, given that tag-recapture methods can be applied only within limitations and given that the poor visibility and strong currents preclude direct underwater observations of this species in this turbid estuary. Rather than collect detailed positional data over short periods of time (days) for a few tautog, we chose to collect seasonal occurrence and site-utilization data for two large groups of tautog ($n$=16 and 11) at four specific sites located within known tautog habitat. Sites were monitored by using a fixed, submerged hydrophone array between November 1998 and September 1999. The first objective of our study was to determine if tautog remained inshore at natural and manmade structures in lower Chesapeake Bay during winter and summer. The second objective was to document and describe site-utilization patterns within inshore study sites. Data for daily activity patterns are presented elsewhere (Arendt et al., in press).




**Figure 1**

Location of study sites for telemetric study of tautog released in lower Chesapeake Bay near Cape Charles, VA. Texeco Wreck (TX) is located in 18 m of water on a plain west of Susquehanna Channel (30–40 m deep). Coral Lump (CL), Ridged Bottom (RB), and Airplane Wreck (AW) are located in 8–15 m of water on a flat east of the Susquehanna Channel.

## Materials and methods

Tautog were caught, tagged, and released at four sites situated within a 1.5 km × 6 km area near Cape Charles, Virginia (Fig. 1). Side-scan sonar (Sea Scan Technology, Ltd., White Marsh, VA) was used to measure dimensions of the four study sites and to map the surrounding seafloor. The Texeco Wreck, a 30 m × 100 m shipwreck, was located in 18 m of water west of the Susquehanna Channel (30–40 m deep) in an area characterized by flat, relatively featureless bottom topography (Wright et al., 1987). The three remaining sites (Airplane Wreck, Coral Lump, and Ridged Bottom) were located in 8–15 m of water east of the Susquehanna Channel in an area characterized by sand flats and deep, mud-bottomed channels (Wright et al., 1987). The Airplane Wreck (40 m × 20 m) consisted of concrete rubble. The Ridged Bottom (30 m × 100 m) and Coral Lump (100 m × 300 m) sites consisted of natural bedforms. Otter trawl, oyster dredge, and underwater video surveys indicated that all sites were densely populated by several species of sponges, colonial bryozoans, mollusks, and crustaceans.

Tautog were caught with standard two-hook bottom rigs baited with pieces of blue crab or clam and were brought aboard with a nylon landing net. Tautog were observed in an aerated live well up to 2 h before transmitters were implanted. Total length (mm) and sex (White, 1996) were recorded. Only tautog >400 mm TL were tagged ultrasonically. This minimum size increased the odds of transmitters weighing less than 1.25% of fish body weight in water (Winter, 1996), based on size-weight relationships for tautog in Virginia (Hostetter and Munroe, 1993; White, 1996). Tautog >400 mm were also reproductively mature (Hostetter and Munroe, 1993; White, 1996).

Surgical procedures were similar to those used in Nemetz and Macmillan (1988), Mortensen (1990), Holland et al. (1993), Szedlmayer (1997), and Thoreau and Baras (1997). In preparation for surgical implantation of transmitters, level-four anesthesia (Mattson and Ripple, 1989; Prince et al., 1995) was induced by immersing tautog in a 325-mg/L solution of MS-222. Once anesthetized, a small

(25-mm) incision was made on each fish immediately dorsal to the ventral midline between pelvic fins and anus with a sterilized, disposable razor blade. The transmitters was coated in sterile mineral oil and placed into the visceral cavities of tautog with the transducer end of the transmitter facing forward. Braided, polyglycolic acid sutures (Dexon®, I-III), surgical staples (Proximate Plus MD 35W®), and acrylic adhesive (Krazy glue®) were used to close the incision. Betadine was used periodically throughout the surgical procedure and antibiotics (Nu-Flor®) were injected intramuscularly to increase postsurgical survival (Schramm and Black, 1984; Bart and Dunham, 1990; Poppe et al., 1996). Tautog were revived in the aerated live well and released within 0.5 h after surgery. Preliminary evaluation of surgical procedures with "dummy" transmitters indicated 100% transmitter retention, 86% survival, and normal swimming, feeding, reproductive behavior, and physiology for tautog >400 mm TL held up to 418 days in captivity (Arendt, 1999). Tautog were fully recovered from surgery <1 to 6 days after release (Arendt, 1999) according to detection patterns recorded by automated acoustic receivers (Arendt and Lucy, 2000).

V-16-1H-R256 coded transmitters (16 mm × 48 mm, 9 g in water; Vemco, Ltd., Shad Bay, Nova Scotia, Canada) were used in our study. Signal repeat intervals for coded transmitters (69 kHz) varied randomly between 45 and 75 s, which extended battery life to 111 d. Transmitters were primarily detected with a submerged array of automated acoustic receivers (VR1, Vemco, Ltd.); however, transmitters were also detected from a research vessel with acoustic hydrophones (V10, VH65, Vemco, Ltd.) and an electronic receiver (VR60, Vemco, Ltd.).

Omnidirectional VR1 receivers were deployed 100–150 m to the west and east of the perimeter of each of the four sites. Detection radius for each receiver was approximately 400 m. Detection areas for both receivers were overlapped to create three distinct reception zones: a central reception zone common to both receivers and two peripheral reception zones unique to either receiver. VR1 receivers were moored 1.5–3.0 m above the seafloor to provide an unobstructed line-of-sight for transmitter signal reception (i.e. positioned above the "structure" associated with each site) and to reduce acoustic interference from suspended material associated with strong bottom currents. Mooring units consisted of a railroad wheel (227 kg), stainless steel aircraft cable (0.64 cm, 7 × 19 strand), and subsurface and surface floats. Receivers were retrieved every three to six weeks and detection data (transmitter ID, date and time of detection) were downloaded directly to a shipboard computer by means of a VR1-PC cable interface (Vemco, Ltd.).

Data for tautog released in fall 1998 were collected for the duration of the transmitter battery life. Data for tautog released in spring 1999 were collected until all VR1 receivers were permanently removed from each site. VR1 detections for each tautog were sorted into hourly bins and examined graphically. Because tautog are diurnally active and nocturnally quiescent (Olla et al., 1974; Arendt et al., in press), only daytime detection was used to determine site-utilization patterns. Tautog were considered resident at a particular site each day if they were detected at the same site throughout the day (morning, mid-day, evening). Total fish-days (sum of all days between date of first and last detection for all fish) for each calendar season were classified as 1) days when tautog were resident at the site of initial release, 2) days when tautog were detected at an alternative site, or 3) days when they were not detected at all. Site-utilization patterns of tautog were examined for each site in fall (from 9 Nov 98 to 20 Dec 98), winter (from 21 Dec 98 to 20 Mar 99), spring (from 21 Mar 99 to 20 Jun 99), and summer (from 21 Jun 99 to 9 Sep 99).

To increase the probability that tautog would be reported as recaptured should recapture occur, transmitters were labeled with the specimen's ID number, a $50 "reward" notice, and a phone number. Tautog were tagged externally with a small, orange t-bar anchor tag (TBA2; Hallprint, Holden Hill, South Australia) used by the Virginia Game Fish Tagging Program and with a larger, green t-bar tag (SHD-95; Floy Mfg., Seattle, WA) containing the specimen's ID number, a $50 reward" notice, and a phone number. Internal and external reward notices were included because Szedlymayer (1997) had observed that internal reward notices persisted longer than external reward notices for red snapper (*Lutjanus campechanus*) in the Gulf of Mexcio, and that internal reward notices were noticed accidentally. In addition to distinct marking of each tautog, colorful reward posters were posted at over 40 marinas, boat ramps, and tackle shops in lower Chesapeake Bay and literature describing the project was mailed to over 5000 homes and businesses.

## Results

Twenty-seven adult tautog (400–514 mm TL) were tagged with ultrasonic transmitters and released (16 in fall 1998, 11 in spring 1999) near Cape Charles, VA (Table 1). Four tautog were released at each of the four sites in fall 1998 and at the Coral Lump and Texeco Wreck in spring 1999. Two tautog were released at the Ridged Bottom and one tautog was released at the Airplane Wreck in spring 1999. Similar numbers of tautog were tagged and released at manmade ($n$=13) and natural bedform ($n$=14) sites.

Eighty-one percent ($n$=22) of all tautog released were males; 19% were females. Thirteen percent ($n$=2) of tautog released in fall 1998 were female; 27% ($n$=3) of all tautog released in spring 1999 were female. Both female tautog released in fall 1998 (ID19, ID28) were released at the Texeco Wreck. One female tautog was released at the Texeco Wreck (ID37) and two female tautog were released at the Coral Lump (ID39, ID40) in spring 1999.

Ninety-four percent ($n$=15 of 16) of tautog released in fall 1998 remained inshore within lower Chesapeake Bay during winter. Ninety-one percent ($n$=10 of 11) of tautog released in spring 1999 remained inshore within the Bay during summer.

All tautog ($n$=14) released at natural sites remained inshore and were detected only at their respective release sites. Tautog released at the Ridged Bottom and Coral Lump sites were detected 99% of fish-days in fall, 71–91% of fish-days in winter, 64–100% of days in spring, and

**Table 1**

Summary of fish-days for data from 27 adult tautog (400–514 mm TL) tagged and released with ultrasonic transmitters at four sites in lower Chesapeake Bay near Cape Charles, Virginia. Recaptured tautog are noted with an asterisk (*). Abbreviations for sites: CL = Coral Lump; TX = Texeco Wreck; RB = Ridged Bottom; AW = Airplane Wreck.

| | | | | | Fish days | | | |
|---|---|---|---|---|---|---|---|---|
| ID | Sex | TL | Site | Date released | At site | Not detected | At alternate site | Total |
| 1 | M | 432 | CL | 9 Nov 1998 | 174 | 8 | 0 | 182 |
| 18 | M | 406 | CL | 9 Nov 1998 | 155 | 19 | 0 | 174 |
| 19 | F | 495 | TX | 10 Nov 1998 | 58 | 105 | 2 | 165 |
| 20* | M | 470 | TX | 10 Nov 1998 | 0 | 167 | 1 | 168 |
| 21 | M | 406 | RB | 10 Nov 1998 | 88 | 10 | 0 | 98 |
| 22 | M | 400 | RB | 10 Nov 1998 | 178 | 0 | 0 | 178 |
| 23 | M | 483 | AW | 13 Nov 1998 | 152 | 13 | 0 | 165 |
| 24 | M | 432 | AW | 13 Nov 1998 | 112 | 45 | 0 | 157 |
| 25 | M | 432 | CL | 3 Dec 1998 | 183 | 3 | 0 | 186 |
| 26 | M | 400 | CL | 3 Dec 1998 | 177 | 4 | 0 | 181 |
| 27 | M | 514 | TX | 4 Dec 1998 | 177 | 0 | 0 | 177 |
| 28 | F | 413 | TX | 4 Dec 1998 | 42 | 130 | 13 | 185 |
| 29* | M | 400 | AW | 7 Dec 1998 | 147 | 14 | 1 | 162 |
| 30 | M | 419 | AW | 7 Dec 1998 | 46 | 21 | 0 | 67 |
| 31 | M | 445 | RB | 8 Dec 1998 | 144 | 24 | 0 | 168 |
| 32 | M | 419 | RB | 8 Dec 1998 | 88 | 39 | 0 | 127 |
| 33 | M | 406 | TX | 21 Apr 1999 | 0 | 136 | 6 | 141 |
| 34* | M | 432 | CL | 28 May 1999 | 69 | 0 | 0 | 69 |
| 35 | M | 445 | TX | 28 May 1999 | 97 | 7 | 0 | 104 |
| 36 | M | — | TX | 28 May 1999 | 103 | 1 | 0 | 104 |
| 37* | F | 445 | TX | 28 May 1999 | 99 | 5 | 0 | 104 |
| 38* | M | 483 | CL | 7 Jun 1999 | 59 | 0 | 0 | 59 |
| 39* | F | 483 | CL | 7 Jun 1999 | 59 | 0 | 0 | 59 |
| 40* | F | 432 | CL | 7 Jun 1999 | 59 | 0 | 0 | 59 |
| 41 | M | 445 | AW | 7 Jun 1999 | 10 | 2 | 2 | 14 |
| 42* | M | 406 | RB | 9 Jun 1999 | 58 | 0 | 0 | 58 |
| 43 | M | 406 | RB | 9 Jun 1999 | 56 | 2 | 0 | 58 |

98–100% of fish-days in summer (Fig. 2). Reduced detection in winter and spring was partially attributed to two tautog (ID21, ID32) that were detected 46 d and 75 d less than the mean (175 d) for other tautog ($n$=12) released at the same time. To provide a more conservative estimate of site utilization, these tautog were listed as "not detected" at these sites for these days.

Eighty-five percent ($n$=11) of tautog released at man-made structures remained inshore and all but three of these were detected only at their respective release sites. Site utilization by tautog at the Texeco Wreck was low (34–71% of fish-days) in all seasons (Fig. 2). One tautog (ID20) released at the Texeco Wreck was only detected at the Texeco Wreck for three hours after being tagged and released in fall 1998. Two additional tautog (ID19, ID28) spent 64–70% of fish-days away from the Texeco Wreck in fall, winter, and spring, and a fourth tautog (ID33) spent 100% of fish-days away from the Texeco Wreck in spring and summer. One of these tautog (ID28) was detected at the Coral Lump for 6 days in January 1999 (VR1 receivers) and all three tautog were regularly detected with the VR60 receiver at a site 2 km south of the Texeco Wreck. Tautog released at the Airplane Wreck were resident 99% of fish-days in fall, 66% of fish-days in winter, and 58% of fish-days in spring (Fig. 2). Reduced detection in winter resulted partially from one tautog (ID30) being detected 108 days less than the mean (175 days) for other tautog ($n$=12) released at the same time. To provide a more conservative estimate of site utilization, this tautog was listed as "not detected" at the Airplane Wreck for these days. No summer data were available for tautog at the Airplane Wreck because the single tautog (ID41) released there in spring 1999 was detected at this site for two days following release, was later detected at the Texeco Wreck, and then was never detected again at any site. Tautog 41 was listed as "not detected" at the Airplane Wreck from the time of




**Figure 2**

Seasonal site-utilization patterns of tautog released at manmade (Airplane Wreck and Texeco Wreck) and natural bedform (Ridged Bottom and Coral Lump) sites near Cape Charles, Virginia, in fall (9 Nov 98 to 20 Dec 98), winter (21 Dec 98 to 20 Mar 99), spring (21 Mar 99 to 20 Jun 99), and summer (21 Jun 99 to 9 Sep 99). Site-utilization patterns were greatest for tautog released at natural bedform sites.

leaving this site until VR1 receivers were removed from this site, 112 d later.

Eight tautog (27%) released in our study were subsequently recaptured (Table 1). Two tautog (ID20, ID29) released in fall 1998 were recaptured (13%) away from release sites by commercial fishermen in spring 1999. Tautog 20 was released at the Texeco Wreck on 10 November 1998 and either left this site the same day or its transmitter failed to transmit data. This tautog was subsequently recaptured 10.2 km northeast of the Texeco Wreck in a crab pot on 27 April 1999, 169 days later. Tautog 29 was first caught at the Airplane Wreck on 13 November 1998 and held in a wire cage with several other tautog at the Airplane Wreck for five days as part of a catch-release mortality study on tautog.[3] After being released at the Airplane Wreck on 18 November 1998, this tautog was recaptured at the Airplane Wreck on 7 December 1998 (19 d later) and ultrasonically tagged and released. Tautog 29 remained at the Airplane Wreck until 12 May 1999, then was recaptured in a gill net 2 km east of the Airplane Wreck on 19 May 1999. Recreational fishermen recaptured six tautog released in spring 1999 (55%) at the same sites where these fish were released 114–211 d earlier and 8–13 weeks after VR1 receivers were removed from sites.

## Discussion

Tautog remained inshore within lower Chesapeake Bay during winter, at sustained water temperatures of 5–8°C (Arendt et al., in press). Although detected at these water temperatures, tautog overall were detected less than during other times of the year, most likely because tautog remained inactive and within structures for several days at a time (Arendt et al., in press). Inshore occurrence of tautog in winter has been observed in eastern Long Island Sound,[4] Delaware Bay (Eklund and Targett, 1991), and lower Chesa-

[3] Lucy, J. A., and M. D. Arendt. 1999. Exploratory field evaluation of hook-release mortality in tautog (*Tautoga onitis*) in Lower Chesapeake Bay, Virginia. Rep. VMRC-99-10, 11 p. [Available from Marine Advisory Program, Virginia Institute of Marine Science, Gloucester Point, VA 23062.]

[4] Auster, P. J. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North and Mid-Atlantic)—tautog and cunner. U.S. Fish and Wildlife Service Biological Report 82 (11.105). U.S. Army Corps of Engineers Rep. TR EL-82-4, 13 p. NOAA's National Undersea Research Program, Univ. Connecticut at Avery Point, Groton, CT 06430.

peake Bay (Hostetter and Munroe, 1993). When water temperatures remains above 9–10°C,[5] a viable winter inshore fishery exists for tautog within lower Chesapeake Bay. Within the winter fishery, inshore catches occur predominantly in December and March, whereas offshore catches occur predominantly in January and February. Occurrence of an inshore winter fishery for tautog in Virginia is unique within this species' geographic distribution.

Tautog remained inshore during the summer at a maximum sustained water temperature of 27°C (Arendt et al., in press). Summer residence data have been supported by direct underwater observations.[6] Infrequent recreational catches of tautog in lower Chesapeake Bay during summer have also been reported.[2] Inshore, summer residence of tautog has been documented for Great South Bay, NY, when water temperatures were 19–24°C (Olla et al., 1974) and Narragansett Bay, RI, at maximum sustained water temperatures of 22°C.[7] These findings contradict reports from Virginia (Adams, 1993), New York (Briggs, 1969), and Rhode Island (Cooper, 1966) that adult tautog may move offshore to cooler water during summer.

Tautog remained inshore during summer in the absence of blue mussels (*Mytilus edulis*), a primary food item of tautog in northern areas (Bigelow and Schroeder, 1953; Olla et al., 1974). In June 1998–99, using underwater video, otter trawl, and oyster dredge, we documented large clusters of live blue mussels at study sites and noted growth of mussels on VR1 mooring units. By July 1998–99, blue mussels were not present at any of these sites. Absence of mussels in July in both years was most likely due to lethal effects of water temperatures ≥27°C (Wells and Gray, 1960).

Because blue mussels are not present in lower Chesapeake Bay year round, the diet of tautog inhabiting lower Chesapeake Bay throughout the year may be more diverse than that of tautog in northern areas. Stomach contents from an ultrasonically tagged tautog recaptured in October 1999 at the Ridged Bottom site consisted primarily of the bryozoan *Alcyinidium verilli*. At an artificial fishing reef near Cape Charles, VA, tautog consumed a variety of crustaceans, shellfish, bryozoans, and hydroids.[8] Similar temporal distributions of blue mussels have been reported in Delaware Bay, where diets of tautog subsequently shift towards alternative, less nutritious food items at times when blue mussels are unavailable.[9]

Year-round occurrence of adult tautog in Chesapeake Bay differs from seasonal (spring and fall) inshore occurrence of adult tautog in Great South Bay, NY (Olla et al., 1974; Briggs, 1977), Narragansett Bay, RI (Cooper, 1966), and the Weweantic River Estuary, MA (Stolgitis, 1970). Year-round occurrence of ultrasonically tagged tautog was consistent with large-scale patterns of occurrence of tautog from conventionally tagged tautog (127–584 mm TL) in the Virginia Game Fish Tagging Program between 1995 and 1999.[2,5] Of 563 recaptured tautog that were originally tagged in lower Chesapeake Bay, excluding Cape Charles, and adjacent coastal waters, 85% ($n$=476) were recaptured at the same sites where released 0–1214 d earlier, including 20 multiple recaptures of individuals at the same sites where originally released. High incidence of recaptures of tautog at the same sites where they were released occurred during all seasons. Only 5% of all recapture events involved movement (8–97 km) of tautog between Chesapeake Bay and adjacent coastal waters.

Daily detection patterns were almost always similar for both VR1 receivers at sites, indicating that tautog remained within the central signal reception area of both VR1 receivers and in the general vicinity of sites throughout the entire day. Tautog were generally not detected by VR1 receivers at night; however, tautog likely remained at sites throughout the night (Arendt et al., in press). Tautog were likely detected less often (or not at all) at night because of nocturnal quiescence in or near structure (Olla et al., 1974) and therefore were effectively out of range of VR1 receivers because of the presence of an acoustic barrier (Matthews, 1992; Pearcy, 1992; Bradbury et al., 1995,1997; Zeller, 1997).

Featureless bottom topography, a known deterrent to emigration for large, temperate labrids (*Notolabrus tetricus, N. fucicola, Pictilabrus laticlavius, Pseudolabrus psittaculus)* in Tasmania (Barrett, 1995), did not act as a deterrent to emigration for tautog in lower Chesapeake Bay. Two tautog (ID20, ID28) released at the Texeco Wreck in fall 1998 traversed a wide (2-km), deep (37–40 m) mud-bottomed channel (Wright et al., 1987) on at least three occasions, and one of these tautog (ID20) was subsequently recaptured. Two tautog (ID19, ID28) traveled between the Texeco Wreck and a site 2 km south of the Texeco Wreck on at least 12 occasions. A third tautog (ID33) left the Texeco Wreck almost immediately after being released and was subsequently detected (VR60 receiver) at this site throughout the remainder of our study.

Movement by tautog was assumed to represent actual movement by tagged tautog as opposed to movement of a

[5] White, G. G., J. E. Kirkley, and J. A. Lucy. 1997. Quantitative assessment of fishing mortality for tautog (*Tautoga onitis*) in Virginia. Preliminary report, 54 p. Department of Fisheries Science and Marine Advisory Program, Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, VA 23062.

[6] Hager, C. 1999 (July). Personal communication of direct underwater observation of tautog at Plantation Light (3–8 m depth), 2 km southeast of Texeco Wreck study site. School of Marine Science, College of William and Mary, Virginia Institute of Marine Science, Gloucester Point, VA 23062.

[7] Castro, K. 1999 (October). Personal communication of water temperature observation for Narragansett Bay. East Farm–Fisheries Center, Univ. Rhode Island, Kingston, RI 02881.

[8] Feigenbaum, D., C. Blair, and A. J. Provenzano. 1985. Artificial reef study—year II report. Virginia Marine Resources Commission rep. VMRC-83-1185-616, 57 p. VA Institute of Marine Science, Gloucester Point, VA 23062.

[9] Steimle, F., K. Foster, W. Muir, and B. Conlin. 1999. The diet of tautog collected on an artificial reef in Delaware Bay and interannual effects of prey availability (and notes on other tautog diet studies in the middle Atlantic Bight). First Biennial Conference on the biology of tautog and cunner, Mystic, CT, 30 November–1 December 1999. National Marine Fisheries Service, Highlands, NJ 07732.

predator that may have preyed upon tagged tautog, even though these three tautog were never recaptured nor visually observed after release. Adult tautog are very infrequently preyed upon by sharks in Virginia (Gelsleichter et al., 1999); however, sharks are not likely present in Chesapeake Bay when water temperatures are 5–8°C. At these water temperatures, large striped bass (*Morone saxatilis*) pose the only possible predatory threat to tautog. A recent study on feeding habits of adult striped bass in Chesapeake Bay found no tautog in the stomachs of more than 2000 striped bass, many of which were collected from the Chesapeake Bay Bridge-Tunnel complex, a well-known fishing area for adult tautog (Walters, 1999).

All tautog detected or recaptured away from release sites were released at manmade sites, which also happened to be the smallest sites. No information was available regarding the origin of these manmade sites, but both have been in place for at least 20 years.[10,11] Stone et al. (1979) concluded that artificial reefs reach a stable state after five years. Benthic macrofauna collected at manmade sites during our study were similar to benthic macrofauna collected at natural sites, suggesting that food may have been similar between manmade and natural sites. Given these observations, habitat size may be an important factor for adult tautog in determining the scale of local movements between adjacent habitats. Understanding the relationship between habitat size and site utilization warrants further investigation, especially with recent increased interest in the construction of artificial habitats for purposes of stock enhancement and enhanced fishing opportunities for tautog.

Sex ratio of female to male tautog in our study was heavily skewed (1:3.5) towards male tautog due to opportunistic sampling, and the preponderance of male tautog likely contributed to the high levels of site utilization observed in our study. In laboratory settings, adult male tautog aggressively defend territories throughout most of the year (Olla et al., 1978, 1980) and only during the spawning season are female tautog permitted to enter territories (Olla and Samet, 1977; Olla et al., 1981). Overall activity, including male agonistic behavior, also decreases as water temperatures approach annual minimum and maximum values (Olla et al., 1978, 1980). Although sample size in our study was too small to distinguish site-utilization patterns by sex, it is worth noting that both tautog that left the Texeco Wreck in Nov–Dec and that periodically returned to this site throughout the winter and spring were females. In contrast, during the spring spawning season, three females (one at Texeco Wreck, two at Coral Lump) remained at release sites throughout the spring–summer monitoring period and all three were subsequently recaptured by recreational fishermen at these same sites in the fall. More sex-specific data are needed to fully comprehend what role reproductive biology and social structure have on seasonal site-utilization patterns.

Site-utilization patterns exhibited by ultrasonically tagged tautog were consistent with patterns reported for tautog released at these same sites from the Virginia Game Fish Tagging Program (VGFTP). Between April 1998 and October 1999, 40 tautog, tagged and released at these sites, were recaptured, including one tautog recaptured twice at the same site. Six of eight (75%) tautog originally released at the Texeco and Airplane Wrecks were recaptured away from these sites. Of these six tautog, three moved to the Coral Lump and Ridged Bottom and three moved to sites located 26.9 to 43.2 km away in lower Chesapeake Bay. In contrast, 32 tautog tagged and released at the Coral Lump and Ridged Bottom sites were recaptured, all but two (which moved from the Ridged Bottom to the Coral Lump) were recaptured where released. One additional fish moved to the Coral Lump from an artificial reef located 4 km to the northeast and within 2 km of where both tautog were recaptured by commercial fishermen in spring 1999.

Ultrasonically and conventionally tagged tautog released near Cape Charles, VA, in lower Chesapeake Bay demonstrated high site utilization at and high site affinity (returned to release sites after short emigration) for release sites. Extended residence by tautog at familiar sites during annual environmental extremes is considered more beneficial than emigration to more optimal environmental conditions because residence at familiar sites reduces the risk of not finding suitable shelter, food, mates, or of encountering predators (Olla et al., 1978). Although directed seasonal offshore movements were not observed in our study, movements between adjacent inshore locations occurred several times, including movement to adjacent locations during the periods of seasonal thermal extremes. Understanding temporal and spatial utilization of habitats is an important first step to identifying essential fish habitat and critical to evaluating and protecting fishery resources within Chesapeake Bay and elsewhere.

## Acknowledgments

This research represents part of M. D. Arendt's Master's thesis; earlier drafts of this manuscript were reviewed by J. Hoenig, J. Musick, D. Evans, and W. DuPaul. J. Jenrette of the FV *Bucaneer*, C. Machen of the RV *Langley*, G. Pongonis, and J. Olney Jr. and VIMS divers (B. Gammisch, W. Reisner, T. Chisholm, and W. Stockhausen) provided valuable field assistance. We thank Vemco, Ltd., and Physical Sciences Department, VIMS, for assistance with equipment operation and deployment. We also thank J. Jenrette and S. Speirs (for collectively tagging and releasing 300+ tautog at study sites and recapturing 5 of 8 tautog tagged ultrasonically), volunteer anglers, and the commercial fishing industry for tagging, releasing, and reporting recaptured tautog. This project was funded by the Recreational Fishing Development Fund of the Virginia Marine Resources Commission and by General Funds from VIMS.

[10] Verry, S. 1998. Automated wreck and obstruction information system, special area report (37°00N–37°30N; 076°00W–076°30W). NOAA/NOS, Hydrographic Services Branch/N/CS31, Silver Spring, MD 20910.

[11] Jenrette, J. 1998. Personal commun. Captain, FV *Buccaneer*, P.O. Box 149, Route 1108, Cape Charles, VA 23310.

## Literature cited


Adams, A. J.
1993. Dynamics of fish assemblages associated with an offshore artificial reef in the southern Mid-Atlantic Bight. M.A. thesis, School of Marine Science, College of William and Mary, Gloucester Point, VA, 97 p.

Arendt, M. D.
1999. Seasonal residence, movement, and activity of adult tautog (*Tautoga onitis*) in lower Chesapeake Bay. M.S. thesis, School of Marine Science, College of William and Mary, Gloucester Point, VA, 104 p.

Arendt, M. D., and J. A. Lucy.
2000. Recovery period and survival of ultrasonically tagged adult tautog in the lower Chesapeake Bay using automated receivers. *In* Biotelemetry 15: proceeding of the 15th international dymposium on biotelemetry, Juneau, Alaska (J. H. Eiler, D. J. Alcorn, and M. R. Neuman, eds.), p. 117–125. International Society on Biotelemetry, Wageningen, The Netherlands.

Arendt, M. D., J. A. Lucy and D. A. Evans.
In press. Diel and seasonal activity patterns of adult tautog, T*autoga onitis*, in lower Chesapeake Bay, inferred from ultrasonic telemetry. Environ. Biol. Fishes

Barrett, N. S.
1995. Short- and long-term movement patterns of six temperate reef fishes (Families Labridae and Monocanthidae). Mar. Fresh. Res. 46(5):853–860.

Bart, A. N., and R. A. Dunham.
1990. Factors affecting survival of channel catfish after surgical removal of testes. Prog. Fish. Cult. 52(4):241–246.

Bigelow, H. B., and W. C. Schroeder.
1953. Bony fishes, class Osteichthyes. Cunner tribe or wrasses. Family Labridae, tautog *Tautoga onitis* (Linnaeus). *In* Fishes of the Gulf of Maine, p. 478–484. Fish. Bull. 53.

Bradbury, C., J. M. Green, and M. Bruce-Lockhart.
1995. Home ranges of female cunner, *Tautogalabrus adspersus* (Labridae), as determined by ultrasonic telemetry. Can. J. Zool. 73(7):1268–1279.
1997. Daily and seasonal activity patterns of female cunner, *Tautogalabrus adspersus* (Labridae), in Newfoundland. Fish. Bull. 95(4): 646–652.

Briggs, P. T.
1969. The sport fisheries for tautog in the inshore waters of eastern Long Island. N.Y. Fish Game J. 16(2):238–254.
1977. Status of tautog populations at artificial reefs in New York waters and effect of fishing. N.Y. Fish Game J. 24(2):154–167

Cooper, R. A.
1966. Migration and population estimation of the tautog, *Tautoga onitis* (Linnaeus), from Rhode Island. Trans. Am. Fish. Soc. 95(3):239–247.

Eklund, A. M., and T. E. Targett.
1991. Seasonality of fish catch rates and species composition from the hard bottom trap fishery in the middle Atlantic Bight (US East Coast). Fish. Res.12(1):1–22.

Gelsleichter, J., J. A. Musick and S. Nichols.
1999. Food habits of the smooth dogfish, *Mustelus canis*, dusky shark, *Carcharhinus obscurus*, Atlantic sharpnose shark, *Rhizoprionodon terraenovae*, and the sand tiger, *Carcharias taurus*, from the northwest Atlantic Ocean. Environ. Biol. Fish. 54:205–217.

Holland, K. N., J. D. Peterson, C. G. Lowe, and B. M. Wetherbee.
1993. Movements, distribution, and growth rates of the white goatfish *Mulloides flavolineatus* in a fisheries conservation zone. Bull. Mar. Sci. 52(3):982–992.

Hostetter, E. B., and T. A. Munroe.
1993. Age, growth, and reproduction of tautog *Tautoga onitis* (Labridae: Perciformes) from coastal waters of Virginia. Fish. Bull. 91(1):45–64.

Lucy, J. A., and C. G. Barr.
1994. Experimental monitoring of Virginia's artificial reefs using fishermen's catch data. Bull. Mar. Sci., 55(2-3):524–537.

Matthews, K. R.
1992. A telemetric study of the home ranges and homing routes of lingcod *Ophidion elongatus* on shallow rocky reefs off Vancouver Island, British Columbia. Fish. Bull. 90(4):784–790.

Mattson, N. S., and T. H. Ripple.
1989. Metomidate, a better anesthetic for cod (*Gadus morhua*) in comparison with benzo-caine, MS-222, chlorobutanol, and phenoxyethanol. Aquaculture 83(1989):89–94.

Mortensen, D. G.
1990. Use of staple sutures to close surgical incisions for transmitter implants. Am. Fish. Soc. Symp. 7:380–383.

Nemetz, T. G., and J. R. MacMillan.
1988. Wound healing of incisions closed with a cynaoacrylate adhesive. Trans. Am. Fish. Soc. 117(2):190–195.

Olla, B. L., A. J. Bejda, and A. D. Martin.
1974. Daily activity, movements, feeding, and seasonal occurrence in the tautog, *Tautoga onitis*. Fish. Bull. 72(1): 27–35.

Olla, B. L., and C. Samet.
1977. Courtship and spawning behavior of the tautog, *Tautoga onitis* (Pisces: Labridae), under laboratory conditions. Fish Bull. 75(3):585–599.

Olla, B. L., C. Samet, and A. L. Studholme.
1981. Correlates between number of mates, shelter availability and reproductive behavior in the tautog *Tautoga onitis*. Mar. Biol. 62(4):239–248.

Olla, B.L., A. L. Studholme, A. J. Bejda, and C. Samet.
1978. Effect of temperature on activity and social behavior of the adult tautog *Tautoga onitis* under laboratory conditions. Mar. Biol. 45(4):369–378.
1980. Role of temperature in triggering migratory behavior of the adult tautog *Tautoga onitis* under laboratory conditions. Mar. Biol. 59(1):23–30.

Parker, R. O., Jr.
1990. Tagging studies and diver observations of fish populations on live-bottom reefs of the U.S. southeastern coast. Bull. Mar. Sci. 46(3):749–760.

Pearcy, W. G.
1992. Movements of acoustically-tagged yellowtail rockfish *Sebastes flavidus* on Heceta Bank, Oregon. Fish. Bull. 90(4):726–735.

Poppe, T. T., T. A. Mo, and K. Langdal.
1996. Surgical implantation of radio tansmitters into the abdominal cavity of brook trout and brown trout. Norsk Veterinaertidsskr. 198(4):241–244.

Prince, A. M. J., S. E. Low, T. J. Lissimore, R. E. Diewert, and S. G. Hinch.
1995. Sodium bicarbonate and acetic acid: an effective anethestic for field use. N. Am. J. Fish. Manage. 15(1):170–172.

Schramm, H. L., Jr., and D. J. Black.
1984. Anesthesia and surgical procedures for implanting radio transmitters into grass carp. Prog. Fish. Cult. 46(3): 185–190.

Stolgitis, J. A.
1970. Some aspects of the biology of the tautog, *Tautoga onitis* (Linnaeus), from the Weweantic River Estuary,

Massachussetts, 1966. M.S. thesis, Univ. Massachusetts, Amherst, MA, 48 p.

Stone, R. B., H. L. Pratt, R. O. Parker Jr., and G. E. Davis.
1979. A comparison of fish populations on an artificial and natural reef in the Florida Keys. Mar. Fish. Rev. 41(9):1–11.

Szedlmayer, S.
1997. Ultrasonic telemetry of red snapper, *Lutjanus campechanus*, at artificial reef sites in the northeast Gulf of Mexico. Copeia 1997(4):846–850.

Thoreau, X., and E. Baras.
1997. Evaluation of surgery procedures for implanting telemetry transmitters into the body cavity of tilapia *Oreochromis aureus*. Aquat. Living Resour. 10(4):207–211.

Walters, J. F., III.
1999. Diet composition and feeding habits of large striped bass (*Morone saxatilis*) in Chesapeake Bay. M.S. thesis, School of Marine Science, College of William and Mary, Gloucester Point, VA, 124 p.

Wells, H. W., and I. E. Gray.
1960. The seasonal occurrences of *Mytilus edulis* on the Carolina coast as a result of transport around Cape Hatteras. Biol. Bull. (Woods Hole) 119:550–559.

White, G. G.
1996. Reproductive biology of tautog, *Tautoga onitis*, in the lower Chesapeake Bay and coastal waters of Virignia. M.A. thesis, School of Marine Science, College of William and Mary, Gloucester Point, VA, 100 p.

Winter, J.
1996. Advances in underwater biotelemetry. *In* Fisheries techniques, 2nd ed. (B. R. Murphy and D. W. Willis, eds.), p. 555–590. Am. Fish. Soc., Bethesda, MD.

Wright, L. D., D. B. Prior, C. H. Hobbs, R. J. Byrne, J. D. Boon, L. C. Schaffner and M. O. Green.
1987. Spatial variability of bottom types in the lower Chesapeake Bay and adjoining estuaries and inner shelf. Estuar. Coast. Shelf Sci. 24(6):765–784.

Zeller, D. C.
1997. Home ranges and activity patterns of coral trout *Plectropomus leopardus* (Serranidae). Mar. Ecol. Prog. Ser. 154(1997):65–77.

# Phylogeographic analysis of mitochondrial DNA variation in Alaskan coho salmon, *Oncorhynchus kisutch*

**Anthony J. Gharrett**
**Andrew K. Gray**
Fisheries Division, School of Fisheries and Ocean Sciences
University of Alaska Fairbanks
11120 Glacier Highway
Juneau, Alaska 99801
E-mail address (for Anthony J. Gharrett): ffajg@uaf.edu

**Vladimir Brykov**
Institute of Marine Biology
Russian Academy of Science, Far East Branch
690041 Vladivostok, Russia

**Abstract**—Mitochondrial DNA (mtDNA) haplotypes of coho salmon (*Oncorhynchus kisutch*) sampled from northern Pacific Ocean and Bering Sea drainages formed two monophyletic clades between which nucleotide divergences averaged 2.95 substitutions per 1000 nucleotides. These data were obtained from restriction endonuclease digestions of PCR products that included over 97% of the mtDNA genome and resolved 16 different haplotypes in 258 fish from 13 locations. Comparisons of haplotype compositions of populations indicated that the Bering Sea drainages and one Kodiak Island population clustered separately from nine other Gulf of Alaska populations, including one from Asia. Rates of gene flow among populations estimated from haplotype frequencies (assuming an equilibrium between gene flow and random drift) were low (about one female per generation between drainages within regions) in relation to allozyme-based estimates of gene flow for other Pacific salmon species. Much of the haplotype frequency variation was within-region variation. Haplotypes from both clades occur in many extant populations, suggesting that gene flow, population movements, or recolonization followed divergence of refugial isolates. Nested clade analysis of the geographic distribution of mtDNA haplotypes indicated that coho salmon demographic history has been influenced by recent isolation by distance and that historic population fragmentation was preceded by range expansion. These observations are consistent with effects expected from Pleistocene glacial advances and retreats.

Manuscript accepted 19 March 2001.
Fish. Bull. 99:528–544 (2001).

In drainages flowing into the Gulf of Alaska and Bering Sea, coho salmon (*Oncorhynchus kisutch*) are the least numerous and their population structure the least understood of Pacific salmon species (*Oncorhynchus* spp.). Many populations spawn in late fall or winter in remote drainages that are difficult to access. Spawning populations are often small and separated widely (Sandercock, 1991). In larger rivers spawning adults may return to small, often transient, headwater streams. After emergence, fry and juveniles may move to rearing areas, usually much lower in the river (Sandercock, 1991), forming complex admixtures from spawning populations in large drainages.

Studies of allozyme variation have provided insight into the population structure of Pacific salmon species (e.g. Zhivotovsky et al., 1994, and references therein). Patterns of genetic variability often provide evidence of relationships between populations resulting from coancestry or gene flow, and genetic divergence among populations may be used for stock identification (Shaklee et al., 1999). In coho salmon, the low level of allozyme variation resolves relatively little population genetic structure (Reisenbichler and Phelps, 1987; Wehrhahn and Powell, 1987; Bartley et al., 1992; Pustavoit, 1995). This low level of allozyme variation is consistent with numerous spawning populations that have small effective sizes and low levels of gene flow, such as that of coho salmon.

Analysis of DNA variation adds dimensions of interpretation not possible with allozyme data (Avise et al., 1988; Avise, 1989). "Gene trees" for mitochondrial DNA (mtDNA) haplotypes are especially informative. The mitochondrial genome is transmitted (primarily) maternally (Gyllensten et al., 1991), and mtDNA is haploid and clonally inherited with no recombination. Consequently, mutations accumulate over time within a clonal mtDNA line or haplotype. Comparison of different haplotypes provides a basis for reconstruction of matriarchal genealogies. There is also evidence that mtDNA sequences may diverge (evolve) faster than many nuclear sequences (Brown et al., 1982). The extent of nucleotide divergence provides a temporal basis for comparing haplotype lineages. The rates of divergence of mtDNA have been roughly estimated for a number of species pairs by comparing observed nucleotide sequence divergence with fossil records that document the emergence of those species (Brown et al., 1979; Shields and Wilson, 1987 and references therein). Although applications of these molecular "clocks" are questionable when extended to species for which the fossil record is poor or missing, deductions about relative (as opposed to absolute) divergence times can be made from the extent of nucleotide change within a species. Furthermore, the geographic distri-




**Figure 1**

Sites from which coho salmon were sampled for mtDNA analysis. Squares denote samples used in both the preliminary and secondary analyses; circles denote collections used only for the secondary analysis. Locations are Hugh Smith River (1), Fish Creek, a Taku River tributary (2), Berner's River (3), Indian River (4), Ford Arm River (5), Crooked Creek (6), Little Susitna River (7), Buskin River (8), Karluk River (9), Eek River (10), Kanektok River (11), Delta Clearwater River, a Yukon River tributary (12), and Kamchatka River (13).

bution of haplotypes and their genealogical relationships can provide information about the historic demography and gene flow of a species. Templeton and colleagues (e.g. Templeton and Sing, 1993, Castelloe and Templeton, 1994; Templeton, 1998) have developed methods to examine both the shape of the "gene tree" and the geographic distribution of haplotypes, which they term "nested clade analysis of geographic distances."

The objectives of our study were to survey the geographic distribution of mtDNA variation in Alaskan coho salmon populations along the Gulf of Alaska and Bering Sea and to use that information and the mtDNA haplotype "gene tree" to deduce the nature of the historic demographic processes that influenced the contemporary geographic distribution of coho salmon.

## Materials and methods

Coho salmon were sampled from 12 drainages in Alaska and one in Asia (Fig. 1). Samples of heart tissue from each specimen were preserved in 95% ethanol or a solution of 20% dimethyl sulfoxide (DMSO) and 0.25M ethylenediaminetetraacetic acid (EDTA) at pH 8, saturated with NaCl (Seutin et al., 1991).

Total genomic DNA was isolated by phenol-chloroform extraction (Wallace, 1987) or with Puregene DNA™ isolation kits (Gentra Systems Inc., Minneapolis, MN). Sequences were PCR-amplified using primers that targeted seven regions of the mtDNA genome in pieces that range from about 2115 to 2689 base pairs (bp) (Fig. 2, Table 1). The regions were designated ND3/ND4 (including genes for the NADH dehydrogenase-3 subunit and NADH dehydrogenase-4L and -4 subunit genes), ND5/ND6 (including genes for the NADH dehydrogenase-5 and -6 subunits), Cytb/D-loop (including the cytochrome *b* gene and the control region), 12S/16S (including 12S rRNA gene and most of the 16S rRNA gene), ND1/ND2 (including the NADH dehydrogenase-1 and NADH dehydrogenase-2 subunit genes), COI/COII (including most of the cytochrome oxidase I subunit gene and the cytochrome oxidase II subunit gene), and A8/COIII (including genes for the ATPase-8 and -6 subunits and the cytochrome oxidase III subunit gene). The seven mtDNA regions were amplified by denaturation at 94°C for 5 min, followed by 30 cycles of 1 min at 94°C, 1 min at 55°C, and 3 min at 72°C [0.2 mM of each dNTP, 0.2μ M of each primer, 2 mM $MgCl_2$, 50 mM KCl, and 10 mM Tris-HCl, pH 8.3 with 1 unit of Taq polymerase (Perkin Elmer, Norwalk, CT) in a 50-μL reaction], except that amplifications of regions A8/COIII and ND3/ND4 required 3 mM instead of 2 mM $MgCl_2$.

Subsamples of PCR products of each mtDNA region were digested with each of 12 restriction enzymes. The endonucleases recognized six bases (*Ase* I), multiple six-base sites (*Ava* I, *Hin*d II, *Sty* I), multiple 5 base sites (*Bst*N I), and four bases (*Bst*U I, *Cfo* I, *Dde* I, *Hin*f I, *Mbo* I, *Msp* I, *Rsa* I). Digestion reactions were carried out under conditions recommended by the manufacturers. The resulting frag-




**Figure 2**

The gene map of linearized mtDNA and fragments that can be PCR amplified in coho salmon. The map is based on the sequence of *O. mykiss* mtDNA (Zardoya et al., 1995). More precise details of the range spanned by each primer pair are provided in Table 1.

**Table 1**

Primers for PCR amplification of coho salmon mitochondrial DNA (mtDNA), location in relation to *O. mykiss* (Zardoya et al., 1995), PCR fragment sizes, and fragment overlaps (bp).

| Region | Sequence | *O. mykiss* locations | Fragment size | Overlap | Source |
|---|---|---|---|---|---|
| 12S/16S | 5′ AATTCAGCAGTGATAAACATT 3′ | 1234–1254 | | | 1 |
| | 5′ AGATAGAAACTGACCTGGATT 3′ | 3615–3635 | 2402 | | 1 |
| | | | | 121 | |
| ND1/ND2 | 5′ ACCTCGATGTTGGATCAGG 3′ | 3515–3533 | | | 1 |
| | 5′ ATTAAAGTGNTTGA(T/G)TTGCATTC 3′ | 6181–6203 | 2689 | | 1, 5 |
| | | | | –430 | |
| COI/COII | 5′TAATCGTCACAGCCCATGCCTTCGT 3′ | 6634–6658 | | | 2 |
| | 5′ GGTCAGTTTCAGGGTTCAGGTTTAGC 3′ | 9079–9104 | 2471 | | 2 |
| | | | | 166 | |
| A8/COIII | 5′ CTAGTGACATGCCCCAACTCAACC 3′ | 8939–8962 | | | 2, 3 |
| | 5′ TCATAAGGCGGTCATGGACTTAAACC 3′ | 11028–11053 | 2115 | | 2, 3 |
| | | | | 480 | |
| ND3/ND4 | 5′ TTACGCGTATAAGTGACTTCCAA 3′ | 10574–10596 | | | 2, 3 |
| | 5′ TTTTGGTTCCTAAGACCAATGGAT 3′ | 12881–12904 | 2331 | | 2, 3 |
| | | | | 32 | |
| ND5/ND6 | 5′ AACAGCTCATCCATTGGTCTTAGG 3′ | 12873–12896 | | | 2, 4, 5 |
| | 5′ TTACAACGATGGTTTTTCATGTCA 3′ | 15319–15342 | 2470 | | 2, 4, 5 |
| | | | | 19 | |
| Cytb/D–loop | 5′ TGAA(G/A)ACCACCGTTGTTATTCAA 3′ | 15324–15347 | | | 2, 4, 5 |
| | 5′ TAGGGCCTCTCGTATAACCG 3′ | 1321–1340 | 2659 | | 2, 4, 5 |
| | | | | 107 | |
| 12S/16S | | | | | |

[1] Consensus from Anderson et al. (1981); Anderson et al. (1982); Bibb et al. (1981); Roe et al. (1985); Chang et al. (1994).
[2] Unpublished mtDNA sequence data from our lab.
[3] Thomas and Beckenbach (1989).
[4] Cronin et al. (1993).
[5] Carney et al. (1997).

ments were separated by electrophoresis through a 1.5% agarose gel (a mixture composed of one part Ultra Pure™ agarose [BRL Gibco, Grand, NY] and two parts Synergel™ [Diversified Biotech Inc., Boston, MA]) in 0.5×TBE buffer (TBE is 90 mM Tris-boric acid, and 2 mM EDTA, pH 7.5). DNA in the gel was stained with ethidium bromide and photographed on an ultraviolet light transilluminator. Digests that produced fragments too small for detection in agarose/Synergel™ gels were resolved in 12% polyacrylamide (29:1 acrylamide:bisacrylamide; 1×TBE) gels. DNA fragments separated in polyacrylamide gels were stained with SYBR Green 1 Nucleic Acid Stain™ (Molecular probes, Eugene, OR), which is more sensitive than ethidium bromide. Either a 1-kilobase (kb) ladder or *Hae* III digested $\phi\chi^{174}$ RF phage DNA (BRL Gibco, Grand, NY) was used as a molecular weight reference for estimating restriction fragment sizes.

Restriction sites were inferred from fragment patterns that could be related to each other by the gain or loss of a single site. Composite haplotypes were constructed from restriction fragment patterns of all restriction enzymes across all mtDNA PCR regions. Using the rules of Castelloe and Templeton (1994) to resolve ambiguities, we constructed the single most probable parsimonious tree depicting restriction site changes between haplotypes. Using REAP (McElroy et al., 1990), we estimated haplotype (nucleon) and nucleotide diversities within populations (Nei, 1987) as well as average nucleotide divergences between populations. Nucleotide divergence between populations takes into account both the haplotype frequency differences between populations and the nucleotide divergences between haplotypes (Nei and Tajima, 1983; Nei, 1987; Nei and Miller, 1990). Homogeneity of haplotype diversities among populations was tested by using the Monte-Carlo simulation in REAP (McElroy et al., 1990)(10,000 iterations; Hedges, 1992) to establish probability levels for goodness-of-fit statistics (Roff and Bentzen, 1989).

Populations were clustered from pair-wise nucleotide divergences by using the Fitch and Margoliash (1967) least-squares method (FITCH in PHYLIP; Felsenstein, 1995). For comparisons between populations, the precision of estimates of nucleotide divergence depends on sample size. Therefore, stability of the topology was examined by bootstrapping (2000 iterations; Hedges, 1992) over individuals within each collection. A consensus tree (CONSENSE in PHYLIP; Felsenstein, 1995) that shows the stability of the topology, but not the branch lengths, was generated from the set of bootstrapped trees.

The hierarchical structure of the expanded set of coho samples was analyzed by analysis of molecular variance (AMOVA; Excoffier et al., 1992) with Arlequin (Schneider et al., 1997). Collections were grouped geographically into four regions: Southeast Alaska, Southcentral Alaska, Bering Sea, and Asia. With appropriate choices of divergence matrices, the analysis can examine the structure from haplotype frequencies (e.g. Weir, 1996) or from nucleotide diversities based on paths between haplotypes traced through a haplotype tree (Excoffier et al., 1992). Significance ($P_{MC}$) of $\Phi$-statistics (Excoffier et al., 1992) was estimated from distributions of the statistics generated by 17,000 permutations (Hedges, 1992) at the appropriate level of hierarchy.

Nested clade analysis of geographical distributions of haplotypes and subclades (e.g. Templeton and Sing, 1993; Castelloe and Templeton, 1994; Templeton, 1998) was conducted with GEODIS 2.0 (Posada et al., 2000).

## Results

Our general approach was to conduct a broad preliminary survey to obtain genome-wide information for mtDNA restriction site variation. Subsequently, we focused on the variable restriction sites and examined larger sample sizes and additional populations. From those results we constructed a fine-scale mtDNA "gene tree" and analyzed the geographic distributions of mtDNA lineages to deduce the nature of the historical demographic changes that influenced present-day population structure. The approach also allowed us to determine the effects on estimates of molecular parameters that occur when analyses focus on variable restriction sites.

### Diversities of coho salmon mtDNA

Ten coho salmon from each of seven drainages (Fig. 1) were analyzed to survey broadly the species' mtDNA variability using 12 restriction endonucleases (Appendix 1). The total number of restriction sites inferred from restriction fragment patterns was 298 (an average of 291.28 per haplotype), which corresponds to 1284 nucleotides (an average of 1254.80), or a maximum of 7.73% (an average of 7.56%) of the coho salmon mtDNA genome (Table 2). Sixteen sites (1.2% of the total) were variable. Individually, the regions averaged between 29 and 57 restriction sites, which correspond to a maximum of between 5.58% and 9.57% of the nucleotides in a region. Although the amplified regions had some overlaps (Table 1), no variable sites were observed in the areas of overlap; and no invariant sites were shared between regions, except possibly *Dde* I sites in the 408-bp overlap between A8/COIII and ND3/ND4 (Table 1). Because of the large total number of sites examined, a few overlapping *Dde* I sites would cause only a slight decrease in nucleotide diversity estimates and have little effect on nucleotide divergence estimates.

Restriction site variation was observed in five of the seven PCR-amplified mtDNA regions for the 12 restriction endonucleases used. Between zero and five variable sites were observed per region. No variation was detected in the A8/COIII and ND3/ND4 regions. The largest number of variable sites (5) and the highest level of nucleotide diversity (5.99 substitutions per 1000 base pairs) were observed in the ND5/ND6 region (Table 2).

Because there is no recombination between heterologous mtDNA molecules, the composite haplotype is the appropriate unit to consider in genetic analysis (e.g. Avise, 1989). Our preliminary survey discovered 11 haplotypes (Table 3). As a whole, the sample of 70 fish had a haplotype diversity of 0.803 and a nucleotide diversity of 1.70 substitutions per 1000 base pairs. Haplotype diversities within

**Table 2**

Number and variability of restriction sites observed in each of the seven mtDNA regions we examined using 12 restriction endonucleases in our preliminary survey.

| Region | Fragment size | Mean number of sites | Mean number of nucleotides | % coverage | Number of variable sites | Number of haplotypes | Haplotype diversity (±SE) | Nucleotide diversity (per 1000) |
|---|---|---|---|---|---|---|---|---|
| 12S/16S | 2402 | 53.5 | 226.7 | 9.44 | 3 | 4 | 0.485 ±0.042 | 2.15 |
| ND1/ND2 | 2689 | 41.5 | 184.7 | 6.87 | 3 | 4 | 0.485 ±0.042 | 2.65 |
| COI/COII | 2471 | 39.7 | 168 | 6.8 | 3 | 3 | 0.057 ±0.038 | 0.25 |
| A8/COIII | 2115 | 29 | 126.7 | 5.99 | 0 | 1 | 0 | 0 |
| ND3/ND4 | 2331 | 31 | 130 | 5.58 | 0 | 1 | 0 | 0 |
| ND5/ND6 | 2470 | 36.5 | 157.2 | 6.36 | 5 | 4 | 0.470 ±0.040 | 5.99 |
| Cytb/D-loop | 2659 | 57 | 248.7 | 9.35 | 2 | 3 | 0.670 ±0.014 | 1.87 |
| Total | 16,642 | 291.28 | 1254.8 | 7.54 | 16 | 11 | 0.803 ±0.024 | 1.70 |

**Table 3**

Observed numbers of each mtDNA haplotype, haplotype diversity, and nucleotide diversity (substitutions per 1000 bp) within collections of coho salmon examined in a preliminary survey of North Pacific Ocean coho salmon (Nei and Tajima, 1983; Nei ,1987). Standard errors are in parentheses. Homogeneity of haplotype frequencies ($P_{MC} < 10^{-4}$) was tested with using Monte-Carlo simulation based on 10,000 resampling iterations to estimate probability (Roff and Bentzen, 1989).

| | | A–D cluster | | | | | | E–H cluster | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Haplotype | | | | | | Haplotype | | | | | | |
| Collection | $n$ | A | A′ | B | C | C′ | D | E | E′ | F | G | H | Haplotype diversity | Nucleotide diversity |
| Hugh Smith River | 10 | 3 | 0 | 0 | 4 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0.78 | 1.31 |
| Fish Creek (Taku River) | 10 | 7 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0.53 | 0.88 |
| Ford Arm River | 10 | 2 | 0 | 4 | 1 | 0 | 0 | 2 | 0 | 0 | 1 | 0 | 0.82 | 1.65 |
| Crooked Creek | 10 | 5 | 0 | 0 | 4 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0.64 | 0.78 |
| Eek River | 10 | 0 | 0 | 0 | 4 | 0 | 0 | 4 | 0 | 0 | 0 | 2 | 0.71 | 1.87 |
| Delta Clearwater River | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0.00 | 0.00 |
| Kamchatka River | 10 | 5 | 0 | 0 | 4 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0.06 | 0.94 |
| Total | 70 | 22 | 1 | 4 | 17 | 1 | 1 | 9 | 1 | 1 | 1 | 12 | 0.80 (±0.024) | 1.70 (±0.000) |
| Average | | | | | | | | | | | | | 0.59 (±0.11) | 1.06 (±0.24) |

collections from individual drainages ranged from 0.00 to 0.82 and nucleotide diversities ranged from 0.00 to 1.87 substitutions per 1000 bp. The distribution of haplotypes among collections was highly heterogeneous, which is noteworthy given the small sample sizes ($n$=10) (Table 3).

Phylogenetically, there were two clusters of haplotypes (haplotypes A–D and haplotypes E–H) (Fig. 3). Haplotypes of the two clusters differed by five or more restriction sites, and the average nucleotide divergence between individual haplotypes in the two clusters averaged 2.72 nucleotide substitutions per 1000 nucleotides, as compared to an average nucleotide diversity within each cluster of 0.87. The cluster of A–D accounted for the majority of fish, but haplotypes of both clusters were observed in collections from all drainages, except for fish from the Delta Clearwater River, which had only haplotype H. Bootstrap estimates of nucleotide divergence between clusters and average nucleotide diversity within clusters, estimated from the entire sample of 70 fish, were 2.98 ±0.06 and 0.35 ±0.05 (substitutions per 1000 nucleotides), respectively.

## Expanded coho mtDNA survey based on variable sites

We increased the number of populations surveyed to 13 and increased sample sizes to 20, except for the Kam-




**Figure 3**

Haplotype tree for coho salmon mtDNA restriction site variation. Haplotypes A–H, including A′, C′, and E′, were observed in the preliminary survey (70 fish); haplotypes I–O were observed in the subsequent survey (188 additional fish); and haplotype P was detected in digests conducted to resolve positions on the tree of the latter haplotypes. Arrows indicate the direction of site gain. Ambiguities in the topology were resolved (Castelloe and Templeton, 1994) and the nested structure was constructed (Templeton and Sing, 1993). Haplotypes A′, C′, and E′ were pooled with A, C, and E, respectively, for the nested clade analysis. The results of the analysis are provided in Table 5.

chatka River ($n$=17) and Delta Clearwater River ($n$=21) populations. To make analysis of larger numbers of samples practical, we focused on restriction sites in five mtDNA regions that defined the most abundant eight (A–H) of the 11 haplotypes. Haplotypes A′, C′, and E′, each of which was represented by a single individual, were eliminated. In this survey, we analyzed site variation for the following PCR region by restriction endonuclease combinations:

| 12S/16S rRNA | Cyt b/D-loop | ND5/ND6 | ND1/ND2 | ND3/ND4 |
|---|---|---|---|---|
| *Cfo* I | *Bst*N I | *Ava* I | *Dde* I | *Cfo* I |
| *Dde* I | | *Dde* I | *Sty* I | |

This survey detected 62 restriction sites (56.47 on average in each haplotype), which correspond to 262.67 nucleotides (238.22 on average). The proportion of the genome screened was a maximum of 1.58% (1.43% on average). To completely resolve the placement of haplotypes I, J, K, and L in the "gene tree," we digested their ND5/ND6 regions with *Mbo* I and their COI/COII regions with *Dde* I. An additional haplotype (P) was resolved. In total, the expanded survey resolved three additional haplotypes (I, J, and P) within the A–D clade and five additional haplotypes (K, L, M, N, and O) within the E–H clade (Table 4, Appendix 2, and Fig. 3). Haplotypes of both clusters appeared in most drainages. It is notable that four drainages included haplotypes of only a single clade: Delta Clearwater had 20 of haplotype H and one of the related haplotype O; Indian River had 19 of haplotype A and one of haplotype C; Berners River had five of hapotype A, 11 of haplotype C, one of haplotype I, two of haplotype J, and one of haplotype P; and the Little Susitna collection had four A haplotypes, 15 C haplotypes, and one I haplotype. Within drainages, haplotype diversities ranged from 0.10 to 0.73 and nucleotide diversities ranged from 0.22 to 9.07 substitutions per 1000 bp. The 13 collections exhibited strong heterogeneity ($P<10^{-4}$, Table 4).

In a Fitch-Margoliash phenogram that estimates relationships among drainages based on haplotype frequencies (Fig. 4), Delta Clearwater River differed strongly from the other collections, and the collections from systems that drain into the Bering Sea and from Karluk Lake on Kodiak Island, clustered separately. The remaining col-

**Table 4**

Observed numbers of each mtDNA haplotype, haplotype diversity, and nucleotide diversity (substitutions per 1000 bp) within collections of North Pacific coho salmon screened for variable sites detected in a preliminary survey (Table 3)(Nei and Tajima, 1983; Nei, 1987). Standard errors are in parentheses. Homogeneity of haplotype frequencies ($P_{MC} < 10^{-4}$) was tested by using Monte-Carlo simulation based on 10,000 resampling iterations to estimate probability (Roff and Bentzen, 1989).

| | | Haplotype | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Collection | $n$ | A | B | C | D | E | F | G | H | I | J | K | L | M | N | O | P | Haplotype diversity | Nucleotide diversity |
| Hugh Smith River | 20 | 6 | 0 | 12 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.57 | 3.64 |
| Fish Creek (Taku River) | 20 | 11 | 0 | 3 | 2 | 0 | 2 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0.68 | 5.68 |
| Berners River | 20 | 5 | 0 | 11 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 1 | 0.64 | 2.40 |
| Indian River | 20 | 19 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.10 | 0.22 |
| Ford Arm River | 20 | 10 | 6 | 1 | 0 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.68 | 4.51 |
| Crooked Creek | 20 | 9 | 0 | 10 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.57 | 2.50 |
| Little Susitna River | 20 | 4 | 0 | 15 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.42 | 1.13 |
| Buskin River | 20 | 5 | 0 | 12 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.58 | 4.69 |
| Karluk River | 20 | 1 | 0 | 9 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 4 | 0 | 0 | 0.73 | 9.07 |
| Eek River | 20 | 1 | 0 | 7 | 0 | 9 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.68 | 8.38 |
| Kanektok River | 20 | 5 | 0 | 8 | 0 | 5 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0.75 | 7.91 |
| Delta Clearwater River | 21 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0.1 | 0.2 |
| Kamchatka River | 17 | 9 | 0 | 7 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0.58 | 2.7 |
| Total | 258 | 85 | 6 | 96 | 2 | 28 | 2 | 1 | 24 | 2 | 3 | 1 | 1 | 1 | 4 | 1 | 1 | 0.734 (±0.016) | 6.73 (±0.00) |
| Average | | | | | | | | | | | | | | | | | | 0.54 (±0.06) | 4.08 (±0.83) |

lections spanned the southern portion of the geographic range from southern Southeast Alaska to the Kamchatka Peninsula. Within that set of collections, the two coastal Southeast Alaskan collections (Ford Arm and Indian River) appeared to form a weak cluster, but there was no obvious structure among the remaining collections.

We conducted AMOVA analyses reflecting a geographical hierarchy: Southeast Alaska, Southcentral Alaska, western and interior Alaska (Bering Sea), and Asia to examine the geographic basis of variation. Although the Karluk River collection resembled Bering Sea collections more than other northern Gulf of Alaska collections, we included it with the Southcentral Alaska group to maintain the geographic basis of the analysis. Analyzing the data based on haplotype frequencies (analogous to allelic differences in analysis of variance described by Weir [1996]) revealed highly significant divergence among collections ($\Phi_{ST}$=0.291, $P_{MC}$<0.0001), most of which is attributable to average divergence among drainages within a region ($\Phi_{SC}$=0.227, $P_{MC}\leq$0.0001), rather than differences between regions ($\Phi_{CT}$=0.083, $P_{MC}$=0.094). Incorporating relationships between the haplotypes into the analysis increased the proportion of the total divergence observed among drainages ($\Phi_{ST}$= 0.449, $P_{MC}\leq$0.0001) and among drainages within regions ($\Phi_{SC}$=0.273 , $P_{MC}$<0.0001). The estimate of the proportion of divergence among regions also increased ($\Phi_{CT}$=0.242, $P_{MC}$=0.083). Estimates of long-term gene flow [$N_{e(f)}m=(\Phi_{XY}^{-1}-1)/2$] from $\Phi_{SC}$ and $\Phi_{CT}$ based on haplotype relationships were about 1 female per generation between collections within regions and about two females per generation between regions. Such estimates assume that an equilibrium between gene flow and random drift exists.

The nested clade analysis (Fig. 3, Table 5) collapses the "gene tree" from the periphery and analyzes each subclade for significantly small or large geographical distributions of the components as compared with the subclade as a whole. The first two levels of nesting are 1-step clades (nesting 0-step haplotypes) and 2-step clades (nesting 1-step clades). The significance in their geographic distributions were consistent with restricted gene flow with isolation by distance for haplotypes of the A–D (plus I, J, and P) subclade and past fragmentation for haplotypes of the E–H (less G plus M, N, and O) subclade. The 3-step clade (nested 2-step clades) that includes all of the E–H (plus L, M, N, and O) haplotypes also indicates past fragmentation. The most interior level of nesting, which contrasts the A group of haplotypes with the E group, assuming that the A group is interior (ancestral) based on the interpretations of Castelloe and Templeton (1994), is consistent with contiguous range expansion. We reanalyzed the data without the Delta Clearwater population, the Kamchatka population, and without either. These two geographically distinct populations did not alter the overall interpretation, but the Delta Clearwater population was responsible for significance of clade 2–4 and the Kamchatka population was responsible for the significance of clade 2–2.




**Figure 4**

A Fitch-Margoliash least-squares phenogram constructed from nucleotide divergences among coho salmon collections and a consensus tree (inset) derived from 2000 iterations (Hedges, 1992) of bootstrap resampling from the individuals within each collection. Geographic regions for the samples are Russia (R), Southcentral Alaska (SC), Kodiak Island (KI), and Southeast Alaska (SE), and Bering Sea (BS), The consensus tree is unscaled and describes the confidence of the topology depicted in the phenogram.

## Discussion

### Coho salmon mtDNA variation

We examined the coho salmon mtDNA genome using restriction analysis of seven PCR products and detected fragments as small as 25 bp, which made it possible to sample nearly all (over 97%). of the mitochondrial genome. We are unaware of other fish species that have been as thoroughly surveyed. Our estimates of species-level parameters of molecular evolution should be quite robust in comparisons with those for other taxa, including fish.

Our results (Table 2) suggest that mtDNA variation is not evenly distributed throughout the genome and that focusing analysis on variable sites overestimates nucleotide diversity estimates (see Tables 3 and 4). Many previous studies of Pacific salmon species have limited the portion of the mtDNA genome surveyed (e.g. chum salmon [Cronin et al.; 1993; Park et al., 1993; Seeb and Crane, 1999]; sockeye salmon [Bickham et al., 1995; Taylor et al. 1996]; chinook [Cronin et al., 1993]). Other studies have included a limited geographic range of samples and have restricted the portion of the mtDNA genome studied (e.g. sockeye [Burger et al., 1997]; chinook [Adams et al., 1994]) or surveyed selected variable sites (e.g. pink salmon [Brykov et al., 1996; Seeb et al., 1999]; chum salmon [Scribner et al., 1998]; sockeye [Taylor et al., 1997]). Consequently, no previous studies have produced results that are appropriate for a broad comparison with our results.

We evaluated our results in the context of other fish species by plotting the effective number of haplotypes observed (the number of haplotypes which, if equally abundant, would result in the observed haplotype diversity, $n_e$, against the average nucleotide divergences between haplotypes for piscine species. For these comparisons, we used only data that were derived from 20 or more individuals and that surveyed the entire mtDNA genome (Fig. 5). Only two studies of Pacific salmon met those criteria; and our estimates of nucleotide diversity (0.4–4.5 per 1000 nucleotides) were generally lower than estimates for chinook salmon (6 haplotypes: 1.3–8.1; Wilson et al., 1987) and chum salmon (2 haplotypes: 2.4; Thomas et al., 1986).

Average nucleotide divergences between coho salmon haplotypes are quite low in relation to those of most other species studied; whereas, the effective number of haplotypes is near the median. The effective number of (selectively neutral) haplotypes ($n_e$) is monotonically related to the product of effective population number (of females

**Table 5**

Geographical nested clade analysis (Templeton, 1998) for Alaskan coho salmon mtDNA haplotpes based on the haplotype network in Figure 3 using GEODIS 2.0 (Posada et al., 2000). Only clades that exhibit significant results are shown. The analysis examines the geographical spread ($D_C$) of tip (T) and interior (I) subclades (Castelloe and Templeton, 1994), their relative spread within the clade ($D_N$), and contrasts their distrubutions (I–T) identifying distances that are significantly small (S), significantly large (L), or not significant (NS). Interpretations are based on Templeton's (1998) inference key.

| 1-step clades | | | | | 2-step clades | | | | | 3-step clades | | | | | 4-step clades | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Clade | Sub-clades | Tip/ Interior | $D_C$ | $D_N$ | Clade | Sub-clades | Tip/ Interior | $D_C$ | $D_N$ | Clade | Sub-clades | Tip/ Interior | $D_C$ | $D_N$ | Clade | Sub-clades | Tip/ Interior | $D_C$ | $D_N$ |
| 1-3 | A | I | NS | NS | 2-1 | 1-2 | I | NS | NS | 3-2 | 2-3 | I | NS | L | 4-1 | 3-1 | I | S | S |
| | B | T | S | S | | 1-1 | T | S | NS | | 2-4 | T | S | S | | 3-2 | T | NS | L |
| | | I–T contrast | L | L | | | I–T contrast | L | NS | | | I–T contrast | NS | L | | | I–T contrast | NS | S |
| | Isolation by distance. | | | | | Isolation by distance. | | | | | Past fragmentation | | | | | Contiguous range expansion | | | |
| 1-4 | E | I | NS | NS | 2-2 | 1-3 | I | L | L | | | | | | | | | | |
| | F | T | NS | L | | I | T | NS | NS | | | | | | | | | | |
| | M | T | NS | NS | | | I–T contrast | NS | NS | | | | | | | | | | |
| | N | T | S | S | | Islation by distance. | | | | | | | | | | | | | |
| | | I–T contrast | L | NS | | | | | | | | | | | | | | | |
| | Past fragmentation | | | | 2-4 | 1-4 | I | NS | L | | | | | | | | | | |
| | | | | | | 1–5 | T | S | NS | | | | | | | | | | |
| | | | | | | K | T | NS | NS | | | | | | | | | | |
| | | | | | | | I–T contrast | L | NS | | | | | | | | | | |
| | | | | | | Past fragmentation | | | | | | | | | | | | | |

in the case of mtDNA – $N_{e(f)}$) and mutation rate (μ) (Kimura and Crow, 1964; Ohta and Kimura, 1973). However, for μ that are relatively constant among species, $n_c$ orders the relative size of historic effective sizes of the species (Avise et al., 1988). Estimates of $n_c$ may be influenced by the number of individuals analyzed, the scope of sampling with respect to both geographic range, and the number of sites detected in the mtDNA genome.

The average nucleotide sequence divergence between haplotype pairs presents a more complex picture. Accumulation of nucleotide differences between haplotypes requires time. If the mutation rates (μ) are reasonably similar (although there is debate about the degree of similarity [e.g. Avise et al., 1988; Gibbons, 1997]), the extent of haplotype divergence should order the species according to the time that has elapsed since a major species bottleneck. Such ordering is a consequence of lineage sorting within isolates, which tends toward a set of related (monophyletic) haplotypes but fosters divergence among isolates. However, as a result of lineage sorting, species composed of isolates are likely to exhibit higher average divergence than species broadly connected by gene flow (Neigel and Avise, 1986). The small nucleotide divergence among haplotypes in coho salmon suggests that extant coho salmon share a relatively recent common female ancestor.

Two distinct clusters of haplotypes account for most of the divergence observed among coho salmon. The clusters are separated by an average of 2.72 substitutions per 1000 nucleotides, but divergence among haplotypes within each cluster average about 0.87 substitutions per thousand nucleotides.

## Expanded coho mtDNA survey from variable sites

By concentrating our effort on variable sites, we evaluated the variation present in a larger number of fish and from additional populations. The two clusters of haplotypes observed in the preliminary survey persisted, each augmented by several new haplotypes. The average divergence between haplotypes within each cluster was 4.95 substitutions per 1000 nucleotides and the clusters averaged 10.28 substitutions per 1000 bp apart. By focussing on variable restriction sites,

estimates of nucleotide divergence were inflated more than threefold. However, the data remain useful for examining intraspecific structure, and strong heterogeneity among collections indicated structure among populations or at higher levels of population hierarchy.

The unrooted tree (Fig. 4) depicting relationships among collections shows that the Bering Sea and the Karluk River collections cluster separately from the other collections. The cluster-based structure of the "gene tree" accentuates the divergence between geographic regions because the highest abundance of E-cluster haplotypes occurs in the Bering Sea (and Karluk) collections. In fact, the Delta Clearwater collection is fixed for E-cluster haplotypes. The AMOVA analyses revealed a population structure that appears stronger within regions than among regions. Both estimates of gene flow (one to two females per generation) are low in relation to allozyme-based estimates of other Pacific salmon species (i.e. an exchange of 6.6 to 16.4 individuals per generation for pink salmon [McGregor, 1983; Beacham et al., 1985; Noll et al., 2001]; 6.3 to 9.0 for chum salmon [Kondzela et al., 1994; Phelps et al., 1994; Wilmot et al., 1994; Winans et al.. 1994]; 1.3 to 7.1 for sockeye salmon [Wood et al., 1994; Varnavskaya et al., 1994]; and 1.2 to 4.0 for chinook salmon [Gharrett et al., 1987; Utter et al., 1989; Bartley and Gall, 1990]) but consistent with estimates from coho salmon allozyme data (Wehrhan and Powell, 1987; Bartley et al., 1992).

It is essential to keep in mind that such gene-flow estimates assume an equilibrium between gene flow and random drift. The distribution of coho salmon haplotypes suggests that a broad equilibrium may not exist. For example, the Delta Clearwater River population was nearly fixed for haplotype H, an E-cluster haplotype found only in the Bering Sea populations. Its haplotype composition reflects its geographic isolation from other coho populations. Also, the haplotype distribution from the Kamchatka River population was surprising because it is so similar to Southeast Alaskan populations, but differed from the geographically closer Bering Sea populations. This makes sense if the genetic structure of extant coho salmon populations is strongly influenced by historic events, such as historic random drift, colonization from eastern populations, or survival in an different glacial refugia, and that an equilibrium between gene flow and random drift has not yet been reached.




**Figure 5**

Comparison of the effective number of haplotypes ($n_c$) and the average nucleotide divergence between haplotypes within a species for 25 species of fish estimated from haplotype diversity ($h$). The number of effective haplotypes is monotonically related to $N_e\mu$ ($\mu$ is the mutation rate); larger average nucleotide divergences require longer times and reduced gene flow to develop. The following species were plotted: A = red grouper (Richardson and Gold, 1993, 1997); B = white marlin (Graves and McDowell, 1995); C = hardhead catfish (Avise et al., 1987, 1989); D = American eel (Avise et al., 1986, 1989); E = oyster toadfish (Avise et al., 1987, 1989); F = American shad (Bentzen et al., 1989); G = redear sunfish (Avise et al., 1987, 1988, 1989, 1992); H = *Sicyopterus stimpsoni* (Zink et al., 1996); I = lake whitefish (Bernatchez and Dodson, 1991); J = red snapper (Gold et al., 1994, 1997); K = coho salmon (Carney et al., 1997); L = black drum (Richardson and Gold, 1993; Gold et al., 1994); M = spotted sunfish (Avise et al., 1987, 1989, 1992; Avise, 1992); N = coho salmon (this study); O = NW Atlantic capelin (Dodson et al., 1991); P = bowfin (Avise et al., 1987, 1989, 1992; Avise, 1992); Q = striped marlin (Graves and McDowell, 1994; Graves and McDowell, 1995); R = *Stenogobius hawaiiensis* (Zink et al., 1996); S = *Lentipes concolor* (Zink et al., 1996); T = greater amberjack (Richardson and Gold, 1993); U = *Awaous guamensis* (Zink et al., 1996); V = Atlantic herring (Kornfield and Bogdanowicz, 1987; Richardson and Gold, 1993); W = largemouth bass (Nedbal and Phillipp, 1994); X = red drum (Gold and Richardson, 1991; Gold et al., 1993); Y = warmouth (Avise et al., 1987, 1989, 1992; Avise, 1992); Z = NE Atlantic capelin (Dodson et al., 1991). Estimates for coho salmon in this study (N) and our previous study (K) are circled.

## Synthesis

The data obtained from restriction site analysis provide a present day "snapshot" of coho salmon mtDNA variation and have two levels of resolution. One level is the geographic distribution of haplotypes and the haplotype compositions of the populations sampled. The other level is the pattern and extent of divergence among the haplotypes. The mtDNA haplotype compositions of populations indicate that coho salmon, at least in their Alaskan range, generally have lower gene flow than other Pacific salmon species, although the comparisons assume equilibria between gene flow and random drift that may not yet have been reached. Coho salmon exhibit divergence among populations within regions, but generally not fixed differences. Many of the haplotypes were found in populations throughout the range examined, which suggests

that at some time the haplotypes were disseminated either through gene flow or colonization. Our data, are generally consistent with the known distributions and abundances of North Pacific coho salmon, that is, coho salmon is a species composed of many small spawning populations that have low levels of gene flow. The low geneflow rates estimated within several geographic ranges indicate that a finer scale study of coho salmon population structure is warranted.

Superimposed on this "snapshot" of population structure is the mtDNA "gene tree," which carries information about the demographic history of coho salmon. Nested clade analysis of the geographic distribution of haplotypes and clades of related haplotypes reveal recent isolation by distance, particularly for the geographical distribution of A-cluster haplotypes. There is also evidence of past population fragmentation in the distribution of E-cluster haplotypes, and the interior of the tree has evidence of range expansion. The topology of the tree also indicates recent demographic expansion preceded by stationary, or more likely, declining populations. Radiation of haplotypes to form a star-like pattern occurs when populations expand (Slatkin and Hudson, 1991), such as the haplotypes that surround haplotypes A and E. Whereas, in stationary or declining populations, the haplotype composition of a population eventually tends toward a single, evolutionarily related (monophyletic) set as a result of stochastic processes (Neigel and Avise, 1986), such as the A- or E-cluster.

It is likely that the demographic history of coho salmon is closely tied to Pleistocene events. During the Pleistocene Epoch, glaciers periodically advanced and retreated in regions bordering the northeastern Pacific Ocean. These advances were accompanied by a drop in sea level (exceeding 100 m), decreased sea surface temperatures, increased coverage of sea ice, and reduced marine surface water productivity (e.g. Porter, 1989; Bartlein et al., 1991; de Vernal and Pedersen, 1997). Much of the present day freshwater range was physically covered with ice (Hamilton, 1986). During the last glacial maximum, coho salmon must have been displaced from most of British Columbia and the Gulf of Alaska coast, now the center of their range. During each glacial advance, it is likely that marine and freshwater habitats of Pacific salmon populations were greatly reduced, especially for species that require freshwater rearing, such as coho salmon. Alteration and reduction of natural ranges during these advances probably resulted in the declines or extirpation of many populations. The patterns observed in the analysis of coho salmon mtDNA variation resulted from major demographic changes involving the range of drainages sampled and are consistent with the effects that would be expected as a result of glacial advances and retreats. For most of the last 500,000 years, the environment was much harsher than now, as indicated by the sea level (a measure of the amount of the world's water tied up in ice), which was more than 50 m lower than today during most of that period (Porter, 1989; Bartlein et al., 1991). Peaks of recent interglacial periods during which the environment probably approached modern conditions have occurred about every 100 thousand years, approximately 120 thousand years ago (120 ka) , 200 ka, 330 ka, and 400 ka (Porter, 1989; Petit et al., 1999).

A linkage between geologic record and molecular evolution requires application of a "molecular clock" to relate observed nucleotide divergences to a mutation rate. The calibration of mtDNA clocks is contentious. A rate of 2% nucleotide substitution per million years (Brown et al., 1982), referred to as the "standard clock," has been broadly applied. However, the rates for salmonids may be slower, and rates of 1% per million years or less (Smith, 1992) may be more appropriate.

An interpretation of our results based on the "standard clock" would be that divergence between clusters began 136 ka at the beginning of the last interglacial period and that divergence within the clusters began 43 ka, at the end of the last interglacial period. The small number of haplotypes within each cluster is consistent with a relatively small effective number of females ($N_{e(f)}$) within refugia followed by rapid expansion or the existence of several isolates within each refugium. Following the last glacial maximum, dispersion from one or more glacial refugia was followed by incomplete inter-refugial introgression. The objection to interpretations based on the "standard clock" is that the divergence rate is much faster than the rate generally accepted for salmonids. However, mtDNA clocks are calibrated from interspecies comparisons, which occur over times measured in millions of years (Thomas and Beckenbach, 1989; Martin et al., 1992; Bentzen et al., 1993; Phillips and Oakley, 1997). The mutation rate is certainly not homogeneous over the mtDNA genome, so it is possible that mutational hot spots dominate the rate in shorter time spans but are saturated and much less important in estimates over long time spans.

Although there are unquestionably strong demographic signals in the molecular evolutionary record of coho salmon, we can not match them unequivocally to specific Pleistocene events. To make such a match, several pieces of information are needed. First, the data available for calibrating a clock for the entire salmonid mtDNA genome are limited. Although there are numerous studies of particular mtDNA regions, rate differences among regions compromise their utility. Second, short-term, intraspecific clocks based on hot spots must be developed and compared with long-term, interspecific clocks. We are now examining data from other Pacific salmon species that share the northern Pacific Ocean range with coho salmon. Concordance among those results interpreted from the well-described Pleistocene environmental history may provide us with independent records of Pleistocene influences on the demography of Pacific salmon, which can be used to calibrate intraspecific mtDNA clocks.

The haplotype compositions of the populations that we studied leave us with questions about patterns of postglacial colonization and influences of the Cascadian and Beringian refugial stocks. Acquisition and analysis of samples from native populations in the Pacific Northwest should resolve questions about the composition of coho salmon in the Wisconsin Cascadian refugium. And, an intensive study of populations skirting the Alaska Peninsula and eastern Bering Sea should provide information about local colonization patterns and the composition of Beringian coho.

## Acknowledgments

We appreciate the constructive and encouraging comments made by A.T. Beckenbach on an early draft. D. Churikov and B. Finney provided valuable discussions. We thank D. Churikov, J. Gharrett, and three anonymous reviewers for reviewing the manuscript. T. Chatto, S. Kelley, D. McBride, L. Schwartz, L. Shaul, C. Skaugstad, and S. Taylor kindly provided samples. This work is a result of research sponsored in part by the Saltonstall-Kennedy Fund NA36 FD0179 and in part by the Alaska Sea Grant College Program, which is cooperatively sponsored by NOAA, Office of Sea Grant and Extramural Programs, Department of Commerce, under grant NA90AA-D-SG066 (project R/02-16), and the University of Alaska with funds appropriated by the State. The manuscript was revised while AJG was at Hokkaido University and supported, in part, by the Japan Society for Promotion of Science.

## Literature cited

Adams, N. S., W. J. Spearman, C. V. Burger, K. P. Currens, C. B. Schreck, and H. W. Li.
1994. Variation in mitochondrial DNA and allozymes discriminates early and late forms of chinook salmon (*Oncorhynchus tshawytscha*) in Kenai and Kasilof rivers, Alaska. Can. J. Fish. Aquat. Sci. 51(suppl. 1):172–181.

Anderson, S., A. T. Bankier, G. T. Barrell, M. H. DeBruijn, and A. R. Coulson.
1981. Sequence and organization of the human mitochondrial genome. Nature 290:457–465.

Anderson, S., M. H. DeBruijn, A. R. Coulson, I. C. Eperon, and F. Sanger.
1982. Complete sequence of bovine mitochondrial DNA. J. Mol. Biol. 156:683–717.

Avise, J. C.
1989. Gene trees and organismal histories: a phylogenetic approach to population biology. Evolution 43:1192–1208.
1992. Molecular population structure and the biogeographic history of a regional fauna: a case history with lessons for conservation biology. Oikos:63–76.

Avise, J. C., J. Arnold, R. M. Ball, E. Bermingham, T. Lamb, J. E. Neigel, C. A. Reeb, and N. C. Saunders.
1987. Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. Ann. Rev. Ecol. Syst. 18:489–522.

Avise, J. C., R. M. Ball, and J. Arnold.
1988. Current versus historical population sizes in vertebrate species with high gene flow: a comparison based on mitochondrial DNA lineages and inbreeding theory for neutral mutations. Mol. Biol. Evol. 5:331–344.

Avise, J. C., B. W. Bowen, and T. Lamb.
1989. DNA fingerprints from hypervariable mitochondrial genotypes. Mol. Biol. Evol. 6:258–269.

Avise, J. C., B. W. Bowen, T. Lamb, A. B. Meylan, and E. Bermingham.
1992. Mitochondrial DNA evolution at a turtle's pace: evidence for low genetic variability and reduced microevolutionary rate in the Testudines. Mol. Biol. Evol. 9:457–473.

Avise, J. C., G. S. Helfman, N. C. Saunders, and L. S. Hales.
1986. Mitochondrial DNA differentiation in North Atlantic eels: population genetic consequences of an unusual life history pattern. Proc. Nat. Acad. Sci. US. 83:4350–4354.

Bartlein, P. J., P. M. Anderson, M. E. Edwards, and P. F. McDowell.
1991. A framework for interpreting paleoclimatic variations in eastern Beringia. Quaternary International 12-1: 73–83.

Bartley, D. M., B. Bentley, P. G. Olin, and G. A. E. Gall.
1992. Population genetic structure of coho salmon (*Oncorhynchus kisutch*). Calif. Fish. Game 78:88–104.

Bartley, D. M., and G. A. E. Gall.
1990. Genetic structure and gene flow in chinook salmon populations in California. Trans. Am. Fish. Soc. 119:55–71.

Beacham, T. D., R. E. Withler, and A. P. Gould.
1985. Biochemical genetic stock identification of pink salmon (*Oncorhynchus gorbuscha*)in southern British Columbia and Puget Sound. Can. J. Fish. Aquat. Sci. 42:1474–1483.

Bentzen, P., G. C. Brown, and W. C. Leggett.
1989. Mitochondrial DNA polymorphism, population structure, and life history variation in American shad (*Alosa sapidissima*). Can. J. Fish. Aquat. Sci. 46:1446–1454.

Bentzen, P., W. C. Leggett, and G. G. Brown.
1993. Genetic relationships among the shads (*Alosa*) revealed by mitochondrial DNA analysis. J. Fish. Biol. 43:909–917.

Bernatchez, L., and J. J. Dodson.
1991. Phylogeographic structure in mitochondrial DNA of the lake whitefish (*Coregonus clupeaformis*) and its relation to Pleistocene glaciations. Evolution 45:1016–1035.

Bibb, M. J., R. A. van Etten, C. T. Wright, M. W. Walberg, and D. A. Clayton.
1981. Sequence and gene organization of mouse mitochondrial DNA. Cell 26:167–180.

Bickham, J. W., C. C. Wood, and J. C. Patton.
1995. Biogeographic implications of cytochrome b sequences and allozymes in sockeye (*Oncorhynchus nerka*). J. Hered. 86:140–144.

Brown, W. M., M. George, Jr., and A. C. Wilson.
1979. Rapid evolution of animal mitochondrial DNA. Proc. Natl. Acad. Sci. USA 76:1967–1971.

Brown, W. M., E. M. Prager, A. Wang, and A. C. Wilson.
1982. Mitochondrial DNA sequences of primates: tempo and mode of evolution. J. Mol. Evol. 18:225–239.

Brykov, V. A., N. Polyakova, L. A. Skurikhina, and A. D. Kukhlevsky.
1996. Geographical and temporal mitochondrial DNA variability in populations of pink salmon. J. Fish Biol. 48:899–909.

Burger, C. V., W. J. Spearman, and M. A. Cronin.
1997. Genetic differentiation of sockeye salmon subpopulations from a geologically young Alaskan lake system. Trans. Am. Fish. Soc. 126:926–938.

Carney, B. L., A. K. Gray, and A. J. Gharrett.
1997. Mitochondrial DNA restriction site variation within and among five populations of Alaskan coho salmon (*Oncorhynchus kisutch*). Can. J. Fish. Aquat. Sci. 54:940–949.

Castelloe, J., and A. R. Templeton.
1994. Root probabilities for intraspecific gene trees under neutral coalescent theory. Mol. Phylogenet. Evol. 3:102–113.

Chang, Y-S., F-L. Huang, and T-B. Lo.
1994. The complete nucleotide sequence and gene organization of carp (*Cyprinus carpio*) mitochondrial genome. J. Mol. Evol. 38:138–155.

Cronin, M. A., W. J. Spearman, R. L. Wilmot, J. C. Patton, and J. W. Bickham.
1993. Mitochondrial DNA variation in chinook (*Oncorhyn-*

*chus tshawytscha*) and chum salmon (*O. keta*) detected by restriction enzyme analysis of polymerase chain reaction (PCR) products. Can. J. Fish. Aquat. Sci. 50:708–715.

de Vernal, A., and T. F. Pedersen.
1997. Micropaleontology and palynology of core PAR87A-10: A 23,000 year record of paleoenvironmental changes in the Gulf of Alaska, northeast Pacific. Paleoceanography 12:821–830.

Dodson, J. J., J. E. Carscadden, L. Bernatchez, and F. Colombani.
1991. Relationship between spawning mode and phylogeny in mitochondrial DNA of North Atlantic capelin *Mallotus villosus*. Mar. Ecol. Prog. Ser. 66:103–113.

Excoffier, L., P. E. Smouse, and J. M. Quattro.
1992. Analysis of molecular inference from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. Genetics 131:479–491.

Felsenstein, J.
1995. PHYLIP: phylogeny inference package, version 3.57c. Univ. Washington, Seattle, WA.

Fitch, W. M., and E. Margoliash.
1967. Construction of phylogenetic trees. Science (Washington, D.C.) 155:279–284.

Gharrett, A. J., S. M. Shirley, and G. R. Tromble.
1987. Genetic relationships among populations of Alaskan chinook salmon (*Oncorhynchus tshawytscha*). Can. J. Fish. Aquat. Sci. 44:765–774.

Gibbons, A.
1997. Calibrating the mitochondrial clock. Science (Washington, D.C.) 279:28–29.

Gold, J. R ., and L. R . Richardson.
1991. Genetic studies in marine fishes. IV. An analysis of population structure of the red drum (*Sciaenops ocellatus*) using mitochondrial DNA. Fish. Res. 12:213–241.

Gold, J. R., L. R. Richardson, C. Furman, and T. L. King.
1993. Mitochondrial DNA differentiation and population structure in red drum (*Sciaenops ocellatus*) from the Gulf of Mexico and Atlantic Ocean. Mar. Biol. 116:175–185.

Gold, J. R., L. R. Richardson, C. Furman, and F. Sun.
1994. Mitochondrial DNA diversity and population structure in marine fish species from the Gulf of Mexico. Can. J. Fish. Aquat. Sci. 51(suppl. 1):205–314.

Gold, J. R., F. Sun, and L.R. Richardson.
1997. Population structure of red snapper from the Gulf of Mexico as inferred from analysis of mitochondrial DNA. Trans. Am. Fish. Soc. 126:386–396.

Graves, J. E., and J. R. McDowell.
1994. Genetic analysis of striped marlin (*Tetrapturus audax*) population structure in the Pacific Ocean. Can. J. Fish. Aquat. Sci. 51:1762–1768.
1995. Inter-ocean genetic divergence of istiophorid billfishes. Mar. Biol. 122:193–203.

Gyllensten, U., D. Wharton, A. Josefsson, and A. C. Wilson.
1991. Paternal inheritance of mitochondrial DNA in mice. Nature 352:255–257.

Hamilton, T. D.
1986. Late Cenozoic glaciation in the central Brooks Range. *In* Glaciation in Alaska: the geologic record (T. D. Hamilton, K. M. Reed, and R. M. Thorson, eds.), p. 9–49. Alaska Geological Soc., Anchorage, AK.

Hedges, S. B.
1992. The number of replications needed for accurate estimation of the bootstrap *P* value in phylogenetic studies. Mol. Biol. Evol. 9:366–369.

Kimura, M., and J. J. Crow.
1964. The number of alleles that can be maintained in a finite population. Genetics 49:725–738.

Kondzela, C. K., C. M. Guthrie, S. L. Hawkins, C. D. Russell, J. H. Helle, and A. J. Gharrett.
1994. Genetic relationships among chum salmon populations in Southeast Alaska and northern British Columbia. Can. J. Fish. Aquat. Sci. 51(suppl. 1):50–64.

Kornfield, I., and S. M. Bogdanowicz.
1987. Differentiation of mitochondrial DNA in Atlantic herring, *Clupea herengus*. Fish. Bull. 85:561–568.

Martin, A. P., G. P. Naylor, and S. R. Palumbi.
1992. Rates of mitochondrial DNA evolution in sharks are slow compared with mammals. Nature 357:153–155.

McElroy, D. M., P. Moran, E. Bermingham, and I. Kornfield.
1990. REAP: an integrated environment for the manipulation and phylogenetic analysis of restriction data. J. Hered. 83:157–158.

McGregor, A. J.
1983. A biochemical genetic analysis of pink salmon (*Oncorhynchus gorbuscha*) from selected streams in northern Southeast Alaska. M.S. thesis, Univ. Alaska-Juneau, Juneau, AK, 94 p.

Nedbal, M. A., and D. P. Phillipp.
1994. Differentiation of mitochondrial DNA in largemouth bass. Trans. Am. Fish. Soc. 123:460–468.

Nei, M.
1987. Molecular evolutionary genetics. Columbia Univ. Press, New York, NY, 512 p.

Nei, M., and J. C. Miller.
1990. A simple method for estimating average number of nucleotide substitutions within and between populations from restriction data. Genetics 97:145–163.

Nei, M., and F. Tajima.
1983. Maximum likelihood estimation of the number of nucleotide substitutions from restriction site data. Genetics 97:145–163.

Neigel, J. E., and J. C. Avise.
1986. Phylogenetic relationships of mitochondrial DNA under various demographic models of speciation. *In* Evolutionary processes and theory (E. Nevo and S. Karlin, eds.), p. 514–534. Academic Press, New York, NY.

Noll, C., N. V. Varnavskaya, E. A. Matzak, S. L. Hawkins, V. V. Midanaya, O. N. Katugin, C. Russell, N. M. Kinas, C. M. Guthrie III, H. Mayama, F. Yamazaki, B. P. Finney, and A. J. Gharrett.
2001. Analysis of contemporary genetic structure of even-broodyear populations of Asian and western Alaskan pink salmon, *Oncorhynchus gorbuscha*. Fish. Bull. 99:123–138.

Ohta, T., and M. Kimura.
1973. A model of mutation appropriate to estimate the number of electrophoretically detectable alleles in a finite population. Genet. Res. 22:201–204.

Park, L. K., M. A. Brainard, D. A. Dightman, and G. A. Winans.
1993. Low levels of intraspecific variation in mitochondrial DNA of chum salmon (*Oncorhynchus keta*). Mol. Mar. Biol. Biotech. 2:362–370.

Petit, J. R., and 18 coauthors.
1999. Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica. Nature 399: 429–436.

Phelps, S. R. , L. L. LeClair, S. Young, and H. L. Blankenship.
1994. Genetic diversity patterns of chum salmon in the Pacific Northwest. Can. J. Fish. Aquat. Sci. 51(suppl. 1):65–83.

Phillips, R. B., and T. H. Oakley.
1997. Phylogenetic relationships among Salmoninae based on nuclear and mitochondrial DNA sequences. *In* Molecu-

lar systematics of fishes (T. D. Kocher and C. A. Stepien, eds.) p. 145–162. Academic Press, San Diego, CA.

Porter, S. C.
1989. Some geological implications of average quaternary glacial conditions. Quaternary Research 32:245–261.

Posada, D, K.A. Crandall, and A. Templeton.
2000. GeoDis: A program for the cladistic nested analysis of the geographical distribution of genetic haplotypes. Mol. Ecol. 9:487–488.

Pustavoit, S. P.
1995. Features of genetic structure of coho salmon (*Oncorhynchus kisutch* Walbaum) populations. Genetika 31: 709–714. [Abstract in English.]

Reisenbichler, R. R., and S. R. Phelps.
1987. Genetic variation in chinook, *Oncorhynchus tshawytscha*, and coho, *O. kisutch*, salmon from the north coast of Washington. Fish. Bull. 85:681–701.

Richardson, L. R., and J. R. Gold.
1993. Mitochondrial DNA variation on red grouper (*Epinephelus morio*) and greater amberjack (*Seriola dumerili*) from the Gulf of Mexico. ICES J. Mar. Sci. 50:53–62.
1997. Mitochondrial DNA diversity in and population structure of red grouper, *Epinephelus morio*, from the Gulf of Mexico. Fish. Bull. 95:174–179.

Roe, B. A., D-P. Ma, R. K. Wilson, and J. F-H. Wong.
1985. The complete nucleotide sequence of the *Xenopus laevis* mitochondrial genome. J. Biol. Chem. 260:9759–9774.

Roff, D. A., and P. Bentzen.
1989. The statistical analysis of mitochondrial DNA polymorphisms: $\chi^2$ and the problem of small samples. Mol. Biol. Evol. 6:539–545.

Sandercock, F. K.
1991. Life history of coho salmon (*Oncorhynchus kisutch*). *In* Pacific salmon life histories (C. Groot and L. Margolis, eds.), p. 395–445. UBC Press, Vancouver, B.C., Canada.

Schneider, S., J-M. Kueffer, D. Roesli, and Laurant Excoffier.
1997. Arlequin version 1.1: a software for population genetic data analysis. Genetics and Biometry Laboratory, Univ. Geneva, Switzerland.

Scribner, K. T., P. A. Crane, W. J. Spearman, and L. W. Seeb.
1998. DNA and allozyme markers provide concordant estimates of population differentiation: analyses of U.S. and Canadian populations of Yukon River fall-run chum salmon (*Oncorhynchus keta*). Can. J. Fish. Aquat. Sci. 55:1748–1758.

Seeb, J. E., C. Habicht, W. D. Templin, L. W. Seeb, J. B. Shaklee, and F. M. Utter.
1999. Allozyme and mitochondrial DNA variation describe ecologically important genetic structure of even-year pink salmon inhabiting Prince William Sound, Alaska. Ecol. Freshwater Fish 8:122–140.

Seeb, L. W., and P. A. Crane.
1999. Allozyme and mitochondrial DNA discriminate Asian and North American populations of chum salmon in mixed stock fisheries along the south coast of the Alaska Peninsula. Trans. Am. Fish. Soc. 128:88–103.

Seutin, G., B. N. White, and P. T. Boag.
1991. Preservation of avian blood and tissue samples for DNA analysis. Can. J. Zool. 69:82–90.

Shaklee, J. B., T. D. Beacham, L. Seeb, and B. A. White.
1999. Managing fisheries using genetic data: case studies from four species of Pacific salmon. Fish. Res. 43:45–78.

Shields, G. F., and A. C. Wilson.
1987. Calibration of mitochondrial DNA evolution in geese. J. Mol. Evol. 24:212–217.

Slatkin, M. and R. R. Hudson.
1991. Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. Genetics 129:555–562.

Smith, G. R.
1992. Introgression in fishes: significance for paleontology, cladistics, and evolutionary rates. Syst. Biol. 41:41–57.

Taylor, E. B., C. J. Foote, and C. C. Wood.
1996. Molecular genetic evidence for parallel life-history evolution within a Pacific salmon (sockeye salmon and kokanee, *Oncorhynchus nerka*). Evolution 50:401–416.

Taylor, E. B., S. Harvey, S. Pollard, and J. Volpe.
1997. Postglacial genetic differentiation of reproductive ecotypes of kokanee *Oncorhynchus nerka* in Okanagan Lake, British Columbia. Mol. Ecol. 6:503–517.

Templeton, A. R.
1998. Nested clade analysis of phylogeographic data: testing hypotheses about gene flow and population history. Mol. Ecol. 7:381–397.

Templeton, A. R., and C. F. Sing.
1993. A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. IV. Nested analysis with cladogram uncertainty and recombination. Genetics 134:659–669.

Thomas, W. K., and A. T. Beckenbach.
1989. Variation in salmonid mitochondrial DNA: evolutionary constraints and mechanisms of substitution. J. Mol. Evol. 29:233–245.

Thomas, W. K., R. E. Withler, and A. T. Beckenbach.
1986. Mitochondrial DNA analysis of Pacific salmonid evolution. Can. J. Zool. 64:1058–1064.

Utter, F. M., G. Milner, G. Ståhl, and D. Teel.
1989. Genetic population structure of chinook salmon, *Oncorhynchus tshawytscha*, in the Pacific Northwest. Fish. Bull. 15:575–576.

Varnavskaya, N. V., C. C. Wood, and R. J. Everett.
1994. Genetic variation in sockeye salmon (*Oncorhynchus nerka*) populations in Asia and North America. Can. J. Fish. Aquat. Sci. 51(suppl. 1):132–146.

Wallace, D. W.
1987. Large- and small-scale phenol extractions. *In* Methods in enzymology, vol. 12: guide to molecular cloning techniques (S. L. Berger and A. R. Kimmel, eds.), p. 33–41. Academic Press, San Diego, CA.

Weir, B.
1996. Genetic data analysis II. Sinauer Press, Sunderland, MA, 445 p. + xii,

Wehrhahn, C. F., and R. Powell.
1987. Electrophoretic variation, regional differences, and gene flow in the coho salmon (*Oncorhynchus kisutch*) of southern British Columbia. Can. J. Fish. Aquat. Sci. 44:822–831.

Wilmot, R. L., R. J. Everett, W. J. Spearman, R. Baccus, N. V. Varnavskaya, and S. V. Putivkin.
1994. Genetic stock structure of western Alaska chum salmon and a comparison with Russian Far East stocks. Can. J. Fish. Aquat. Sci. 51(suppl. 1):84–94.

Wilson, G. M., W. K. Thomas, and A. T. Beckenbach.
1987. Mitochondrial DNA analysis of Pacific Northwest populations of *Oncorhynchus tshawytscha*. Can. J. Fish. Aquat. Sci. 44:1301–1305.

Winans, G. A., P. B. Aebersold, S. Urawa, and N. V. Varnavskaya.
1994. Determining the continent of origin of chum salmon (*Oncorhynchus keta*) using genetic stock identification techniques: status of allozyme baseline in Asia. Can. J. Fish. Aquat. Sci. 51(suppl. 1):105–113.

Wood, C. C., B. E. Riddell, D. T. Rutherford, and R. E. Withler.
1994. Biochemical genetic survey of sockeye salmon (*Oncorhynchus nerka*) in Canada. Can. J. Fish. Aquat. Sci. 51(suppl. 1):114–131.

Zardoya, R., A. Garrido-Pertierra, and J. M. Bautista.
1995. The complete nucleotide sequence of the mitochondrial DNA genome of the rainbow trout, *Oncorhynchus mykiss*. J. Mol. Evol. 41:942–951.

Zhivotovsky, L. A., A. J. Gharrett, M. K. Glubokovsky, and M. W. Feldman.
1994. Gene differentiation in Pacific salmon (*Oncorhynchus* sp.): facts and models with reference to pink salmon (*O. gorbuscha*). Can. J. Fish. Aquat. Sci. 51 (suppl. 1):223–232.

Zink, R. M., J. M. Fitzsimmons, D. L. Dittmann, D. R. Reynolds, and R. T. Nishimoto.
1996. Evolutionary genetics of Hawaiian freshwater fish. Copeia 1996:330–335.

## Appendix 1

Restriction fragment sizes for seven PCR-amplified coho salmon mtDNA regions for twelve restriction endonucleases. Composite haplotypes are shown in Appendix 2.

2S/165

| *Ase* I | *Ava* I | *Bst*N I | *Bst*U1 | *Cfo* I | | *Dde* I | | | *Hind* II | *Hinf* I | *Mbo* I | | *Msp* I | *Rsa* I | *Sty* I |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 | 2 | 1 | 2 | 3 | 1 | 1 | 1 | 2 | 1 | 1 | 1 |
| 2400 | 1200 | 1000 | 1300 | 1210 | 1100 | 455 | 400 | 400 | 750 | 1800 | 1475 | 900 | 950 | 930 | 1500 |
| | 1200 | 410 | 475 | 380 | 380 | 400 | 280 | 280 | 720 | 600 | 375 | 575 | 650 | 600 | 500 |
| | | 380 | 300 | 350 | 350 | 280 | 250 | 205 | 720 | | 210 | 375 | 400 | 290 | 200 |
| | | 310 | 275 | 150 | 150 | 155 | 205 | 205 | 210 | | 170 | 210 | 190 | 200 | 200 |
| | | 200 | | 120 | 120 | 145 | 155 | 155 | | | 170 | 170 | 130 | 200 | |
| | | 150 | | | 110 | 130 | 145 | 145 | | | | 170 | 80 | 180 | |
| | | | | | | 125 | 130 | 130 | | | | | | | |
| | | | | | | 120 | 125 | 125 | | | | | | | |
| | | | | | | 104 | 120 | 120 | | | | | | | |
| | | | | | | 80 | 104 | 104 | | | | | | | |
| | | | | | | 65 | 80 | 80 | | | | | | | |
| | | | | | | 55 | 65 | 65 | | | | | | | |
| | | | | | | 48 | 55 | 55 | | | | | | | |
| | | | | | | 45 | 48 | 48 | | | | | | | |
| | | | | | | 37 | 45 | 45 | | | | | | | |
| | | | | | | 35 | 37 | 45 | | | | | | | |
| | | | | | | 33 | 35 | 37 | | | | | | | |
| | | | | | | 30 | 33 | 35 | | | | | | | |
| | | | | | | 27 | 30 | 33 | | | | | | | |
| | | | | | | | 27 | 30 | | | | | | | |
| | | | | | | | | 27 | | | | | | | |

ND1/ND2

| *Ase* I | *Ava*I | *Bst*N I | *Bst*U I | *Cfo* I | *Dde* I | | | *Hind* II | *Hinf* I | *Mbo* I | *Msp* I | *Rsa* I | *Sty* I | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 | 1 | 2 | 3 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 3 | 4 |
| 1786 | 1596 | 1496 | 2026 | 1146 | 780 | 780 | 780 | 1846 | 1546 | 1026 | 1926 | 1176 | 650 | 650 | 550 | 550 |
| 520 | 600 | 1150 | 500 | 900 | 480 | 480 | 480 | 800 | 850 | 590 | 780 | 375 | 550 | 550 | 530 | 530 |
| 230 | 450 | | 120 | 600 | 240 | 240 | 240 | | 250 | 510 | | 275 | 530 | 530 | 470 | 470 |
| 110 | | | | | 220 | 210 | 180 | | | 310 | | 250 | 395 | 305 | 395 | 305 |
| | | | | | 180 | 180 | 148 | | | 210 | | 220 | 305 | 305 | 305 | 305 |
| | | | | | 145 | 145 | 145 | | | | | 220 | 230 | 230 | 230 | 230 |
| | | | | | 123 | 123 | 123 | | | | | 130 | | 90 | 180 | 180 |
| | | | | | 120 | 120 | 120 | | | | | | | | | 90 |
| | | | | | 115 | 115 | 115 | | | | | | | | | |
| | | | | | 95 | 95 | 95 | | | | | | | | | |
| | | | | | 78 | 78 | 78 | | | | | | | | | |
| | | | | | | (10) | 62 | | | | | | | | | |
| | | | | | | | (10) | | | | | | | | | |

*continued*

**Appendix 1 (continued)**

A8/A6/COIII/ND3

| *Ase* I | *Ava* I | *Bst*N I | *Bst*U I | *Cfo* I | *Dde* I | *Hin*d II | *Hinf* I | *Mbo* I | *Msp* I | *Rsa* I | *Sty* I |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 2210 | 2210 | 1200 | 1700 | 1300 | 970 | 1300 | 500 | 1800 | 800 | 780 | 500 |
| | | 700 | 500 | 900 | 970 | 900 | 498 | 210 | 600 | 680 | 450 |
| | | 310 | | | 180 | | 400 | 200 | 490 | 345 | 350 |
| | | | | | 80 | | 300 | | 320 | 240 | 340 |
| | | | | | | | 160 | | | 150 | 240 |
| | | | | | | | 150 | | | | 220 |
| | | | | | | | 150 | | | | 90 |

COI/COII

| *Ase* I | *Ava* I | *Bst*N I | *Bst*U I | *Cfo* I | *Dde* I | | | *Hin*d II | *Hinf* I | | *Mbo* I | | *Msp* I | *Rsa* I | *Sty* I |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 | 1 | 2 | 3 | 1 | 1 | 2 | 1 | 2 | 1 | 1 | 1 |
| 1562 | 2432 | 1152 | 1422 | 1522 | 530 | 530 | 530 | 1682 | 575 | 500 | 650 | 550 | 2000 | 872 | 1282 |
| 650 | | 890 | 500 | 1010 | 360 | 360 | 360 | 750 | 500 | 500 | 550 | 520 | 220 | 350 | 1000 |
| 220 | | 390 | 410 | | 330 | 330 | 330 | | 410 | 410 | 480 | 480 | 212 | 350 | 150 |
| | | | | | 293 | 293 | 293 | | 295 | 295 | 270 | 270 | | 300 | |
| | | | | | 255 | 255 | 255 | | 150 | 150 | 220 | 220 | | 225 | |
| | | | | | 235 | 235 | 235 | | 150 | 150 | 175 | 175 | | 215 | |
| | | | | | 196 | 196 | 209 | | 150 | 150 | 75 | 130 | | 120 | |
| | | | | | 119 | 110 | 110 | | 90 | 90 | | 75 | | | |
| | | | | | 110 | 80 | 80 | | | 75 | | | | | |
| | | | | | 26 | 39 | 39 | | | | | | | | |
| | | | | | (13) | 26 | 26 | | | | | | | | |
| | | | | | | (13) | | | | | | | | | |

ND3/ND4

| *Ase* I | *Ava* I | *Bst*N I | *Bst*U I | *Cfo* I | | *Dde* I | *Hin*d II | *Hinf* I | *Mbo* I | *Msp* I | *Rsa* I | *Sty* I |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | 1 | 2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 1955 | 2105 | 1050 | 1800 | 1600 | 1100 | 525 | 2305 | 1730 | 525 | 900 | 875 | 1970 |
| 350 | 200 | 750 | 500 | 700 | 700 | 490 | | 550 | 500 | 800 | 490 | 330 |
| | | 500 | | | 500 | 400 | | | 400 | 300 | 365 | |
| | | | | | | 290 | | | 350 | 250 | 230 | |
| | | | | | | 150 | | | 190 | 115 | 140 | |
| | | | | | | 150 | | | 190 | | 110 | |
| | | | | | | 90 | | | 150 | | 90 | |
| | | | | | | 90 | | | | | | |

ND5/ND6

| *Ase* I | *Ava* I | | *Bst*N I | | *Bst*U I | *Cfo* I | *Dde* I | | | | *Hin*d II | *Hinf* I | *Mbo* I | | *Msp* I | *Rsa* I | *Sty* I |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 2 | 1 | 2 | 1 | 1 | 1 | 2 | 3 | 4 | 1 | 1 | 1 | 2 | 1 | 1 | 1 |
| 1100 | 2530 | 2230 | 900 | 900 | 2025 | 1000 | 665 | 600 | 600 | 600 | 1950 | 550 | 1442 | 1442 | 1525 | 2530 | 1675 |
| 850 | | 300 | 900 | 900 | 525 | 800 | 490 | 570 | 490 | 490 | 550 | 475 | 410 | 410 | 410 | | 600 |
| 550 | | | 650 | 400 | | 500 | 300 | 490 | 320 | 300 | | 420 | 312 | 312 | 285 | | 275 |
| | | | 100 | 250 | | 200 | 270 | 320 | 300 | 270 | | 320 | 286 | 281 | 200 | | |
| | | | | 100 | | | 200 | 160 | 270 | 200 | | 275 | 100 | 100 | 130 | | |
| | | | | | | | 160 | 150 | 160 | 160 | | 210 | | (5) | | | |
| | | | | | | | 150 | 145 | 150 | 150 | | 175 | | | | | |
| | | | | | | | 145 | 65 | 145 | 145 | | 125 | | | | | |
| | | | | | | | 120 | | 65 | 120 | | | | | | | |
| | | | | | | | | | | 65 | | | | | | | |

*continued*

**Appendix 1 (continued)**

| CytB/D-Loop | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *Ase* I | *Ava* I | *Bst*N I | | | *Bst*U I | *Cfo* I | *Dde* I | *Hin*d II | *Hin*f I | *Mbo* I | *Msp* I | *Rsa* I | *Sty* I |
| 1 | 1 | 1 | 2 | 3 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 1524 | 1534 | 900 | 900 | 900 | 2100 | 900 | 550 | 2094 | 690 | 810 | 700 | 1240 | 1050 |
| 600 | 520 | 850 | 540 | 540 | 444 | 694 | 430 | 450 | 650 | 610 | 330 | 400 | 580 |
| 420 | 315 | 415 | 415 | 400 | | 510 | 310 | | 470 | 400 | 240 | 390 | 450 |
| | 175 | 400 | 400 | 310 | | 300 | 290 | | 310 | 170 | 225 | 195 | 220 |
| | | 130 | 310 | 220 | | 140 | 280 | | 250 | 165 | 210 | 140 | 140 |
| | | | 130 | 195 | | | 280 | | 230 | 165 | 205 | 125 | 110 |
| | | | | 130 | | | 175 | | | 150 | 205 | 105 | 70 |
| | | | | | | | 130 | | | 72 | 180 | | |
| | | | | | | | 90 | | | | 180 | | |
| | | | | | | | | | | | 120 | | |

**Appendix 2**

Key to haplotype composition inferred from variable fragment patterns in Appendix 1. No variation was observed in the A8/COIII region (Table 2). Numbers in parentheses are presumed haplotypes.

| | ND1/ND2 | | 12S/16S | | | ND3/ ND4 | Cytb/ Dloop | ND5/ND6 | | | | COI/A8 | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Haplotype | *Dde* I | *Sty* I | *Cfo* I | *Mbo* I | *Dde* I | *Cfo* I | *Bst*N I | *Ava* I | *Bst*N I | *Dde* I | *Mbo* I | *Dde* I | *Hin*F I | *Mbo* I |
| A | 2 | 4 | 2 | 1 | 2 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 1 |
| A′ | 2 | 4 | 2 | 1 | 2 | 1 | 2 | 2 | 1 | 2 | 2 | 2 | 2 | 1 |
| B | 2 | 4 | 1 | 1 | 2 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 1 |
| C | 2 | 4 | 2 | 1 | 2 | 1 | 3 | 2 | 2 | 2 | 2 | 2 | 2 | 1 |
| C′ | 2 | 4 | 2 | 2 | 2 | 1 | 3 | 2 | 2 | 2 | 2 | 2 | 2 | 1 |
| D | 2 | 3 | 2 | 1 | 2 | 1 | 3 | 2 | 2 | 2 | 2 | 2 | 2 | 1 |
| E | 3 | 2 | 2 | 1 | 2 | 1 | 1 | 1 | 2 | 4 | 1 | 2 | 2 | 1 |
| E′ | 3 | 2 | 2 | 1 | 2 | 1 | 1 | 1 | 2 | 4 | 1 | 2 | 1 | 2 |
| F | 3 | 1 | 2 | 1 | 2 | 1 | 1 | 1 | 2 | 4 | 1 | 2 | 2 | 1 |
| G | 3 | 2 | 2 | 1 | 2 | 1 | 1 | 1 | 2 | 3 | 1 | 1 | 2 | 1 |
| H | 3 | 2 | 2 | 1 | 1 | 1 | 1 | 1 | 2 | 4 | 1 | 2 | 2 | 1 |
| I | 2 | 4 | 2 | | 2 | 1 | 1 | 2 | | 2 | 2 | 2 | | |
| J | 2 | 4 | 2 | | 2 | 1 | 3 | 1 | | 3 | 2 | 3 | | |
| K | 1 | 2 | 2 | | 2 | 1 | 1 | 1 | | 4 | 2 | 2 | | |
| L | 3 | 1 | 2 | | 2 | 1 | 1 | 1 | | 3 | 1 | 2 | | |
| M | 3 | 2 | 2 | | 2 | 1 | 1 | 1 | | 1 | (1) | (2) | | |
| N | 3 | 2 | 2 | | 3 | 1 | 1 | 1 | | 5 | (1) | (2) | | |
| O | 3 | 2 | 2 | | 1 | 2 | 1 | 1 | | 5 | (1) | (2) | | |
| P | 2 | 4 | 2 | | 2 | 1 | 3 | 1 | | 3 | 2 | 2 | | |

# Diet and consumption rates of overwintering YOY striped bass, *Morone saxatilis*, in the Hudson River*

**Thomas P. Hurst**
**David O. Conover**
Marine Sciences Research Center
State University of New York
Stony Brook, New York 11794-5000
E-mail address (for T. P. Hurst): hurst@msrc.sunysb.edu

**Abstract**–Starvation is often proposed as the mechanism of size-dependent mortality of overwintering temperate fishes, yet little is known about the energetics of fish at low temperatures. Young-of-the-year (YOY) Hudson River striped bass (*Morone saxatilis*) suffer a winter energy deficit and experience size-selective winter mortality in some years that may influence recruitment. To better understand the role of energetic stress in winter mortality, we determined diet composition and measured consumption rates of wild fish. Gastric evacuation rates were measured in the laboratory at 2°, 5°, 8°, and 11°C. Measured evacuation rates for YOY striped bass were among the lowest reported, with 25% of the initial meal remaining after 150 hours at 2°C. Variation in evacuation rate data increased as temperature decreased. Diets of fish captured in winter were dominated by gammarid amphipods and shrimp species. Gut fullness ranged from 0% to 7%, averaging 0.4% body weight. Evacuation rates, gut fullness values, and river temperatures were combined to estimate daily consumption rates on 29 dates over five winters. Consumption estimates ranged from 0% to 0.29%bw/day and were generally higher in early winter than in late winter and not correlated with river temperatures. Stomach fullness was negatively correlated with the level of lipid energy reserves at the individual level. The patterns may reflect an internal control on appetite, or a depression of prey availability in late winter. These findings indicate that the potential for winter starvation may be influenced by both internal and external constraints on consumption rate.

Manuscript accepted 15 March 2001.
Fish. Bull. 99:545–553 (2001)

Although the feeding and growth of many temperate fish species have been extensively examined during summer months, details of the feeding ecology of most species during winter remain largely unknown. Knowledge of the winter energetics of many fish species is restricted to the observation that growth decreases markedly in autumn and often becomes negligible during winter. There are only scattered reports of the feeding patterns of overwintering temperate fishes (Daiber, 1956; Keast, 1968; Foltz and Norden, 1977; Diana, 1979), and few estimates of consumption rates (but see Diana, 1979; Minton and McLean, 1982). This lack of information on the winter energetics of fishes applies also to laboratory-based physiological measurements, including gastric evacuation rates and metabolic rates. Despite the observation that temperature is the predominant factor regulating digestion, for none of the 22 species listed in a compilation of gastric evacuation data (He and Wurtsbaugh, 1993) had rates been measured at the minimum temperatures likely to be encountered in winter.

Many temperate fish species lose energy throughout the winter, relying upon stored lipid reserves to fuel metabolism (Toneys and Coble, 1980; Schultz and Conover, 1997; Hurst et al., 2000). Reliance on lipid reserves is believed to be necessary because consumption rate is severely limited by the ability of fish to digest food at low temperatures. However low consumption rates of overwintering fish may also be limited by low food availability (Cunjak and Power, 1987; Foy and Paul, 1999).

Winter mortality that selects against smaller fish has been documented in a number of temperate species (Sogard, 1997) and is most commonly attributed to depletion of lipid energy reserves and starvation. Smaller fish have higher starvation mortality rates because they tend to have lower energy reserves and higher weight-specific metabolic rates than larger fish. An implicit assumption in the "starvation hypothesis" is that overwintering fish have little or no capacity to obtain energy by feeding. However, in several experiments, fish allowed to feed *ad libitum* had higher winter survival rates than unfed fish (Post and Evans, 1989; Thompson et al., 1991; Hurst and Conover, 1998). Although wild fish are unlikely to have access to unlimited food, these results suggest that fish benefit from feeding during winter. Furthermore, some overwintering fish increase consumption rates in response to depletion of internal reserves (Metcalfe and Thorpe, 1992; Bull et al., 1996; Hurst and Conover, 2001). These findings suggest the need for a more complete understanding of the energetics of overwintering fish, including consumption rates of wild fish and an evaluation of factors that regulate consumption in winter.

The striped bass (*Morone saxatilis*) is a relatively well studied, commercially important species along the east coast of the United States and Canada (Boreman and Austin, 1985). Previous work has shown that recruitment to the Hudson River population may be regulated by the severity of the winter that age-0 fish encounter (Hurst and Conover, 1998). In addition, size-selec-

* Contribution 1226 of the Marine Sciences Research Center, State University of New York, Stony Brook, New York 11794.

tive winter mortality has been documented among young-of-the-year (YOY) striped bass in both the Hudson River and Miramichi River populations (Hurst and Conover, 1998; Bradford and Chaput[1]).

Young-of-the-year striped bass in the Hudson River experience a winter energy deficit that varies in severity among years (Hurst et al., 2000). This interannual variability may be related to feeding conditions in the environment. Although studies of the summer feeding habits of YOY striped bass are available from most east coast estuaries (Markle and Grant, 1970; Boynton et al., 1981; Gardinier and Hoff, 1982; Rulifson and McKenna, 1987), information for overwintering fish is extremely limited (Hartman and Brandt, 1995a). A bioenergetics model has been developed for juvenile striped bass (Hartman and Brandt, 1995b), describing metabolic rates and maximum consumption rates of fish across a wide range of temperatures. This type of deterministic bioenergetic modeling does not take into account the observed variability in energy storage and depletion cycles (Hurst et al., 2000) or compensatory feeding responses observed in response to depletion of reserves (Metcalfe and Thorpe, 1992). A better understanding of the energetics of overwintering juvenile striped bass, including *in situ* estimates of consumption rates, is required to fully evaluate the potential for winter starvation in this species.

In our study, we present results of experiments measuring gastric evacuation rates at winter temperatures necessary to determine consumption rates of wild fish. We also describe the diet of overwintering YOY striped bass and estimate consumption rates of overwintering fish on 29 dates over five winters. Finally, we analyze feeding patterns at both the individual and population level to determine the factors regulating consumption rates of overwintering fish and discuss these results as they relate to the potential for winter starvation.

## Methods

### Gastric evacuation experiments

Wild fish were captured from the Hudson River estuary and transported in river water to the Flax Pond Marine Laboratory of the State University of New York at Stony Brook in Old Field, New York. Fish were treated with 0.60 ppm copper sulphate for 5 minutes and 15 ppm oxytetracycline for 5 days to reduce risk of mortality from infection. Fish were acclimated to laboratory conditions for at least 3 weeks prior to use and only fish appearing healthy and behaving normally were used in the experiment. Temperatures during the acclimation period were maintained between 1° and 5°C, and salinities were maintained at 15 ppt. Fish were fed frozen adult brine shrimp (*Artemia* sp.) and sand shrimp (*Crangon septemspinosa*) daily during the acclimation period. Prior to experimentation, groups of fish were acclimated to the test temperature (±0.5°C) for at least one week. Fish rarely fed voluntarily at low temperatures and were not offered food for between 3 and 5 days prior to use in the experiment.

Evacuation rates were measured at 2°, 5°, 8° and 11°C. Temperatures in the experimental tanks were maintained by recirculating fluid chillers; they never differed from the prescribed temperature by more than 0.5° and were generally within 0.2°C. Fish used in the experiment ranged in size from 88 to 150 mm TL (5.6 to 31.4 g wet weight), the natural size range of YOY striped bass in winter. To reduce stress during the feeding process, fish were weighed (to 0.01 g wet weight) and acclimated to 65-L test tanks for 12 hours prior to feeding. Individual fish were captured with a dip-net from the test tank, force-fed the meal, and returned to the test tank within 1 minute. Fish were fed by opening the mouth and forcing the meal through the esophagus with a pair of blunt forceps. A meal comprised a single whole or partial *C. septemspinosa* weighing approximately 2% of the fish's body weight. A ration level of 2% body weight was chosen because it falls in the upper range of, but is well below the maximum, gut fullness levels observed among wild fish. Each fish remained in the test tank until the time of sampling, when it was netted and sacrificed with an overdose of MS-222 anesthetic and measured (to 1.0 mm TL). The remaining stomach contents were dissected from the fish, dried of excess water, weighed (to 0.001 g), and dried to a constant weight at 60°C. In no cases did fish regurgitate the meal following feeding or during the netting and sacrifice procedure.

The evacuation rate of at least 3 fish was measured at each of 10 or more time points at each temperature (min. 33 fish at 11°C; max. 43 fish at 2°C). The maximum interval between feeding and sacrifice encompassed the majority of the evacuation time at each temperature and ranged from 72 hours at 11°C to 168 hours at 2°C. The minimum interval used was 0.25 hours at all temperatures. Fish were exposed to a 10:14 light:dark cycle to mimic the natural winter photoperiod. Feeding time was standardized to the light:dark cycle, and fish were exposed to light for the first 8–10 hours of digestion.

Evacuation of the meal was described with an exponential model allowing a time-lag prior to the beginning of evacuation because of its utility in estimating rations among wild fish (Elliott and Persson, 1978; Bromley, 1994). The model was fitted by using biphasic nonlinear regression of untransformed data on the percentage of initial meal remaining over time with the equation

$$\%\ remaining = \begin{cases} 100 & if\ \ t < (c_0 + c_1 T) \\ 100e^{-(b_0 e^{b_1 T})[t-(C_0+C_1 T)]} & if\ \ t \geq (c_0 + c_1 T), \end{cases}$$

where $T$ = temperature;
$t$ = time in hours since ingestion;
$b_0$ and $b_1$ = the coefficients of the exponential relationship between evacuation rate and temperature; and

[1] Bradford, R. G.,and G. Chaput. 1997. Status of striped bass (*Morone saxatilis*) in the Gulf of St. Lawrence in 1996 and revised estimates of spawner abundance for 1994 and 1995. Dep. Fish. and Oceans, Can. Stock Assess Secretariat Res. Doc. 97/16, 31 p. Dep. Fisheries and Oceans, Science Branch, Gulf Region, P.O. Box 5030, Moncton, New Brunswick E1C 9B6, Canada.

**Table 1**

Summary of collections and diets of overwintering YOY striped bass collected in the Hudson River Estuary, 1993–97. %F = mean frequency of occurrence, %W = mean percentage of wet weight; $n$ = number of fish sampled per date.

| | | Winter | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 1993 | | 1994 | | 1995 | | 1996 | | 1997 | |
| No. of dates sampled | | 4 | | 8 | | 7 | | 6 | | 4 | |
| Temperature (°C) | | 2.0–5.1 | | 4.2–10.4 | | 1.0–9.4 | | 3.3–6.2 | | 5.0–6.1 | |
| $n$ | | 20–34 | | 15–53 | | 8–37 | | 16–29 | | 18–52 | |
| Percentage empty stomachs | | 71.0 | | 52.5 | | 72.3 | | 57.6 | | 73.4 | |
| Mean gut fullness ±SE | | 0.44 ±0.103 | | 0.51 ±0.056 | | 0.34 ±0.066 | | 0.18 ±0.037 | | 0.26 ±0.060 | |
| Prey type | Scientific name | %F | %W | %F | %W | %F | %W | %F | %W | %F | %W |
| Amphipods | *Gammarus* sp. | 66.7 | 59.1 | 92.1 | 90.2 | 58.2 | 39.5 | 86.3 | 52.4 | 34.3 | 15.7 |
| Sand shrimp | *Crangon septimspinosa* | 37.0 | 27.1 | 4.2 | 4.2 | 7.1 | 8.1 | 0 | 0 | 3.1 | 5.3 |
| Grass shrimp | *Palaemonetes pugio, Palaemonetes vulgaris* | 38.9 | 6.4 | 0 | 0 | 4.3 | 4.1 | 0 | 0 | 3.1 | 7.5 |
| Mysid shrimp | | 0 | 0 | 0 | 0 | 0 | 0 | 3.0 | 7.1 | 18.8 | 4.3 |
| Unidentifiable shrimp | *C. septimspinosa, P. pugio, P. vulgaris* | 0 | 0 | 0.7 | 0.4 | 27.5 | 22.8 | 16.4 | 32.5 | 21.4 | 26.4 |
| Crabs | | 33.3 | 5.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Polychaetes | | 0 | 0 | 0 | 0 | 17.9 | 23.3 | 0 | 0 | 0 | 0 |
| Oligochaetes | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 23.9 | 38.8 |
| Fish | *Anchoa mitchilli, Menidia menidia, Ammodytes americanus* | 1.9 | 1.7 | 5.4 | 5.3 | 2.9 | 2.2 | 7.1 | 8.0 | 3.1 | 1.9 |
| UIR[1] | | 40.7 | 24.4 | 11.9 | 8.5 | 28.2 | 6.0 | 31.8 | 29.2 | 42.2 | 25.1 |

[1] UIR = unidentified invertebrate remains. These were not included in the calculation of %W of other prey items.

$c_0$ and $c_1$ = the intercept and slope of the linear relationship between temperature and the lag prior to the beginning of evacuation.

If no lag is present in the data, the parameters $c_0$ and $c_1$ approach 0. The effect of body size on evacuation was assessed by examining the relationship between fish length and deviation from the best fitting evacuation model.

## Diet of YOY striped bass

Overwintering YOY striped bass were collected with a 9-m bottom trawl (38-mm stretch mesh codend) from the lower Hudson River estuary, the only known wintering aggregation for the Hudson River population (Dovel, 1992). Sampling occurred throughout the overwintering period of five consecutive years. Fish were captured during daylight hours between river mile 0 and 9 in conjunction with the New York Power Authority's hatchery evaluation and tagging program. Sampling in each year began in mid-December and ended in late March or early April. Bottom water temperatures were measured during sampling. The number of fish analyzed for diet per date ranged from 8 to 53 depending on catch rates (Table 1). Captured fish were individually wrapped and immediately frozen for preservation. In the laboratory, fish were thawed, measured (mm TL), weighed (g wet weight), and stomach contents were removed. Prey were identified to the lowest possible taxon and weighed. All prey in each category in each stomach were weighed as a group because the abundance and small size of the dominant prey item made individual measurements unfeasible (gammarid amphipods; often >50/stomach and <0.005 g each). Diets of YOY striped bass were described by the contribution of items expressed on the basis of both weight ($\%W_i$=*weight of prey$_i$ / weight of all identifiable prey*) and frequency ($\%F_i$=*100 × no. of stomachs with prey$_i$/no. of stomachs with prey*). Unidentifiable items were measured separately and are expressed as a percentage of the total weight of all stomach contents.

## Estimation of consumption rates

Daily consumption rate estimates were generated by combining gut fullness values (S=total weight of stomach contents/fish weight) of overwintering YOY striped bass with laboratory-determined gastric evacuation rates with the method of Eggers (1979):

$$C = 24R_e \times \overline{S},$$

where $\overline{S}$ = average gut fullness of field-caught fish; and
$R_e$ = exponential evacuation rate at measured field temperatures determined from laboratory experiments described above.

This simplified version of the Elliott and Persson (1978) model is appropriate when sampling is not conducted in discrete time intervals. We used gut-fullness measures from fish collected throughout the day, as described above. Although fish were sampled only during daylight hours, the sampling interval (24 hours) is substantially shorter than the evacuation time (>72 hours), even at the warmest temperatures observed; hence any diel feeding patterns will have little effect on our consumption estimates.

Standard deviations of consumption estimates were generated from a Taylor expansion of the consumption model

$$\sigma^2_{\overline{C}} = \left(\frac{d\overline{C}}{d\overline{S}}\right)^2 \sigma^2_{R_e} + \left(\frac{d\overline{C}}{dR_e}\right)^2 \sigma^2_{\overline{S}} + 2\left(\frac{d\overline{C}}{dR_e}\right)\left(\frac{d\overline{C}}{d\overline{S}}\right)\mathrm{cov}(R_e, \overline{S}).$$

The variance in the evacuation rate parameter ($R_e$) was estimated through a bootstrap procedure by fitting the evacuation model to 1000 sets of 152 observations sampled with replacement from the evacuation rate data. Since $R_e$ is a function of temperature, the variance in $R_e$ at a given temperature was determined by inserting the temperature into the 1000 model fits. The variance in $\overline{S}$ was estimated for each date as $\mathrm{var}(S_i)/n$.

The standard deviation of the consumption estimate depends upon the covariance between evacuation rate ($R_e$), measured in laboratory experiments, and gut fullness ($\overline{S}$), observed in wild fish. Because we have no way of measuring the covariance between these parameters, we evaluated the importance of this term by calculating the variance under three assumptions: 1) $\overline{S}$ and $R_e$ are not correlated, 2) they covary perfectly, and 3) they display perfect negative covariance.

### Analysis of feeding patterns

We investigated feeding patterns of YOY striped bass by examining the relationship between several factors (body size, time of year, water temperature, and energy storage) and gut fullness, at the individual and population level. Gut fullness was chosen over consumption rate for three reasons. First, gut fullnesses were measured directly, whereas consumption rates were estimated from a model by using gut fullnesses. Second, consumption estimates are directly dependent on temperature, an independent variable in these analyses. Finally, we were interested in determining the conditions that stimulate feeding, which we believe are more immediately reflected in the gut fullness measures.

Feeding patterns at the individual level were investigated by examining the relationships between gut fullness and the independent variables of body size, lipid level, water temperature, and time of year. Gut fullness values of individual fish were dominated by zeros (empty stomachs) and could not be transformed appropriately, preventing the use of parametric statistics. Because empty stomachs may reflect either a lack of prey availability or reduced appetite, we concentrated analyses at the individual level on the slope of the 95% quantile of gut fullness regressed against the independent variables (Scharf et al., 1998). This technique has been used to examine scatter-plots, when boundaries of a relationship between two factors are of interest, rather than the mean. In our case we were interested in the factors that stimulate feeding as opposed to developing a predictive relationship between independent variables and gut fullness. The 95% quantile line describes the relationship between maximum observed gut fullness and the independent variables. Mean gut fullness of all fish captured on a given date was used to investigate feeding patterns at the population level and was compared to average lipid level of fish captured on that date, time of year, and water temperature, by using Kendall's coefficient of rank correlation (Sokal and Rohlf, 1981).

Lipid levels were determined for a subsample of fish from the diet analysis. This involved determination of the percentage of dry body weight comprising non polar lipids (those used primarily for energy storage). Details of the compositional analysis procedure can be found in Hurst et al. (2000) and Schultz and Conover (1997). Compositional analysis was performed on 15 to 40 fish from each sampling date (except 13 December 1995). The relationship between lipid level and gut fullness at the individual level was examined by using all fish for which both pieces of information were available ($n$=587), whereas analyses at the population level used the average lipid level observed on a given date ($n$=28).

## Results

### Gastric evacuation rate

Evacuation rates of overwintering YOY striped bass declined greatly with temperature. At mid-winter temperatures, evacuation rates were among the lowest reported for any fish species. Time to 50% evacuation ranged from 31 hours at 11°C to 101 hours at 2°C (Fig. 1). The best fit parameters in the evacuation model were $b_0$=–0.00685, $b_1$=0.126, $c_0$=25.757, and $c_1$=–1.8033.

We observed a lag between feeding and a measurable loss of material from the stomach. The length of the observed lag decreased as temperature increased from 18.2 hours at 2°C to 6.1 hours at 11°C. The exponential models used here to describe digestion fitted the experimental data as well as, or significantly better than, other common models (linear and square root; Bromley 1994).

The amount of variability among individuals in evacuation rate increased as temperature decreased, leading to a poorer fit of the evacuation models at the lower temperatures (Fig. 1). Deviations from the model increased significantly as temperature decreased ($P$=0.018; ANOVA of absolute value of sample deviations from evacuation model). Body size had no effect on evacuation rate among juvenile striped bass. We found no correlation between residual values from the evacuation model and fish length ($r$=–0.03 $P$=0.685).




**Figure 1**

Gastric evacuation patterns of YOY striped bass fed *Crangon septemspinosa* at four temperatures. Points are observations of percentage of initial meal remaining in the stomach after a predetermined time. Lines represent evacuation model fitted by using all data.

## Diet composition

Benthic invertebrates were the dominant prey of YOY striped bass overwintering in the lower Hudson River estuary, making up 95.0% of the diet by weight; the remaining 5.0% were various fish prey (Table 1). Most striped bass captured during winter had empty stomachs (64.8%), and on two dates, all fish sampled had empty stomachs (30 March 1994 and 6 December 1994). Unidentifiable stomach contents ranged from 6.0% weight (%W) in 1995 to 29.2%W in 1996. The most common item in the diet was gammarid amphipods (those identified to genus were *Gammarus* sp., but most were not identified) making up 51.3%W and occurring in 81% of stomachs with food (%F). Several species of shrimp were also common dietary items, including sand shrimp (*C. septemspinosa*) and grass shrimps (*Palaemonetes* spp.). Other invertebrates that were important items on several dates included mysid shrimp, polychaetes, and oligochaetes. Fish were rarely found in the stomachs of YOY striped bass (3.8%F) but included bay anchovy (*Anchoa mitchilli*), Atlantic silversides (*Menidia menidia*), and American sand lance (*Ammodytes americanus*).

The importance of individual prey taxa varied among years and among dates within each year, although low sample sizes of fish with prey prevent drawing strong conclusions from these data. In 1994, striped bass fed almost exclusively on gammarid amphipods (90.2%W and 92.1%F), but these prey were less important in other years. Shrimp were important in 1993 (33.5%W, mostly *C. septemspinosa*) and 1995–97 (34.9 to 43.6%W all species). Polychaetes were observed in striped bass diets only in 1995 (23.3%W and 17.9%F). Oligochaetes were found commonly in 1997, averaging 18.5%F and 23.9%W of the diet but did not occur in other years.

## Consumption rates

Assuming an exponential evacuation pattern, consumption rate estimates for overwintering YOY striped bass ranged from 0 to 0.29% body weight per day (%bw/day, Fig. 2). Standard deviations of estimated consumption averaged 46% of the estimate (range: 20.2–100.9%). The standard deviation of the consumption estimate was relatively insensitive to assumptions of the covariance between $\overline{S}$ and $R_e$. The ratio of estimates under the least conservative assumption (perfect covariance) to that under the most conservative assumption (perfect negative covariance) was less than 1.08, with one exception (1.17 on 20 December 1993). The standard deviations presented in Figure 2 are the intermediate values, based on the assumption of no covariance.

## Feeding patterns

Among individuals, we observed a strong negative relationship between maximum gut fullness and the level of storage lipids. The 95$^{th}$ quantile of gut fullness was 3% for fish with lipid reserves of 2% dry weight and decreased to under 1% for fish with lipid levels in excess of 19% dry weight (Fig. 3). No significant patterns were observed between individual gut fullness levels and water temperature, time

of year, or body size. At the population level, only date was significantly correlated with mean gut fullness (Kendall's rank correlation: $P$=0.032; Fig. 4). Addition of temperature and average lipid level did not provide significant improvements in the fit of the model ($P$>0.20 for both).

## Discussion

### Evacuation rates

Laboratory-measured evacuation rates of YOY striped bass at representative winter temperatures were among the lowest reported for any fish species. None of the estimates compiled from multiple studies by He and Wurtsbaugh (1993) included rates as low as the 0.027 we measured at 11°C. The reason for this observation is not that striped bass have exceedingly low digestion rates compared with other fishes, but rather that evacuation rate is strongly related to temperature and is rarely measured at the lower end of temperatures encountered by each species. For example, the lowest temperature at which evacuation rates in white perch (*Morone americana*) have been measured is 12.6°C (Parrish and Margraf 1990), despite the fact that this fish often occupies waters near 0°C. The only evacuation rates we found in the literature comparable to those we measured were those for largemouth bass at 4°C ($R_e$=0.006; Cochran and Adelman, 1982, from data in Markus, 1932) and for brown trout at 1.8°C ($R_e$=0.026; Jensen and Berg, 1993), both measured near the species' thermal minima.




**Figure 2**

Consumption estimates (%body weight/day [±1 SD]) for YOY striped bass in the Hudson River estuary through five winters, and bottom water temperatures observed during sampling.

Factors other than temperature have been found to influence evacuation rates such as prey type, method of feeding (voluntary vs. force-feeding), and body size. All of our experiments were conducted with a single prey type (*C. septemspinosa*) at one ration level (2% of body weight). Meal size has been found to affect evacuation rate in laboratory studies (Smith et al., 1989; Andersen, 1998), but this effect is substantially smaller than that of temperature (He and Wurtsbaugh, 1993). Our meal size was slightly higher than the mean, but well within the range observed among individual fish. We found that body size did not affect evacuation rate over the size range of overwintering YOY striped bass (88–150 mm). In several studies comparing evacuation rates of various prey, large bodied shrimp, including *C. septemspinosa*, were found to be evacuated at lower rates than soft bodied prey or smaller shrimp species (Nelson and Ross, 1995; Singh-Renton and Bromley, 1996; Lankford and Targett, 1997), whereas other studies found no differences among prey types (Juanes and Conover, 1994; dos Santos and Jobling, 1995). Although *C. septemspinosa* and other shrimp species are common diet items of YOY striped bass, they were not the dominant item. If other prey items, such as amphipods, are digested and evacuated more rapidly, consumption rates of wild fish will be underestimated when non-shrimp prey dominate the diet.

We allowed a lag in our description of meal evacuation based on laboratory observations. Such a lag has been observed in several other studies, including both those where fish fed voluntarily (Gerald, 1973; Grove et al., 1985) and were force-fed (Vondracek, 1987). The length of the lag observed in juvenile striped bass decreased as temperature increased, as seen in juvenile turbot (*Scopthalmus maximus*; Grove et al., 1985). The lag prior to beginning of evacuation could be an artifact of experimental conditions or a natural delay in the passing of food from the stomach to the intestine following ingestion of shelled prey. Our model estimated the evacuation rate parameter ($R_e$) after the lag because this parameter represents the rate of passage of food from the stomach. Consumption rates based on gut fullness levels may be overestimated if there is a considerable lag prior to the beginning of digestion. However, the digestive lag we observed was short compared with

the total evacuation time, averaging only 11% of the time required to reach 75% digestion. Simulations that varied the length of the digestive lag time and the evacuation rate showed that for lag times in the range we observed, consumption rates were overestimated by less than 5%. If the lag observed in the laboratory experiments does not occur in the wild, estimated consumption rates are unbiased.




**Figure 3**

Relationship between gut fullness and lipid level of overwintering YOY striped bass through five winters. Points are gut fullness and lipid levels of individual fish.




**Figure 4**

Relationship between mean gut fullness (±1 SE) observed on a given date and date for collections of YOY striped bass through five winters.

## Diet

Diets of overwintering YOY striped bass in the lower Hudson River estuary were dominated by benthic invertebrates such as gammarid amphipods and several shrimp species. Juvenile striped bass do not appear to undergo a major diet shift from summer to winter but appear to focus more heavily on amphipods in winter. Studies of summer diets of striped bass in the Hudson River from the 1970s and 1990s found a slightly more diverse diet than we observed in winter. Summer diets included copepods, chironomids, and isopods, and more commonly incorporated fish and polychaetes (Gardinier and Hoff, 1982; Hurst, unpubl. data). Data from other estuaries suggest that the diets of overwintering juvenile striped bass may vary regionally. In Chesapeake Bay, Hartman and Brandt (1995a) found that fish prey accounted for 20–25%W of YOY striped bass diets in winter, substantially more than the 2–9%W we observed in the Hudson River. In the Miramichi River estuary, overwintering fish fed primarily on shrimp (mysids and *C. septemspinosa*) and ceased feeding when temperatures fell below 3°C in late November (Robichaud-LeBlanc et al., 1997). The studies from Chesapeake Bay and the Miramichi River estuary did not examine interannual variability in diets.

Overwintering juvenile striped bass appear to be opportunistic feeders, their diets reflective of the epibenthic invertebrate community in the lower Hudson River and similar to published information on the diets of co-occurring species. Although there have been no surveys of the benthic community in the lower Hudson River in winter, data available from summer surveys in the Hudson River estuary and the adjacent Raritan estuary suggest dominance of the epibenthic community by gammarid amphipods, shrimp, and annelids (Ristich et al., 1977; Steimle and Caracciolo-Ward, 1989). The winter fish community in the lower Hudson River is composed primarily of striped bass, white perch, Atlantic tomcod (*Microgadus tomcod*), and winter flounder (*Pseudopleuronectes americanus*). Winter diets of Atlantic tomcod were dominated by gammarid amphipods and copepods and were significantly less diverse in winter than in spring and autumn (Grabe, 1977, 1980). Diets of age-1 and older striped bass overwintering in the lower Hudson River estuary are similar to those of YOY fish, although the incidence of juvenile fish prey (including YOY striped bass) increased with body size (Dunning et al., 1997).

## Consumption rates

Consumption estimates of overwintering YOY striped bass in the Hudson River were consistently below 1%bw/day, with only 34% of captured fish containing prey items. Our field estimates of the consumption rates of overwintering striped bass are significantly lower than pub-

lished estimates of maximum consumption measured in laboratory experiments (Hartman and Brandt, 1995b). Their estimates of maximum consumption (for a 10-g fish) increased from 0.45%bw/day at 1°C to 7.10%bw/day at 10°C. The highest percentages of predicted maximum consumption achieved by YOY striped bass in the Hudson River occurred on 6 January 1994 (25.8%), and on three dates in early winter 1995 (15.5–18.0%). On other dates consumption was generally below 10% of maximum estimated from the Hartman and Brandt (1995b) model.

Bull et al. (1996) developed a model of consumption for overwintering juvenile Atlantic salmon in which appetite was related to anticipated metabolic requirements. To minimize the risks of starvation and predation associated with foraging, they predicted that appetite of overwintering fish should be highest in early winter (when future metabolic needs are greatest) or when internal energy reserves are low. Our results suggest that a similar model might be appropriate for overwintering striped bass. Gut fullness levels of YOY striped bass were related to level of lipid reserves at the individual level (Fig. 3). Although some fish had empty stomachs at all energy levels, the highest observed gut fullnesses were found in fish with low lipid levels. This finding suggests that overwintering striped bass increase feeding activity when energy reserves become depleted. Such feeding patterns have been documented in laboratory experiments with Atlantic salmon (Metcalfe and Thorpe, 1992) and striped bass (Hurst and Conover, 2001) but have not previously been observed among fish feeding in the wild.

At the population level, we observed a negative relationship between mean gut fullness and date; gut fullnesses were higher in early winter than late winter (Fig. 4). This pattern was predicted in the Atlantic salmon model (Bull et al., 1996) but could also be due to external factors such as depleted food resources at the end of winter. Benthic prey production is likely reduced by low winter temperatures, and standing stocks may become depleted as winter progresses. Further work is required to determine fully the causes and implications of reduced gut fullnesses observed in later winter. Reduced feeding in late winter may reflect the availability of sufficient energetic reserves and suggests that starvation is unlikely. Conversely, reduced feeding due to depressed prey availability in late winter would indicate a strong potential for winter starvation. The role of starvation in winter mortality will depend greatly on determining if variable feeding patterns among years are due to internally controlled variations in feeding motivation or environmentally imposed constraints on prey availability.

## Acknowledgments

We thank J. Powers for assistance with stomach content and lipid analyses. Significant assistance with field collections was provided by the New York Power Authority and the crews of the RV *Pannaway* and the RV *Heather M II*. S. Munch provided advice on data analysis. R. Cowen, R. Cerrato, D. Lonsdale, E. Houde, and two anonymous reviewers provided valuable comments on this manuscript. This work was funded by a graduate fellowship in population biology from the Electric Power Research Institute (T.P. Hurst) and by research grants from the Hudson River Foundation for Science and Environmental Research and the National Science Foundation (OCE9530209 to D.O. Conover).

## Literature cited

Andersen, N. G.
1998. The effect of meal size on gastric evacuation in whiting. J. Fish Biol. 52:743–755.

Boreman, J., and H. M. Austin.
1985. Production and harvest of anadromous striped bass stocks along the Atlantic coast. Trans. Am. Fish. Soc. 114:3–7.

Boynton, W. R., T. T. Polgar, and H. H. Zion.
1981. Importance of juvenile striped bass food habits in the Potomac estuary. Trans. Am. Fish. Soc. 110:56–63.

Bromley, P. J.
1994. The role of gastric evacuation experiments in quantifying the feeding rates of predatory fish. Rev. Fish Biol. Fish. 4:33–66.

Bull, C. D., N. B. Metcalfe, and M. Mangel.
1996. Seasonal matching of foraging to anticipated energy requirements in anorexic juvenile salmon. Proc. Royal Soc. London Series B, 263:13–18.

Cochran, P. A., and I. R. Adelman.
1982. Seasonal aspects of daily ration and diet of largemouth bass, *Micropterus salmoides*, with an evaluation of gastric evacuation rates. Environ. Biol. Fishes 7:265–275.

Cunjak, R. A., and G. Power.
1987. The feeding and energetics of stream-resident trout in winter. J. Fish Biol. 31:493–511.

Daiber, F.
1956. A comparative analysis of the winter feeding habits of two benthic stream fishes. Copeia 1956:141–151.

Diana, J. S.
1979. The feeding pattern and daily ration of a top carnivore, the northern pike (*Esox lucius*). Can. J. Zool. 57: 2121–2127.

dos Santos, J., and M. Jobling.
1995. Test of a food consumption model for the Atlantic cod. ICES J. Mar. Sci. 52:209–219.

Dovel, W. L.
1992. Movements of immature striped bass in the Hudson estuary: estuarine research in the 1980's. State Univ. New York Press, Albany, NY, p. 276–300.

Dunning, D. J., J. R. Waldman, Q. E. Ross, and M. T. Mattson.
1997. Use of Atlantic tomcod and other prey by striped bass in the lower Hudson River estuary during winter. Trans. Am. Fish. Soc. 126:857–861.

Eggers, D. M.
1979. Comments on some recent methods for estimating food consumption by fish. J. Fish. Res. Board Can. 36: 1018–1020.

Elliott, J. M., and L. Persson.
1978. The estimation of daily rates of food consumption for fish. J. Anim. Ecol. 1978:977–991.

Foltz, J. W., and C. R. Norden.
1977. Seasonal changes in food consumption and energy content of smelt (*Osmerus mordax*) in Lake Michigan. Trans. Am. Fish. Soc. 106:230–234.

Foy, R. J., and A. J. Paul.
1999. Winter feeding and changes in somatic energy content

of age-0 Pacific herring in Prince William Sound, Alaska. Trans. Am. Fish. Soc. 128:1193–1200.

Gardinier, M. N., and T. B. Hoff.
1982. Diet of striped bass in the Hudson River estuary. N.Y. Fish Game J. 29:152–165.

Gerald, V. M.
1973. Rate of digestion in *Ophiocephalus punctatus* Bloch. Comp. Biochem. Physiol. 46A:195–205.

Grabe, S. A.
1977. Food and feeding habits of juvenile Atlantic tomcod, *Microgadus tomcod*, from Haverstraw Bay, Hudson River. Fish. Bull. 76:89–94.
1980. Food of age 1 and 2 Atlantic tomcod, *Microgadus tomcod*, from Haverstraw Bay, Hudson River, New York. Fish. Bull. 77:1003–1006.

Grove, D. J., M. A. Moctezuma, H. R. J. Flett, J. S. Foott, T. Watson, and M. W. Flowedew.
1985. Gastric emptying and the return of appetite in juvenile turbot, *Scopthalmus maximus* L., fed on artificial diets. J. Fish Biol. 26:339–354.

Hartman, K. J., and S. B. Brandt.
1995a. Trophic resource partitioning, diets, and growth of sympatric estuarine predators. Trans. Am. Fish. Soc. 124:520–537.
1995b. Comparative energetics and the development of bioenergetics models for sympatric estuarine piscivores. Can. J. Fish. Aquat. Sci. 52:1647–1666.

He, E., and W. A. Wurtsbaugh.
1993. An empirical model of gastric evacuation rates for fish and an analysis of digestion in piscivorous brown trout. Trans. Am. Fish. Soc. 122:717–730.

Hurst, T. P., and D. O. Conover.
1998. Winter mortality of young-of-the-year Hudson River striped bass (*Morone saxatilis*): size-dependent patterns and effects on recruitment. Can. J. Fish. Aquat. Sci. 55:1122–1130.
2001. Activity-related constraints on overwintering young-of-the-year striped bass (*Morone saxatilis*). Can. J. Zool. 79:129–136.

Hurst, T. P., E. T. Schultz, and D. O. Conover.
2000. Seasonal energy dynamics of young-of-the-year Hudson River striped bass. Trans. Am. Fish. Soc. 129:145–157.

Jensen, J. W., and T. Berg.
1993. Food rations and rate of gastric emptying in brown trout fed pellets. Prog. Fish-Cult. 55:244–249.

Juanes, F., and D O. Conover.
1994. Rapid growth, high feeding rates, and early piscivory in young-of-the-year bluefish (*Pomatomus saltatrix*). Can. J. Fish. Aquat. Sci. 51:1752–1761.

Keast, A.
1968. Feeding of some Great Lakes fishes at low temperatures. J. Fish. Res. Board Can. 25:1199–1218.

Lankford, T. E., and T. E. Targett.
1997. Selective predation by juvenile weakfish: post-consumptive constraints on energy maximization and growth. Ecology 75:1049–1061.

Markle, D. F., G. C. Grant.
1970. The summer food habits of young-of-the-year striped bass in three Virginia rivers. Chesapeake Sci. 11:50–54.

Markus, H. C.
1932. The extent to which temperature changes influence food consumption in largemouth bass. Trans. Am. Fish. Soc. 62:202–210.

Metcalfe, N. B., and J. E. Thorpe.
1992. Anorexia and defended energy levels in over-wintering juvenile salmon. J. Anim. Ecol. 61:175–181.

Minton, J. W., and R. B. McLean.
1982. Measurement of growth and consumption of sauger (*Stizostedion canadense*): implications for fish energetics studies. Can. J. Fish. Aquat. Sci. 39:1396–1403.

Nelson, G. A., and M. R. Ross.
1995. Gastric evacuation in little skate. J. Fish Biol. 46: 977–986.

Parrish, D. L., and F. J. Margraf.
1990. Gastric evacuation estimates of white perch, *Morone americana*, determined from laboratory and field data. Environ. Biol. Fishes 29:155–158.

Post, J. R., and D. O. Evans.
1989. Size-dependent overwinter mortality of young-of-the-year yellow perch (*Perca flavescens*): laboratory, in situ, and field experiments. Can. J. Fish. Aquat. Sci. 46:1958–1968.

Ristich, S. S., M. Crandall, and J. Fortier.
1977. Benthic and epibenthic macroinvertebrates of the Hudson River. I. Distribution, natural history and community structure. Estuar. Coast. Mar. Sci. 5:255–266.

Robichaud-LeBlanc, K. A., S. C. Courtenay, and J. M. Hanson.
1997. Ontogentetic diet shifts in age-0 striped bass, *Morone saxatilis*, from the Miramichi River estuary, Gulf of St. Lawrence. Can. J. Zool. 75:1300–1309.

Rulifson, R. A., and S. A. McKenna.
1987. Food of striped bass in the upper Bay of Fundy, Canada. Trans. Am. Fish. Soc. 116:119–122.

Scharf, F. S., F. Juanes, and M. Sutherland.
1998. Inferring ecological relationships from the edges of scatter diagrams: comparisons of regression techniques. Ecology 79:448–460.

Schultz, E. T., and D. O. Conover.
1997. Latitudinal differences in somatic energy storage: adaptive responses to seasonality in an estuarine fish (*Atherinidae: Menidia menidia*). Oecologia 109:516–529.

Singh-Renton, S., and P. J. Bromley.
1996. Effects of temperature, prey type and prey size on gastric evacuation in small cod and whiting. J. Fish Biol. 49:702–713.

Smith, R. L., J. M. Paul, and A. J. Paul.
1989. Gastric evacuation in walleye pollock, *Theragra chalcogramma*. Can. J. Fish. Aquat. Sci. 46:489–493.

Sogard, S. M.
1997. Size-selective mortality in the juvenile stage of teleost fishes: a review. Bull. Mar. Sci. 60:1129–1157.

Sokal, R. R., and F. J. Rohlf.
1981. Biometry, second ed. W. H. Freeman and Co., New York, NY, 859 p.

Steimle, F. W., and J. Caracciolo-Ward.
1989. A reassessment of the status of the benthic macrofauna of the Raritan Estuary. Estuaries 12:145–156.

Thompson, J. T., E. P. Bergersen, C. A. Carlson, and L. R. Kaeding.
1991. Role of size, condition, and lipid content in the overwinter survival of age-0 Colorado squawfish. Trans. Am. Fish. Soc. 120:346–353.

Toneys, M. L., and D. W. Coble.
1980. Mortality, hematocrit, osmolality, electrolyte regulation, and fat depletion of young-of-the-year freshwater fishes under simulated winter conditions. Can. J. Fish. Aquat. Sci. 37:225–232.

Vondracek, B.
1987. Digestion rates and gastric evacuation times in relation to temperature of the Sacramento squawfish, *Ptychocheilus grandis*. Fish. Bull. 85:159–163.

# Use of a continuous egg sampler for ichthyoplankton surveys: application to the estimation of daily egg production of Pacific sardine (*Sardinops sagax*) off California

**Nancy C. H. Lo**
**John R. Hunter**
**Richard Charter**

Southwest Fisheries Science Center
National Marine Fisheries Service, NOAA
8604 La Jolla Shores
La Jolla, California 92037
E-mail address (for N. C. H. Lo): Nancy.Lo@noaa.gov

**Abstract**–In 1996, using the continuous underway fish egg sampler (CUFES), we carried out a preliminary study with the daily egg production method (DEPM) for estimating fish biomass of Pacific sardine, *Sardinops sagax*. Full-water-column abundance of sardine eggs was correlated with the abundance of eggs taken at 3-m depth with a CUFES; however direct conversion of CUFES samples to the full-water-column abundance, required by the DEPM, would add considerable variance to an estimate of daily egg production. Our preliminary study also indicated that the average size of an egg patch for Pacific sardine was 22 km diameter and that all stages of sardine eggs were not equally vulnerable to a CUFES at 3-m depth.

Using these findings as a guide, in 1997 we carried out an adaptive allocation DEPM survey in which the numbers of sardine eggs collected with the CUFES were used to determine subsequent locations of vertical net tows. All the vertical net tows were taken in a high-density stratum where the CUFES collected at least two eggs per minute; egg density in this stratum was computed by using only vertical tows. The remaining survey area, where the CUFES collected fewer than two eggs per minute, constituted a low-density stratum where egg density was estimated by using the ratio of CUFES egg abundance in these two strata multiplied by the egg density in the high-density stratum. Conventional statistics were used because the sampling units were survey lines spaced at 22-km or greater intervals. An acceptable level of precision (CV=21%) for a daily production of 2.57 eggs 0.05 $m^2$/d (51.4 eggs $m^2$/d), was achieved by using only 141 vertical net tows. Therefore, the CUFES will enhance the DEPM by increasing precision of the estimates, or by reducing costs in relation to a survey composed of only vertical water-column tows, when the CUFES is used adaptively to establish sampling strata for vertical water column tows.

Manuscript accepted 19 March 2001.
Fish. Bull. 99:554–571 (2001).

The continuous underway fish egg sampler (CUFES) (Checkley et al., 1997; van der Lingen et al., 1998; Checkley et al., 1999; Watson et al., 1999) is a new device that provides high-resolution spatial maps of fish eggs by sieving the eggs from water pumped from a fixed depth while a survey ship is underway. These data may be used as an index of fish abundance, or to study spawning habitats, and if converted to the numbers of eggs in the full-water-column, they may be used to estimate fish biomass by using one of the egg production methods (Hunter and Lo, 1997).

The objective of our study was to evaluate the use of the CUFES in the daily egg production method (DEPM) of estimating the spawning biomass of pelagic fishes (Lasker et al., 1985; Parker, 1985). In the DEPM, biomass is calculated from the number of staged eggs taken in plankton samples and the daily fecundity of the parents. In a standard DEPM estimate, eggs are sampled in vertical net tows, starting from a point below the maximum depth of the eggs (typically 70 m) and ending at the surface. These vertical samples are taken within a grid of stations located 4 nmi apart, a sampling interval known to produce uncorrelated samples of anchovy eggs (Smith and Hewitt, 1985). Use of the CUFES in the DEPM has several potential advantages over the use of a standard fixed-grid survey. Continuous sampling may increase the precision of the biomass estimate because it provides an increased spatial resolution of egg patches (Hunter and Lo, 1997). Continuous sampling may save ship time and thereby reduce the cost of sampling per transect mile, and the increased spatial resolution of egg patches provides new knowledge regarding the spawning behavior of the species. Several potential disadvantages of using a CUFES in the DEPM also exist: the gains in precision provided by continuous sampling may be diminished because of the increases in variance due to converting numbers of eggs taken in a CUFES to a full-water-column abundance; a much more complicated formula for variance of the estimate may be needed because of correlated CUFES samples; and the numbers of staged eggs taken in a CUFES may be biased, either because a CUFES damages eggs, making the staging of them more subject to error or because all stages may not be equally vulnerable to sampling at the 3-m depth. Clearly these issues need to be resolved if a CUFES is to be used in a DEPM survey.

We used the following approach to evaluate the use of a CUFES in the estimation of daily egg production. We conducted a pilot CUFES survey in 1996 with the objective of examining the spatial properties of Pacific sardine (*Sardinops sagax*) eggs sampled by a CUFES. We determined how the numbers of eggs taken in a CUFES were related to their abundance in the full-water-column, the spatial correlation of egg samples, and the condition of eggs after passing through a CUFES. Our

**Table 1**

Total number of samples and positive samples of sardine eggs sampled in CUFES and CalVET surveys. The range and mean of duration (minutes) for CUFES collections for 1994, 1996, and 1997 surveys.

| | | 1996 pilot CUFES survey | | 1997 DEPM survey | |
|---|---|---|---|---|---|
| | 1994 (18 Apr—11 May) | leg 1 (15–21 Mar) | leg 2 (21 Mar–6 Apr) | leg 1 (11 Mar–27 Mar) | leg 2 (28 Mar–7 Apr) |
| CUFES[1] (positive) | — | 1396 (889) | 905 (568) | 896 (550) | 331 (137) |
| Duration mean range (min) | — | 3.5 2–5 | 18.13 3–35 | 30 1–54 | 26.9 1–34 |
| CalVET[2] (positive) | 684 (74) | 91 (66) | — — | 141 (102) | — — |
| Survey area ($km^2$) | 380,175 (253,850 in USA) | | 157,000 | 174,196 | |

[1] Total collection in leg 1, 1996, was 1437. The first 41 tows were experimental.

[2] CalVET surveys in 1996 and 1997 were taken in the high-density stratum; in 1994 they were taken in a fixed grid over the whole survey area.

analysis of the 1996 cruise was used to develop an adaptive allocation survey design, similar to that proposed by Thompson et al. (1992), to estimate the biomass of sardine that incorporates a CUFES into the DEPM. Using the adaptive allocation survey design, we conducted a DEPM survey for Pacific sardine with a CUFES in 1997, and the precision of our estimate of daily production of eggs was compared with the precision of the estimate provided by a DEPM survey carried out in 1994 under standard methods (Lo et al., 1996). We also considered the accuracy of shipboard counts of eggs collected in a CUFES in relation to counts of preserved samples made after cruise end. We present our results in chronological order, beginning with those from the 1996 pilot survey which provided the rational for the new survey design in 1997; we next describe the new survey design, and end with a comparison of the 1997 CUFES and DEPM survey with the conventional DEPM survey in 1994.

## Survey data and spatial models

### Survey data

The data used in our study were taken from three ichthyoplankton surveys (Table 1):

1 A pilot CUFES cruise in 1996. This cruise consisted of leg 1 (during which both a CUFES was used and full-water-column tows were taken) and leg 2 (during which only a CUFES was used to survey the large geographic area of spawning sardine (Fig. 1);
2 A DEPM survey in 1997. This survey for Pacific sardine employed a new survey design with the CUFES and the California vertical tow (CalVET, see below) (Smith et al., 1985) (Fig. 2). The allocation of CalVETs was determined by the egg density observed from the CUFES.
3 Results of the 1994 DEPM survey off California, U.S., and Baja California, Mexico (Lo et al., 1996). This was a conventional fixed-grid DEPM survey employing only CalVETs. This cruise was used as a standard for comparing DEPM surveys with and without the CUFES.

The CalVET net consisted of 150-μm nylon netting. The diameter of the CalVET net frame was 25 cm; the tow was lowered to a depth of 70 m and was retrieved vertically. The CUFES was installed midship on the NOAA vessel *David Starr Jordan* onto the intake pipe over the side of the vessel; it extended 3 m below the water surface (see illustration in Checkley et al., 1997). Eggs were sieved from the water flow with the 500-μm nylon mesh of the CUFES concentrator.

The density of eggs taken in the CalVET net was expressed as the number of eggs/0.05 $m^2$ of sea surface water, a standard procedure in the DEPM, where 0.05 $m^2$ is the area of the mouth opening of the CalVET net. All eggs taken in the CalVET samples, regardless of survey, were counted and staged in the laboratory. The density of eggs taken in the CUFES was expressed as the number of eggs taken per minute. The interval over which eggs accumulated in CUFES samples varied depending on their abundance. When abundance was low, samples were collected over intervals of 0.5 h equivalent to 7.4 km or 4 nmi on the transect line; when abundance was high, they were collected over intervals of at least 1 minute (0.24 km or 0.13 nmi). All CUFES samples were counted at sea, preserved, and recounted in the laboratory. All eggs taken by the CUFES in the pilot survey, but not the subsequent DEPM survey, were staged in the laboratory. We used the system detailed in Lo et al. (1996) and grouped eggs by their ages into half-day age classes for egg mortality computation and one-day age classes for spatial statistical analysis.

In our study, we considered the estimates of only the daily production of eggs, $P_0$, one of the key parameters in




**Figure 1**

Sardine eggs/minute from CUFES samples and survey pattern in cruise 9603. Leg 1: 13–21 March 1996 (**A**) and leg 2 (**B**): 6–21 March 1996. The star on the lower right corner of section B is the reference point (0,0) to compute variograms (see text).




**Figure 2**

Sardine eggs/minute from CUFES samples and survey pattern in cruise 9703, 11 March–6 April 1997, with two strata: a high-density stratum (open area) and a low-density stratum (shaded area).

the DEPM. $P_0$ is only one of six parameters used in estimating biomass with the DEPM. In the DEPM, biomass is related to egg production using the model $B_s = P_0 A / (R / W_f) SF$, where $B_s$ = biomass for area $A$; $P_0$ = the daily egg production at age 0 day per unit sea surface area; $W_f$ = the average female weight; $S$ = the fraction of females spawning per day; $F$ =batch fecundity; and $R$ = the fraction of the biomass that is female (Alheit, 1993; Hunter and Lo, 1997). The denominator, $(R / W_f)SF$, is the number of eggs/biomass in grams and is also called the daily specific fecundity. In the 1994 survey all parameters were estimated but in the 1997 survey we estimated only $P_0$ and $A$ and used historical data for the other parameters (Hill et al.[1]).

## Variograms of sardine density (eggs/minute)

As the first step in developing a DEPM survey design for the CUFES, we used geostatistical techniques (Cressie, 1991; Barange and Hampton, 1997; and Fletcher and Sumner, 1999) to describe the spatial structure of the egg distribution and estimated the major diameters of sardines egg patches from the 1996 survey. We applied variogram models (Cressie, 1991; Petitgas, 1993) to CUFE samples (in units of eggs/minute) grouped into three age groups: 1-day (4–27 h), 2-day (28–51 h), and 3-day (52–75 h).

Variogram ($\gamma(h)$) is defined as the variance of difference between values that are $h$ units apart and is a function of variance and covariance:

$$\begin{aligned} 2\gamma(h) &= \mathrm{var}\left[u(x) - u(x+h)\right] \\ &= 2\left(\mathrm{var}(u) - \mathrm{cov}(h)\right) \qquad \text{if } \mathrm{var}\left[u(x)\right] = \mathrm{var}\left[u(x+h)\right], \end{aligned} \tag{1}$$

where $u(x)$ = the eggs/min at location $x$;
$u(x+h)$ = the eggs/min at the location $h$ (nmi) away from $x$;
$\mathrm{var}(u)$ = the variance; and
$\mathrm{cov}(h)$ = the covariance of eggs/min that are $h$ (nmi) apart.

$\gamma(h)$ is the semivariogram. For simplicity, we refer to $\gamma(h)$ as the variogram. We used S+SpatialStats (Kaluzny et al.,

[1] Hill, Kevin, T. M. Yaremko, L. D. Jacobson, N. C. H. Lo, and D. A. Hannan. 1998. Stock assessment and management recommendations for Pacific sardine (*Sardinops sagax*) in 1997. Marine region, Admin. Rep. 98-5, Cal. Dep. of Fish and Game, 8604 La Jolla Shores Dr., La Jolla, CA 92037.

1996) and EVA software (Petitgas and Prampant[2]) to analyze and visualize spatial distributions of sardine eggs/min and to compute a variogram for each of the three age groups of sardine eggs. Two basic assumptions of the variogram (Eq. 1) are intrinsic stationarity and isotropy. Intrinsic stationarity means that a constant mean exists and the variance of egg density is defined by the magnitude of $h$. Isotropy means that spatial correlation and the range of correlation do not change with direction. The variogram is normally expressed as a function of three parameters: range, sill, and nugget effect. The range is the distance beyond which the observations are not correlated. The sill is the variance of the random field and is the asymptotic value of $\gamma(h)$ . The nugget effect measures the micromeasurement error and the white noise for $h$ close to 0 (Cressie, 1991).

Ideally, for $h$ close to 0, the variogram ($\gamma(h)$) will be close to zero because the observations tend to be similar. As $h$ increases, the observations become $h$ units apart and tend to be different, or cov($h$) decreases, and the variogram increases. At a certain distance, $h^*$, cov($h$) approaches zero, and for $h > h^*$, the variogram approaches its asymptote (sill). The distance, $h^*$, is the range. The range ($h^*$) was estimated from a model that best fits the data and was used to estimate the diameter of the patch of sardine eggs because eggs whose distances are less than $h^*$ nmi are correlated and thus are likely to be in the same patch. Conversely, eggs that are more than $h^*$ nmi apart are no longer correlated and thus are assumed to be in different patches. We chose the robust (or stable) estimator of the variogram (Cressie and Howkins, 1980; Cressie, 1991):

$$2\hat{\gamma}(h) = \frac{\left[\sum_{N(h)} |u(x) - u(x+h)|^{1/2}\right]^4}{\left[0.457 + 0.494/|N(h)|\right]N(h)^4}, \quad (2)$$

where $|N(h)|$ = the number of distinctive pairs; and
$h$ = the distance (nmi) between any two locations.

To avoid possible trends, we first ran a local regression model (LOESS) of ln(eggs/min+1) against line distance (the distance computed from the survey lines, $y$-axis) and station distance ($x$-axis) (Chamber and Hastie, 1992) where the reference point (0,0) is a pseudo station in Mexico (station 260 on CalCOFI line 980; Fig. 1B). We then constructed a variogram for the residuals from the local regression model (LOESS) in four directions clockwise from the transect (0° ,45°, 90°, and 135°) to examine possible anisotrophy. We chose natural logarithm (ln) transformed data because the distribution of eggs was skewed. Finally, we used an interactive S+ function (model.variogram function) to determine the estimates of parameters for each variogram: range, sill and nugget effect. The range was then used as the estimate of the diameter of the patch for each of the three age groups and total number of eggs.

[2] Petitgas, P., and A. Prampant. 1993. EVA (estimation variance), a geostatistical software on IBM-PC for structure characterization and variance computation. CM1993/D:65, 81st meeting of ICES. IFREMER laboratoire d'Ecologie Halieutique, BP 21105,44311 Nantes (France), Cedex 03, 33 p.

## Results of the 1996 pilot CUFES survey

### Spatial correlation and patch size of sardine eggs

For each of the three age groups and the total number of eggs, the four-directional variograms of the residuals of ln(eggs/min+1) from LOESS (Fig. 3) indicated that the variograms for transects at 0°, 45°, and 135° were clearer than the cross-transect (90°), particularly for the 1-day-old eggs, 2-day-old eggs, and total egg category. The variogram in the direction of within-transect (0 degree) had the clearest signal because intervals between adjacent collections were the shortest. Because the variograms for each direction looked similar, we used the variogram in the within-transect direction to assess the spatial correlation of sardine eggs.

The spherical model was chosen to fit the variogram (Cressie, 1991):

$$\gamma(h;\theta) = c_0 + c_s\left[(3/2)(\|h\|/a_s) - (1/2)(\|h\|/a_s)^3\right] \begin{array}{l} 0 \quad h = 0 \\ 0 < \|h\| < a_s \\ C_0 + C_s, \quad \|h\| \geq a_s. \end{array} \quad (3)$$

where $c_0$ = the nugget effect;
$c_s$ = the practical sill (variance – nugget);
$a_s$ = the range; and
$\|h\|$ = Euclidean distance.

For total number of eggs (all eggs combined), the range of the residuals of ln(eggs/min+1) was 22.2 km (12 nmi), sill (variance in the random field) was 0.4, and nugget was 0.05. For 1-day-old eggs, the range was 14.8 km (8 nmi) and the sill was 0.3. For 2-day-old eggs, the range was 18.5 km (10 nmi), the sill was 0.15, and nugget was 0.005, and for 3-day-old eggs, range was 22.2 km (12 nmi), the sill was 0.065, and nugget was 0.005 (Fig. 4). Because the maximum range was 22.2 km (12 nmi), eggs collected more than 22.2 km (12 nmi) apart were considered uncorrelated; therefore, transect lines spaced intervals of 12 nmi or greater were considered independent. The data also indicated the gradual dispersion of egg sardine patches with time as described by Smith (1973) because the diameter of sardine egg patches increased from 14.8 km for 4–27 h old eggs to 22.2 km for eggs 52–75 h old.

### Conversion of CUFES egg density to full-water-column abundance and distribution of egg stages

Egg counts from 91 paired samples collected with the CUFES and CalVETs during leg 1 of the1996 survey (Table 1) were used to derive a conversion factor from eggs/minute of CUFES sample to CalVET catch ($R$). We used a regression estimator to compute the ratio of eggs/minute from the CUFES to eggs/tows from CalVETs, $R = \mu_y/\mu_x$, where $y$ = eggs/minute; $x$ = eggs/tow; and $R$ = the catch ratio. The estimator of $R$ is $\hat{R} = \Sigma(x \times y)/\Sigma(x^2)$.




**Figure 3**

Variogram of residuals of ln (*sardine eggs*/minute+1) for age groups: 4–27 h (**A**), 28–51 h (**B**), and 52–75 h (**C**), and total eggs for four directions (**D**). Degree 0 is the direction perpendicular to the coastal line, degree 90 is the direction along the coastal line during leg 2 of cruise 9603. On the *x*-axis, the inner ticks are in km and the outer ticks are in nmi.

Paired samples were taken wherever high abundance of eggs appeared in samples collected with the CUFES. We obtained egg/minute = 0.73 eggs/0.05m$^2$ (CV=0.16) (Fig. 5). This means that for one egg observed from a CalVET, one would expect to see, on average, 0.73 eggs/min. Or for one egg/min from a CUFES, one would expect to see 1.5 eggs/tow. A striking difference existed between the data from the 1996 pilot survey and the full survey carried out in 1997. In 1997, the catch ratio of eggs/minute to eggs/tows was 0.25 (CV=0.08) from 110 pairs of CalVET and CUFES of which at least one sample was positive (Fig. 5). This means that one egg/tow from a CalVET tow was equivalent to approximately 0.25 egg/min from a CUFES, or one egg/minute from the CUFES was equivalent to 4 eggs/tow from a CalVET sample. The ephemeral nature of such conversion coefficients was not known to us when we developed the design for the 1997 survey. However, the variance associated with the direct 1996 conversions was a strong incentive to reduce the effects of direct conversions in the design of the 1997 survey and in the calculation of biomass.

To determine if the CUFES provides an unbiased sample of all sardine eggs stages, we compared the distributions of developmental stages between the two samplers taken in 91 paired CUFES and CalVET samples during leg 1 of the 1996 survey (Fig. 1). A 2 × 11 contingency table was constructed for total counts of each of 11 stages of eggs collected with the CalVET and the CUFES. A chi-square statistic was computed to test the null hypothesis that the distribution of stages was independent of the samplers.

The chi-square ($\chi^2$) analysis showed that the distribution of stages was not the same between the two samplers ($\chi^2$=188.47, df =10, *P*-value <0.01) (Table 2). The difference was primarily due to eggs of stages I, III, V, and VI. The CUFES caught only two stage-I eggs; therefore we decided to run a $\chi^2$ test with stage-I and stage-II eggs collapsed into one group. A similar conclusion was reached for the later case ($\chi^2$=160.31, df =9, *P*-value <0.01). A large $\chi^2$ value indicates that staged sardine eggs in the upper 3 m may not be representative of the sardine egg stages in the whole-water-column and that the egg production computed from the CUFES survey would be biased.

## Design for the 1997 CUFES and DEPM survey

The foregoing analysis of the 1996 survey data indicated that adequate correlation exists in overall egg abundance between the CUFES and the full-water-column egg samples, but a direct conversion of CUFES data to full-water-column abundance would add considerable variance to an estimate of daily egg production. The foregoing analysis




**Figure 4**

Spherical model fitted to variograms of residuals of ln (*sardine eggs*/minute+1) for three age groups and total eggs, in the direction perpendicular to the coastal line during leg 2 of cruise 9603. On the *x*-axis, the inner ticks are in km and the outer ticks are in nmi.

**Table 2**

Contingency table of sardine egg counts by developmental stages and two samplers: the CalVET and the CUFES, leg 1, 1996.

| | Egg stages | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | I | II | III | IV | V | VI | VII | VIII | IX | X | XI |
| Total count | | | | | | | | | | | |
| CalVET | 31 | 63 | 116 | 46 | 36 | 95 | 23 | 27 | 37 | 15 | 13 |
| CUFES | 2 | 74 | 466 | 44 | 148 | 104 | 103 | 22 | 75 | 21 | 14 |
| Percent | | | | | | | | | | | |
| CalVET | 6 | 13 | 23 | 9 | 7 | 9 | 5 | 5 | 7 | 3 | 3 |
| CUFES | 0 | 7 | 43 | 4 | 14 | 10 | 10 | 2 | 7 | 2 | 1 |

also indicated that a potential bias may exist if staged CUFES eggs are used to estimate daily egg production ($P_0$) because all stages of eggs may not be equally vulnerable to the CUFES. A further complication was found, in that egg samples taken in the CUFES were strongly correlated up to distances of 22.2 km, necessitating the use of geostatistics if individual samples were to be used for the estimates. We concluded from these findings that the most effective way to use the CUFES in the DEPM design was to use it to allocate full-water-column tows with a CalVET. This adaptive allocation design, similar to one proposed by Thompson et al. (1992), takes advantage of the rapid continuous monitoring capability of a CUFES, while preserving the quantitative features of the CalVET.

We decided to allocate a CalVET at 4-nmi intervals, whenever catch from the CUFES was greater than or equal to 2 eggs/min[1] When the catch from the CUFES began to decrease to fewer than 2 eggs/min, then CalVETs were stopped (Fig. 6). This critical value for the allocation of CalVETs was equivalent to slightly more than three eggs per CalVET tow and was based on the conversion factor from the pilot study of CUFES eggs/min = CalVET 0.73 eggs/tow. The CUFES was in operation continuously along each transect line, and samples were taken at an interval




**Figure 5**

Conversion coefficient: CUFES eggs/min = CalVET 0.73 eggs/tow in 1996 and CUFES eggs/min = CalVET 0.25 eggs/tow in 1997. Insert is an expansion of the graph near the origin. x = data obtained from 1996; o = data obtained from 1997.




**Figure 6**

Sardine eggs/0.05 m² from CalVET samples in cruise 9703 survey,11 March–7 April 1997. Lines (perpendicular to the coast) are the cruise tracks.

ranging from 1 to 54 minutes with a mean of 30 minutes and median of 15 minutes (Table 1). Only samples from CalVETs were used to model the sardine embryonic mortality curve, thereby avoiding a possible stage-specific bias with the CUFES.

Spacing of transect lines was also determined to some extent by samples from the CUFES. The scheduled spacing of lines was 40 nmi apart but if a high density of sardine eggs was encountered, an additional transect line was added between the two scheduled lines. As a result of this procedure, two lines were added (lines 4 and 8) in the 1997 DEPM survey (Fig. 2).

To estimate $P_0$, the survey area was then stratified into two strata: a high-density stratum where the density of eggs taken by the CUFES was at least 2 eggs/min; and a low-density stratum (Fig. 2) where only CalVET samples were taken and where the density of eggs was less than 2 eggs per minute. Within each stratum, the transect line was selected as the sampling unit both for CalVET samples and for CUFES samples. Because lines were spaced at intervals greater than 12 nmi, statistical procedures based on uncorrelated data were used (correlation should not be ignored if the distances are less than 12 nmi). We describe below the two different estimation methods used to estimate $P_0$: one for the high-density stratum and the other for the low-density stratum; we combined the two methods to estimate $P_0$ combined for the total survey area.

We used transects as the primary sampling units and applied conventional statistical methods to construct the mortality curve from CalVET samples (Armstrong, 1988). CalVETs, the secondary sampling units, were taken for the first ten transects, except transect number eight. The $P_0$ was estimated for each of two strata by using different methods, and an weighted average was obtained for the whole survey.




**Figure 7**

Sardine eggs/0.05 m² for each developmental stage in the high-density stratum, cruise 9703, 11 March–7 April 1997.

**Table 3**

Daily sardine egg and yolksac larval production and their standard error in the high-density stratum in 1997.

| | Age (day) | Daily production/ 0.05 m² | Standard error |
|---|---|---|---|
| Eggs | 0.32 | 3.29 | 1.32 |
| | 0.76 | 4.48 | 2.10 |
| | 1.17 | 4.32 | 2.05 |
| | 1.71 | 6.12 | 2.88 |
| | 1.92 | 3.21 | 1.60 |
| | 2.52 | 1.86 | 0.66 |
| Yolksac larvae | 5.62 | 0.65 | 0.16 |

## Egg production in the high-density stratum ($P_{0,1}$)

Ages of staged sardine eggs (Fig. 7) were assigned according to a temperature-dependent time-to-stage model (Lo et al., 1996). Sardine eggs were grouped in half-day categories excluding those eggs that were newly spawned during the first three hours. An average eggs-per-tow and an average age were obtained for each half-day category within each transect. A weighted average number of eggs in each half-day category was obtained where the weight is duration of the CUFES sampling interval for each transect line (Eq. 4, Table 3).

$$P_t = \frac{\sum_i P_{it} m_i}{\sum_i m_i} \qquad \mathrm{var}(P_t) = \frac{n/(n-1)\sum_i m_i^2 (P_{it} - P_t)^2}{\left(\sum_i m_i\right)^2}. \quad (4)$$

where $P_{it}$ = the eggs or yolksac larvae/0.05 m²; and
$m_i$ = the total CUFES sampling time (minutes) for the $i$th transect for $i$ = 1,..., 7, 9, and 10.

For a weighted nonlinear regression, the weight is $1/\mathrm{SE}(p_t)$.

The yolksac larval stage includes larvae from time of hatching to the time they form functional jaws, i.e. larvae ≤6 mm live length or 5 mm captured length (Zweifel and Lasker, 1976). As with the egg data, the average number of yolksac larvae per tow was calculated for each transect and the overall mean catch of yolksac larvae was computed as a weighted average. Yolksac larva production was the mean catch of larvae ≤5.00 mmi (captured size) per CalVET divided by its temperature-dependent stage duration (Zweifer and Lasker, 1976). The age of yolksac larvae was estimated by the average of minimum age and maximum age, i.e. hatching time + age of forming functional jaw)/2 (Lo et al., 1996) (Table 3).

With these data, an embryonic mortality was modeled by an exponential curve (Eq. 5) (Lo et al., 1996). The mortality curve was fitted to eggs/0.05 $m^2$ in each age class and numbers of yolksac larvae/0.05 $m^2$ with an assigned age based on observed mean water temperature. The estimates of the intercept ($P_{0,1}$) and the instantaneous mortality rate from the mortality curve were obtained by a nonlinear regression of S+ nonlinear regression function (nls()).

The exponential embryonic mortality curve is

$$P_t = P_{0,1} \exp(-zt), \quad (5)$$

where $P_t$ = a weighted average of eggs – yolksac larvae /0.05$m^2$; and

$t$ = the mean age (d) for each of 6 half-day age groups of eggs and yolksac larvae. The weights are total CUFES sampling time (minutes) for each transect (Eq. 4).

### Daily egg production in the low-density stratum ($P_{0,2}$)

Because no CalVET samples were taken in the low-density stratum, we estimated daily egg production ($P_{0,2}$) in that stratum as the product of the egg production in the high-density stratum ($P_{0,1}$) and the ratio (q) of egg/min in the low-density stratum to that in the high density stratum from the CUFES samples. Here we assumed that $q$ is the same no matter whether it was computed from eggs/min by the CUFES or from eggs/tow by the CalVET:

$$P_{0,2} = P_{0,1}\, q, \quad (6)$$

$$q = \frac{\sum_i \frac{\bar{x}_{2,i}}{\bar{x}_{1,i}} m_i}{\sum_i m_i}, \quad (7)$$

where $m_i$ = the total CUFES time (minutes) in the $i$th transect; and

$x_{ji}$ = was eggs/min in the $j$th stratum and $i$th transect.

The variance of $q$ was computed according to that of a ratio estimator (Eq. 4).

### Daily egg production for the total survey area ($P_0$)

$P_0$ was computed as a weighted average of $P_{0,1}$ and $P_{0,2}$, where

$$\begin{aligned} P_0 &= \frac{P_{0,1}A_1 + P_{0,2}A_2}{A_1 + A_2} \\ &= P_{0,1}w_1 + P_{0,2}w_2 \\ &= P_{0,1}[w_1 + qw_2] \end{aligned} \quad (8)$$

and according to Goodman (1960), the unbiased estimate of var($P_0$) is

$$v(P_0) = v(P_{0,1})(w_1 + w_2 q)^2 + P_{0,1}^2 w_2^2 v(q) - v(P_{0,1}) w_2^2 v(q), \quad (9)$$

where $w_i = \frac{A_i}{A_1 + A_2}$, $i = 1,2$, and $A_i$ = the area size.

### Simulation

Bootstrap simulations were conducted to provide the possible biases and another estimate of the standard error of daily egg production ($P_0$) and the instantaneous mortality rate ($z$) for each stratum and the entire survey area under the adaptive allocation sampling scheme. As mentioned earlier, CalVETs were taken on nine transect lines and not on line 8 and line 11. In the simulation, nine transects with CalVETs were sampled with replacement and estimation procedures described in previous section were followed. To evaluate the effect of weighting, we included weighted and unweighted nonlinear regression where the weight was the inverse of the standard error of egg production of each age group and yolksac larvae. One thousand iterations were run, and the standard deviation of 1000 estimates was the bootstrapped standard error of the estimates. Bias was the difference between the average of 1000 estimates and the estimate from the original data. The bias-corrected estimate was the original estimate minus the bias.

## Results of the 1997 CUFES and DEPM survey

### Daily egg production

The daily egg production for each half-day category and yolksac larval production and their ages (d) were used to construct an embryonic mortality curve for the high-density stratum (Eq. 5, Fig. 8, Table 3). The daily egg production in the high-density stratum ($P_{0,1}$) based on unweighted nonlinear regression was 5.04 eggs/0.05 $m^2$/d (100.8 eggs/$m^2$/d ,CV=0.25) and egg mortality was $z$=0.21 (CV=0.73) for an area ($A_1$) of 66,841 $km^2$ (19,530 $nmi^2$) (Eq. 8, Fig. 9). The ratio ($q$) of egg density between the low-density stratum and high-density stratum from CUFES samples was 0.211 (CV=0.43) (Eq. 7). Therefore, in the low-density stratum, the egg production ($P_{0,2}$) was 1.064 eggs/0.05 $m^2$/d (21.28 eggs/$m^2$/d, CV=0.49) for an area ($A_2$) of 107,255 $km^2$ (31,338 $nmi^2$). The estimate of the daily egg production for the entire survey area was 2.57/0.05 $m^2$ (51.4/$m^2$, CV=0.27) (Eq. 8, Table 4). The weighted nonlinear regression produced estimates slightly different from those with unweighted nonlinear regression: $P_{0,1}$ =4.76/0.05 $m^2$ (59.2/$m^2$, CV=0.18), $z$ = 0.35(CV=0.14), and the $P_0$ for the entire survey area was 2.43/0.05 $m^2$ (48.6/$m^2$, CV=0.21).

The bootstrap estimate of $P_0$ (5.10) was similar to the original estimate $P_0$ (5.04) in the high-density stratum. The standard error of $P_0$ (1.6) from the bootstrap analysis was higher than the estimate from the original data (1.28). The bias of $P_0$ (0.06) was negligible because the ratio




**Figure 8**

Embryonic mortality curve of sardine eggs and yolksac larvae in the high-density stratum, 9703. The last data point represents yolksac larvae (Table 3).




**Figure 9**

Relationship between egg counts at sea and those taken in the laboratory from CUFES samples, cruise 9703. The diagonal line is the fitted regression line: egg counts at sea = 0.8 egg counts in laboratory.

of bias to standard error was less than 0.25 (Efron and Tibshirani, 1993). However, the daily instantaneous mortality rate ($z$=0.29) from the bootstrap was higher than 0.21 from the original data. The bias-corrected $z$ was 0.13 (SE=0.12). For the entire survey area, the bootstrapped $P_0$ was 2.60/0.05 m$^2$ (SE=0.71; CV=0.27), similar to the original estimate (2.57/0.05 m$^2$).

The bootstrap results indicated that the weighted nonlinear regression produced a downward biased estimate of $P_0$ (bias=–1.26=3.50–4.76) in the high-density stratum. The standard error (1.60) of $P_0$ from bootstrap was also higher than that computed from the original data (0.86). The fact that the ratio of bias to standard error (1.26:1.60) was greater than 0.25 may indicate that the weighted non-

**Table 4**

Estimates of daily egg production ($P_0$: number of eggs/0.05 m²/day at age 0), daily instantaneous mortality rate ($z$) for the high-density stratum (66,841 km²) and low-density stratum (107,255 km²), $P_0$ for the entire survey area (174,096 km²), the catch ratio of eggs/min in the low-density stratum to eggs/min in the high-density stratum ($q$), their standard errors (SE), estimated bias, and the bias-corrected estimates for the unweighted and weighted nonlinear regression from 1000 iterations of bootstrap simulation.

| | High-density stratum | | | | Low-density stratum | | Entire survey area | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | $P_{0,1}$ | SE ($P_{0,1}$) | $z$ | SE ($z$) | $P_{0,2}$ | SE ($P_{0,2}$) | $P_0$ | SE ($P_0$) | $q$ | SE ($q$) |
| Unweighted nonlinear regression | | | | | | | | | | |
| survey | 5.04 | 1.28 | –0.21 | 0.16 | 1.06 | 0.52 | 2.57 | 0.72 | 0.211 | 0.09 |
| bootstrap | | | | | | | | | | |
| mean | 5.10 | 1.60 | –0.29 | 0.28 | 1.10 | 0.53 | 2.60 | 0.85 | 0.22 | 0.08 |
| SE | 1.60 | 0.72 | 0.12 | 0.11 | 0.48 | 0.28 | 0.71 | 0.34 | 0.09 | 0.03 |
| CV | 0.31 | 0.46 | –0.40 | 0.38 | 0.45 | 0.52 | 0.27 | 0.41 | 0.41 | 0.40 |
| bias | 0.06 | 0.32 | –0.08 | 0.12 | 0.03 | 0.01 | 0.03 | 0.13 | 0.009 | –0.01 |
| bias corrected | 4.98 | 0.96 | –0.13 | 0.04 | 1.04 | 0.51 | 2.54 | 0.58 | 0.202 | 0.10 |
| Weighted nonlinear regression | | | | | | | | | | |
| survey | 4.76 | 0.86 | –0.35 | 0.050 | 1.004 | 0.45 | 2.43 | 0.51 | 0.211 | 0.09 |
| bootstrap | | | | | | | | | | |
| mean | 3.50 | 0.72 | –0.27 | 0.056 | 0.730 | 0.33 | 1.80 | 0.41 | 0.22 | 0.08 |
| SE | 1.60 | 0.35 | 0.10 | 0.025 | 0.400 | 0.21 | 0.76 | 0.20 | 0.09 | 0.03 |
| CV | 0.46 | 0.49 | –0.36 | 0.440 | 0.550 | 0.63 | 0.42 | 0.48 | 0.42 | 0.40 |
| bias | –1.26 | –0.14 | 0.08 | 0.006 | –0.274 | –0.12 | –0.63 | –0.10 | 0.009 | 0.01 |
| bias corrected | 6.02 | 1.00 | –0.43 | 0.044 | 1.278 | 0.57 | 3.06 | 0.61 | 0.202 | 0.10 |

linear regression was unwarranted in our case. One possible reason is that the standard errors (SE) of egg production in each age group were between 1.3 and 2.8, whereas the SE for yolksac larvae was 0.16 (Table 3). Although the weighted regression should be used when the variances of data are unequal, we believe that there is too large a disparity between variance of eggs and yolksac larvae, and too much weight was assigned to yolksac larvae for a weighted nonlinear regression.

The spawning biomass was computed from the daily egg production from the 1997 survey, and the historical daily specific fecundity (number of eggs/gram of biomass) was 23.55 (Macewicz et al., 1996) assuming the daily specific fecundity was the same as that for 1994–96. Sardine spawning biomass in 1997 would be 379,940 t[3] for an area of 174,096 km² (50,868 nmi²) from San Diego to San Francisco. No variance of $B_s$ was computed because no variance of the number of eggs per population weight (g)/day was available.

## Comparison of results from the 1997 CUFES and DEPM survey with results from a conventional DEPM survey

We compared the results of the 1997 CUFES and DEPM design with those of a conventional DEPM (1994), in which only CalVET samples were taken, to illustrate how the CUFES allocation design may affect the performance of a DEPM survey of Pacific sardine. We believe the comparison is instructive, even though the two surveys differed somewhat in area and population size; the conventional 1994 survey covered a larger area than the 1997 survey (380,175 km² vs. 174,096 km²), and, the total biomass of sardine was smaller in 1994 than in 1997 (111,493 t (Lo et. al., 1996) vs. 379,940 t, respectively). In both surveys staged eggs and yolksac larvae from CalVET samples were used in the calculation of $P_0$.

An obvious difference in the results of the two surveys was that only 11% (74/684) of CalVET samples were positive for sardine eggs in the 1994 conventional survey, whereas in the 1997 survey 72% (102/141) were positive (Table 5). These results indicate that CUFES was effective in allocating CalVET samples and thereby reducing ship-time costs per survey mile. The coefficients of variation (CV) for the estimates of $P_0$ were similar: 0.22 for the conventional survey compared with 0.27 for the CUFES-based DEPM. Thus, variance penalty for using the ratio estimator $q$ did not greatly diminish the benefit in using the CUFES in the DEPM. This simple statistical comparison, however, does not reveal the greatest potential benefits in using a CUFES. The allocation of CalVETs would be most useful when the population is at a lower level, as it was in 1994, because at such levels one must cover a large survey area to assure an unbiased estimate, but the population is probably concentrated in a very small fraction of the area where CalVET samples will be allocated. In addition, the high-resolution maps of the spatial distribution

[3] The spawning biomass would be 359,280 metric tons if $P_0$ of 2.43/0.05 m² from weighted nonlinear regression was used.

**Table 5**

Sardine daily egg production ($P_0$) from a conventional survey (1994), compared with a CUFES and DEPM survey (1997).

| 1994 Conventional DEPM survey Area: 380,175 km$^2$ | | 1997 CUFES and DEPM survey Area: 174,096 km$^2$ | |
|---|---|---|---|
| CalVET samples | | CalVET samples | |
| total | 684 | total | 141 |
| positive for eggs | 72 | positive for eggs | 102 |
| positive percent | 11% | percent positive | 72% |
| CUFES samples | None | CUFES samples | |
| | | total | 1227 |
| | | total positive | 687 |
| | | percent positive | 56% |
| | | high-density stratum | 84% |
| | | low-density stratum | 40% |
| Daily egg production | | Daily egg production | |
| $P_0$ | 0.169/0.05 m$^2$ | $P_0$ | 2.57/0.05 m$^2$ |
| CV | 0.22 | CV | 0.27 |
| spawning biomass | 111,493 t | spawning biomass | 379,940 t |

of eggs provided by the CUFES have not as yet been incorporated into the DEPM design, but we believe in the long-term it will be possible to improve the accuracy of surveys by doing so, as well as possible to develope new insights into the processes involved in selection of spawning habitats by parent fish.

## Comparison of shipboard egg counts with preserved egg counts

A key element of the allocation design was that the allocation of CalVET samples was based on near real-time counts and on the identification of eggs by CUFES operators. After having been counted by CUFES operators at sea, the eggs were preserved in vials, and later recounted and identified in the laboratory by experts. We compared egg counts in the laboratory with those taken at sea from the 1997 survey to determine the reliability of shipboard counting and identification and to indicate the extent to which a difference affected the final estimate of the daily egg production.

Within each stratum, we first computed an overall mean eggs/min for the laboratory and ship count by using a ratio estimator (where $y$=total number of eggs and $x$=the total minutes for each transect summed over all transects) (Table 6). We also computed a ratio of eggs/min from the laboratory to that from the ship for each transect line and obtained a weighted ratio for each stratum, where weight was the duration for each transect within a stratum (Eq. 7). The ratio for the entire survey area was a weighted average of two ratios, one from each stratum, and weight was the area of each stratum, as in Equation 5.

In the high-density stratum, the ratio from the laboratory counts to the ship counts was 1.19:1 (CV=0.03). In the low-density stratum, the ratio from the laboratory to the ship was 1.22:1 (CV=0.05) (Table 6). The overall ratio for the entire survey area was 1.20:1 (CV=0.10); therefore the laboratory count was higher than the ship count by 20% (Table 6).

Of a total 1227 collections, 687 were positive according to laboratory counts. There were 16 collections where ship counts were positive but counts in the laboratory were zero. Eggs of other species in those 16 CUFES collections were obviously misidentified as sardine eggs. (Table 6). A total of 658 pairs had positive counts, out of which 130 pairs had equal positive counts. A total of 524 pairs had zero counts. Therefore there were 654 equal counts between laboratory and ship (524 zeros and 130 positive counts) and 573 pairs (=1227–654) with a mismatch. The relationship between the total counts from the laboratory and the ship confirmed the undercount from the ship (Fig. 9). The variance of undercounts increased with the total counts from the laboratory. One outlier was a collection where the ship count was zero and the laboratory count was 341. Although the absolute undercounts increased with the total number of eggs, the percent of undercount decreased with the total egg count.

## Conversion coefficient between two strata ($q$)

In the 1997 CUFES and DEPM survey, one of the primary functions of the CUFES collections was to provide a conversion coefficient ($q$) of egg density between two strata to convert the egg production in the high-density stratum ($P_{0,1}$) to the low-density stratum ($P_{0,2}$), i.e. $P_{0,2} = P_{0,1} \times q$ (Eq. 6). The conversion coefficients ($q$) computed from the laboratory and ship counts were similar: 0.213 (CV=0.44) from the laboratory and 0.211 (CV=0.43) from the ship counts. Therefore, the bias of using egg counts at sea to calibrate egg production per unit area in the low-density stratum would be small: if the ratio of 0.213:1 were used, the egg production for the entire survey area would be 2.60/0.05 m$^2$ instead of 2.57/0.05 m.$^2$

**Table 6**

Comparison of egg density and distribution of zero and positive egg counts from CUFES samples with those taken in the laboratory and at sea (ship), 1997. CVs are in shown in parentheses.

| Egg density (eggs/min) | High-density stratum | Low-density stratum | Overall | $q$ |
|---|---|---|---|---|
| ship | 4.16 (0.41) | 0.47 (0.45) | | 0.213 (0.44) |
| laboratory | 4.91 (0.4) | 0.57 (0.45) | | 0.211 (0.43) |
| ratio | 1.19 (0.03) | 1.22 (0.05) | 1.20 (0.10) | |

Distribution of zero and positive egg counts

| | Laboratory | | |
|---|---|---|---|
| Ship | zero | positive | total |
| zero | 524 | 29 | 553 |
| positive | 16 | 658[1] | 674 |
| total | 540 | 687 | 1,227 |

[1] 658 = 130 with exact match + 428 without match.

## Discussion

Presently the most productive use of the CUFES in the DEPM is to use the CUFES as an efficient auxiliary information provider and the CalVET, or another full-water-column tow, as the primary sampler. Given our present level of knowledge of egg distributions and accuracy of predicting them, it would be folly to depend only on the CUFES for an estimate of $P_0$. Predictive egg-distribution models (Sundby, 1983; Westgard, 1989) are promising and sometime in the future might make exclusive use of the CUFES practical in a DEPM. Exclusive use of the CUFES as a sampler in the DEPM also would require detailed knowledge of the stage-specific vulnerability of eggs to the sampler because a bias seemed to exist for sardine. In short, full-water-column tows are essential for accurately estimating egg production today, but the CUFES can greatly facilitate the allocation of such tows.

The CUFES enhancement of the basic DEPM design (Lasker, 1985) may have broad application because the DEPM is used world-wide for estimating the spawning biomass of sardines, anchovies, and other species of fishes (Priede and Watson, 1993; Zeldis, 1993; Lo, 1997). Thus, we feel it is useful to discuss some of the new features of the DEPM survey that we developed for estimating $P_0$ with the CUFES.

### Critical value for allocation sampling

To use the CUFES in our DEPM survey design requires selecting an egg density that triggers full-water-column sampling (CalVET sampling). We used a critical value of 2 eggs/min, which was equivalent to 3 or 8 eggs/tow, depending on the conversion factor. If the 1997 conversion factor (eggs/min=0.25 egg/tow) was correct, we could lower the critical value to 1 egg/min, equivalent to 4 eggs/tow. This would create a larger high-density stratum and more CalVET tows would be allocated. This range of critical values (3–8 eggs/tow) was similar to the value (5 eggs/tow) used in a stratified sampling design for an anchovy survey in Biscay Bay in Spain (Petitgas, 1997). As a result, the precision of $P_0$ and $z$ would likely be improved. Increasing the area for the high-density sampling also reduces the potential for bias from the assumption of a constant egg mortality between strata. On the other hand, lowering the critical value diminishes the gain from using the CUFES. Clearly, an optimal critical value exists for each species and survey area. The critical value can be determined prior to the survey or during the survey by using order statistics (Thompson and Seber, 1996; Quinn et al., 1999).

The extent that the critical value can be fine tuned to deliver an optimum balance between the CUFES and CalVET samples for a particular region, species, and season is unknown. The large difference in catch ratios between our 1996 (0.73) and 1997 (0.25) surveys certainly does not support the idea of fine tuning. These differences may overstate the expected variability for sardine because the areas were different; the 1996 samples were taken over a very limited portion of the survey area , whereas in 1997, the sample pairs were taken where high-density spawning occurred (Figs. 1A, 2, and 6). Interestingly, our 1997 estimate (0.25) is similar to that computed by us for sardine off South Africa (van der Lingen et al., 1998).

### Variability in catch ratios

The extent of vertical mixing of eggs is probably the main factor affecting the variation of the catch ratio between surveys. Because the selection of the optimal critical value for CalVET sample allocation depends upon on the relationship between catches in the CUFES and those in the CalVET, or the catch ratio, it seems useful to consider what

the ratio would be if water were perfectly mixed in the whole-water-column. If the water were perfectly mixed, fish eggs would be distributed randomly and the catch ratio (between CalVET and CUFES) would be a constant, because egg count would be proportional to the volume of water filtered through the two samplers. A CUFES filters on the average 0.64 $m^3$ of water/min and the CalVET filters 3.5 $m^3$ of water. Therefore under perfect mixing, the ratio of eggs/min to eggs/tow = 0.64 $m^3$/3.5 $m^3$ = 0.18 for the daytime when fish schools are in deeper water, assuming the vertical distribution of sardine eggs was similar to that of anchovy eggs (Moser and Pommeranz, 1999). At night, when fish schools are in the upper 50 meters,[4] the ratio would be 0.64 $m^3$/2.5$m^3$ = 0.26. This means that on the average, for four to six eggs seen in a CalVET catch, we would expect one egg/min in the CUFES. The fact that the estimated ratio for two years were 0.76 in 1996 and 0.25 in 1997, indicates that more sardine eggs appeared in the upper 3 m than would be expected under perfect mixing and less mixing occurred in 1996 than in 1997. Clearly, if all the eggs were in the upper 3 m of water column, the ratio would be 1:1 for the same surface area. As the extent of vertical mixing in the sea is highly variable, we believe that calibration tows are always needed, even if the CUFES is used only as an index of egg abundance. Vertical egg mixing models might eventually help to reduce calibration requirements.

## Identification and counting of eggs at sea

Identification and counting of fish eggs while the ship is underway was an essential ingredient of our adaptive allocation design. The eggs of some commercially important clupeid species are often difficult to distinguish from those of co-occurring species (Ahlstrom and Moser, 1980; Matarese and Sandknop, 1984; Watson and Sandknop, 1996). Further, many melanostomiin species have eggs with characteristics (e.g. large diameter, wide perivitelline space, segmented yolk) that are similar to those of co-occurring sardines and other clupeid eggs. During a DEPM survey off Oregon, Bentley et al. (1996) encountered a type of melanostomiin egg similar to the egg of Pacific sardine at 24 of 46 stations. Our experience has shown that the risk of misidentification increases when identifications are made at sea; during our CUFES and DEPM survey in 1997, about 1% of the eggs initially identified as sardines turned out to be melanostomiin eggs after examination in the laboratory. Fortunately, melanostomiin eggs were in such low abundance that they had no effect on the critical values of density. The possibility of misidentification differs depending on season, location, and target species. Clearly all shipboard positive records for areas that lie outside of the known spawning range and season of the target species should be checked in the laboratory after the cruise. Owing to the great abundance of sardines and anchovy eggs, the effect of misidentification on biomass estimation is probably trivial if the survey is conducted during peak spawning months.

Egg counts on shipboard were somewhat lower than shoreside counts. Even though the effect of the difference is negligible from the standpoint of biomass estimation, we recommend that shoreside measurement be maintained. In particular, for collections that contain a large amount of other organisms (e.g. salps) which makes it difficult to count the eggs on aboard the ship, shoreside counting would be necessary.

## Stratified design

An important feature of the survey design was the stratification of sampling by egg density and sampler type. In the high-density stratum, we used only staged eggs from CalVET samples to estimate $z$ and $P_0$, whereas in the low-density stratum we collected only CUFES samples. We believe it would not be useful to use staged eggs from CUFES samples in the low-density stratum to estimate $z$ and $P_0$ directly because of the low egg abundance, possibility of stage-specific bias, and lack of yolksac larvae (the CUFES does not sample yolksac larvae well). It seemed preferable to use, for the low-density stratum, the estimate of $P_0$ for the high-density stratum, adjusted by the ratio of egg densities taken in CUFES at high and low strata. This, of course, requires the assumption that egg mortality did not differ between strata. A direct test of this assumption is impractical because of the large sampling effort needed at low density to obtain a sufficient number of positive samples to detect a difference in mortality rates. Fortunately, the effect of this potential bias is diminished because the low-density stratum contributes fewer eggs. In our example, the low-density stratum contributed about 25%[5] of the daily production. An alternative approach would be to allocate CalVET sampling to the low-density stratum. We believe this approach would not be cost effective because the number of positive CalVETs would be so low.

Another important element of the stratified design was the use of transect lines as the sampling unit. Model-based geostatistics are needed (Fletcher and Sumner, 1999) for data from continuous samplers such as echo-sounders and CUFESs, unless sampling units are defined such that data are uncorrelated among sampling units in the survey design (Armstrong et al., 1988). Because conventional design-based statistical procedures are easier to apply, we preferred using a transect line as our sampling unit, which requires that the minimum sample size allows a between-transect distance greater than the diameter of the egg patch. Fortunately the within-transect CUFES collections provided the information needed on the spatial structure to determine the distance of CalVET lines to insure samples are uncorrelated. In our case, tows, a minimum of 22.2 km

[4] Castillo Valderrama, P. R. 1995. Distribucion de los principales recursos pelagicos durante los veranlos de 1992 a 1994. Instituo Del Mar Del Peru, Informe No 114. Instituto del Mar delPeru (IMARPE), Esquina Gamarra Y General Valle Chucuito, Callao Peru, 24 p.

[5] 25% =100 × (1.064×107,255)/(2.57×174,096) (Table 4).

(12 nmi) apart, would ensure uncorrelated samples and provide the unbiased estimates of abundance of eggs in each development stage of sardine eggs.

### Use of yolksac larvae

Although yolksac larvae have been used to estimate $P_0$ with the DEPM method (Lo, 1985; 1986; Lo et al., 1996; Hunter and Lo, 1997), but are not a requirement for using the CUFES, we included yolksac larvae to estimate $P_0$ because the mortality rate of eggs and yolksac larvae are similar for northern anchovy (Lo, 1985, 1986) and because the development of early stages of anchovy and sardine is similar (Ahlstrom, 1943; Zweifel and Lasker, 1976; Moser and Alhstrom, 1985; Lo et al., 1996). Our threshold for taking CalVET samples, 2 eggs per minute, generated 102 positive CalVETs (72% were positive). This number is far fewer positive tows than the number needed to be assured a significant slope for the regression of numbers of eggs on their ages. In anchovy, where the eggs are less patchy than sardine, about 500 positive tows were required to assure a CV of the estimate of mortality rate close to 0.6 (Lo, 1997). To obtain a significant slope in the 1997 survey, we used the number and average age of yolksac larvae (adjusted for observed mean temperature) as well as staged eggs. This introduced a potential bias, because by doing this we assumed that yolksac larvae have the same mortality rate as eggs. To avoid this bias, the number of staged egg samples should be increased. One approach would be to stage the eggs taken with a CUFES in the high-density stratum and combine them with the CalVET samples from the same stratum. This would greatly increase the number of staged eggs available but requires the assumption that CUFES staged eggs are an unbiased sample of the full-water-column, a condition not met in our 1996 pilot study. Another approach would be to increase the number of CalVET samples per mile in the high-density stratum from the present one sample per 4 nmi to a higher frequency. A doubling of the CalVET sampling rate would significantly increase survey costs and may not increase the number of positive samples sufficiently to be able to use only eggs for estimation of $P_0$. Considering the relative risks and costs of these approaches, we feel that the use of the yolksac larvae in the estimation of $P_0$ was preferable. In other species, the distribution of eggs may be less concentrated than they are for sardine and sampling at the 4 nmi sampling rate may be adequate. Clearly, other solutions may exist, and we recommend considering these issues with each new application.

## Conclusions

We conclude that the CUFES is a useful tool with the DEPM when it is used adaptively to establish sampling strata for CalVETs. In this mode, the CUFES increases precision and reduces cost per transect mile. Perhaps one of the major benefits of using a CUFES in the DEPM is that one can better afford to expand survey boundaries and thereby reduce the potential bias of not enclosing the entire population, a very common bias in field surveys in general (Gunderson, 1993). Clearly, the more contiguous the distribution of spawned eggs within the surveyed habitat, the greater the benefits in using the CUFES.

## Acknowledgments

We thank three unanimous reviewers for their comments; David Griffith, Ron Dotson, and Amy Hays for operating the CUFES for 1996 and 1997; David Ambrose, William Watson, and Elaine Acuna for counting and staging eggs of CUFES and CalVET samples; Geoffrey Moser for providing the information on misidentification of sardine eggs; and crew members of NOAA RV *David Starr Jordan* for their cooperation.

## Literature cited

Alheit, J.
1993. Use of the daily egg production method for estimating biomass of clupeoid fishes, a review and evaluation. Bull. Mar. Sci. 53(2)750–767.

Ahlstrom, E. H.
1943. Studies on the Pacific pilchard or sardine (*Sardinpos caerulea*). 4. Influence of temperature on the rate of development of pilchard eggs in nature. U.S. Fish. Wildl. Serv., Spec. Sci. Rep. 23, 26 p.

Ahlstrom, E. H., and H. G. Moser.
1980. Characters useful in identification of pelagic marine fish eggs. Calif. Coop. Oceanic Fish. Investig. Rep. 21:121–131.

Armstrong, M., P. Shelton, I. Hampton, G. Jolly, and Y. Melo.
1988. Egg production estimates of anchovy biomass in the southern Benguela. Calif. Coop. Oceanic. Fish. Investig. Rep. 29:137–157.

Barange, M., and I. Hampton.
1997. Spatial structure of co-occuring anchovy and sardine populations from acoustic data: implications for survey design. Fish. Oceangr. 6:294–108.

Bentley. P. J., R. L. Emmett, N. C. H. Lo, and H. G. Moser.
1996. Egg production of Pacific sardine (*Sardinops sagax*) off Oregon in 1994. Calif. Coop. Oceanic. Fish. Investig. Rep. 37:193–200.

Chambers, J. M., and T. J. Hastie (ed).
1992. Statistical models in S. Wadsworth and Brooks/Cole Advanced Books and Software, Pacific Grove, CA, 608 p.

Checkley, D. M., Jr, R. C. Dotson, and D. A. Griffith.
1999. Continuous, Underway sampling of eggs of northern anchovy (*Engraulis mordax*) and Pacific sardine (*Sardinops sagax*) in spring 1996 and 1997 off southern and central California. Deep-Sea Res. (special rep.) II 47(2000) 1139–1155.

Checkley, D. M., Jr., P. B. Ortner, L. R. Settle, L.R., and S. R. Cummings.
1997. A continuous, underway fish egg sampler. Fish. Oceanogr. 6:58–73.

Cressie, N.
1991. Statistics for spatial data. John Wiley and Sons, Inc. New York, Chichester, Toronto, Brisbane, Singapore, 900 p.

Cressie, N., and D. M. Howkins.
1980. Robust estimation of the variogram. I. J. Int. Assoc. Math. Geology 12:115–125.

Efron, B., and R. J. Tibshirani.
1993. An introduction to the bootstrap. Monographs on statistics and applied probability 57. Chapman and Hall, New York, London, 436 p.

Fletcher, W. J., and N. R. Sumner.
1999. Spatial distribution of sardine (*Sardinops sagax*) eggs and larvae: an application of geostatistics and resampling to survey data. Can. J. Aquat. Sci. 56:907–914.

Gunderson. D. R.
1993. Surveys of fisheries resources. John Wiley and Sons, Inc., New York, Chichester, Brisbane, Toronto, Singapore, 248 p.

Goodman, L. A.
1960. On the exact variance of products. J. Am. Stat. Assoc. 55(N292):708–713.

Hunter, J. R., and N. C. H. Lo.
1997. The daily egg production methods of biomass estimation: some problems and potential improvements. OZEANOGRAFIKA, Boletin De La Sociedad De oceanografica De Gipuzkoa, Medalla De Oro De La Ciudad De San Sebastian 2:9–40.

Kaluzny, S. P., C. Vega, T. P. Cardoso, and A. A. Shelly.
1996. S+ spatialstats. User's manual, version 1. Mathsoft, Inc., Seattle, WA, 226 p.

Lasker, R (ed).
1985. An egg production method for estimating spawning biomass of pelagic fish: application to the northern anchovy (*Engraulis mordax*). U.S. Dep. Commer., NOAA Tech. Rep. NMFS 36, 99 p.

Lo, N. C. H.
1985. Egg production of the central stock off northern anchovy, *Engraulis mordax*, 1951–82. Fish. Bull. 83(2)137–150.
1986. Modeling life-stage-specific instantaneous mortality rates, an application to northern anchovy, *Engraulis mordax*, eggs and larvae. Fish. Bull. 84(2):395–406.
1997. Empirical analyses of the precision of regression estimates of daily egg production and egg mortality of pelagic fish for daily egg production method. OZEANOGRAFIKA, Boletin De La Sociedad De Oceanografica De Gipuzkoa,Medalla De Oro De La Ciudad De San Sebastian 2:71–89.

Lo, N. C. H., Y. A. Geen Ruiz, M. Jaacob Cervantes, H. G. Moser, and R. J. Lynn.
1996. Egg production and spawning biomass of Pacific sardine (*Sardinops sagax*) in 1994, determined by the daily egg production method. Calif. Coop. Oceanic. Fish. Investig. Rep. 37:160–174.

Macewicz, B. J., J. J. Castro Gonzalez, C. E. Cotero Altamirano, and J. R. Hunter,
1996. Adult reproductive parameters of Pacific sardine, *sardinops sagax*, during 1994. Calif. Coop. Oceanic. Fish. Investig. Rep. 37:140–151.

Matarese, A. C., and E. M. Sandknop.
1984. Identification of fish eggs. *In* Ontogeny and systematics of fishes (H. G. Moser, W. J. Richards, D. M. Cohen, M. P. Fahay, A. W. Kendall Jr., and S. L. Richardson, eds.), p. 27–31. Am. Soc. Ichthyol. Herpetol. Spec. Publ. 1.

Moser, H. G., and E. H. Ahlstrom.
1985. Staging anchovy eggs. *In* An egg production method for estimating spawning biomass of pelagic fish: application to the northern anchovy, *Engraulis mordax* (R. Lasker, ed.), p. 37–42. U.S. Dep. Commer., NOAA Technical Report NMFS 36.

Moser, H. G., and T. Pommeranz.
1999. Vertical distribution of eggs and larvae of northern anchovy, *Engraulis mordax*, and of the larvae of associated fishes at two sites in the Southern California Bight. Fish. Bull. 97:920–943.

Parker, K.
1985. Biomass model for the egg production method. *In* An egg production method for estimating spawning biomass of pelagic fish: application to the northern anchovy, *Engraulis mordax* (R. Lasker, ed.), p. 5–6. U.S. Dep. Commer., NOAA Technical Rep. NMFS 36.

Petitgas, P.
1993. Geostatistics for fish stock assessments: a review and an acoustic application. ICES J.Mar. Sci. 50:285–298.
1997. Use of disjunctive Kriging to analyse an adaptive survey design for anchovy, *Engraulis encraicolus*, eggs in Biscay Bay. OZEANOGRAFIKA, Boletin De La Sociedad De Oceanografica De Gipuzkoa,Medalla De Oro De La Ciudad De San Sebastian 2:121–132.

Priede, I. G., and J. J. Watson.
1993. An evaluation of the daily egg production method for estimating biomass of Atlantic mackerel (*Scomber scombrus*). Bull. Mar. Sci. 53(2)891–911.

Quinn, T .J., II, D. Hanselman, D. Clausen, C. Lunsford, J. Heifetz.
1999. Adaptive cluster sampling of rockfish populations: proceedings of the American Statistical Association, 1999 Joint Statistical Meeting, Biometrics section. Am. Stat. Assoc., Alexandria, VA.

Smith, P. E.
1973. The mortality and dispersal of sardine eggs and larvae. Rapp. P.-V. Reun. Cons. Perm. Int. Explor. Mer 164:282–292.

Smith, P. E., and R. P. Hewitt.
1985. Anchovy egg dispersal and mortality as interred from close interval observations. Calif. Coop. Oceanic. Fish. Investig. Rep. 26:97–112.

Smith, P. E., W. Flerx, and R. P. Hewitt.
1985. The CalCOFI vertical egg tow (CalVET). *In* An egg production method for estimating spawning spawning biomass of pelagic fish: application to the northern anchovy (*Engraulis mordax*) (R. Lasker, ed.), p. 27–32. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 36.

Sundby, B.
1983. A one-dimensional model for the vertical distribution of pelagic fish eggs in the mixed layer. Deep-Sea Res., 30: 645–661.

Thompson, S. K., F. L. Ramsey, and G. A. F. Seber.
1992. An adaptive procedure for sampling animal populations. Biometrics 48:1195–1199.

Thompson, S. K., and G. A. F. Seber.
1996. Adaptive sampling. John Wiley and Sons, Inc., New York, Chichester, Brisbane, Toronto, Singapore, 265 p.

van der Lingen, C. D., D. Checkley Jr, M. Barange, L. Hutchings, and K. Osgood.
1998. Assessing the abundance and distribution of eggs of sardine, *Sardinops sagas*, and round herring, *Etrumenus whiteheadi*, on the western Agulhas Bank, South Africa, using a continuous, underway fish egg sampler. Fish. Oceanogr. 7(1):35–47.

Watson, W., and E. M. Sandknop.
1996. Clupeidae. *In* The early stages of fishes in the California Current region (H. G. Moser, ed.), 159–171 p. Calif. Coop. Oceanic Fish. Invest. Atlas 33.

Watson, W., R. L. Charter, H. G. Moser, R. D. Vetter, D. A. Ambrose, S. R. Charter, L. I. Robertson, E. M. Sandknop, E. A. Lynn, and J. Stannard.
1999. Fine-scale distributions of planktonic fish eggs in the

vicinities of Big Sycamore Canyon and Vandenberg Ecological Reserves, and Anacapa and San Miguel Island, California. Calif. Coop. Oceanic. Fish. Investig. Rep. 40:128–153.

Westgard, T.

1989. Two models of the vertical distribution of pelagic fish egg in the turbulent layer of the ocean. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 191:195–200.

Zeldis, J. R.

1993. Applicability to egg surveys for spawning-stock biomass estimation of snapper, orange roughy, and hoki in New Zealand. Bull. Mar. Sci.:53(2) 864–890.

Zweifel, J. R., and R. Lasker.

1976. Prehatch and posthatch growth of fishes—a general model. Fish. Bull. 74(3):609–621.

# Growth of juvenile red king crab (*Paralithodes camtschaticus*) in Bristol Bay (Alaska) elucidated from field sampling and analysis of trawl-survey data

**Timothy Loher**
**David A. Armstrong**
School of Fisheries and Aquatic Sciences
Box 355020
University of Washington
Seattle, Washington 98195
Present address (for T. Loher): International Pacific Halibut Commission
P.O. Box 95009
Seattle, Washington 98145-2009
E-mail address (for T. Loher): loher@u.washington.edu

**Bradley G. Stevens**
AFSC Kodiak Laboratory
Kodiak Fisheries Research Center
National Marine Fisheries Service
301 Research Court
Kodiak, Alaska 99615

**Abstract**–An analysis of *in situ* growth rate was conducted for juvenile red king crab (*Paralithodes camtschaticus*) in Bristol Bay, Alaska. Growth of early juveniles (~2–40 mm CL; age 0–3 yr) was determined by fitting seasonalized Gompertz growth models to length-frequency data. The parameters of the growth model and resulting size-at-age estimates were compared with those from studies conducted at Unalaska and Kodiak Islands by fitting the same growth model to published length-frequency data from separate sources. Growth of late juvenile and early reproductive crabs, ~30–100 mm carapace length (CL), was examined by analyzing length-frequency data from the National Marine Fisheries Service annual Bering Sea trawl survey from 1975 through 1999. Mean CL associated with strong size modes of crabs in Bristol Bay length-frequency distributions was resolved by using the FiSAT software package (FAO-ICLARM Stock Assessment Tools) to track the modal size progression of strong year classes and assign mean size-at-age characteristics to the stock.

Growth of early juvenile crabs was slower in Bristol Bay than that observed by other researchers at Unalaska or Kodiak. Sizes at 1, 2, and 3 years after settlement were estimated to be ~9 mm, 23 mm, and 47 mm CL in Bristol Bay compared with 16 mm, 38 mm, and 66 mm CL at Unalaska; at Kodiak, estimated sizes of 12 mm and 42 mm were obtained for age-1 and age-2 crabs, respectively. Within the Bristol Bay trawl survey data, a total of 24 modes were identified for both males and females <~100 mm CL, which included the modal progression of two year classes that presumably settled in 1976 and 1990. The 1976 year class grew slowly and would not have recruited to the reproductive stock until ~9 years after settlement, whereas the 1990 year class appeared to recruit at ~8 years after settlement. Both estimates indicate that Bristol Bay red king crabs are older at reproductive maturity than the ~6 years after settlement presently assumed. An attempt to resolve discrete mean size-at-age from the length-frequency data met with little success because variability in growth between year classes markedly obscured size-at-age characteristics in the stock.

Manuscript accepted 20 March 2001.
Fish. Bull. 99:572–587 (2001).

Population abundance of red king crab (*Paralithodes camtschaticus*) in Bristol Bay, Alaska (Fig. 1) is typified by great variability. The maximum abundance of harvestable male crabs in the stock over the last 25 years has fluctuated by over an order of magnitude, peaking at nearly 60 million individuals in 1977, and has fallen to less than 3 million from 1983 to 1985 (Loher et al., 1998; Zheng and Kruse[1]). Strong fisheries in the late 1970s and early 1980s were followed by substantially reduced abundance in recent years, and fishery closures in 1981 and 1994, leading to concern over the status of the population by both management agencies and fishermen. Such concern prompted the Alaska Department of Fish and Game to develop a detailed harvest-based recovery plan for the fishery (Zheng et al.[2]) that considers the stock to be fully recovered once it reaches an effective spawning biomass of 55 million pounds. This biomass has not occurred in Bristol Bay since 1981 (Zheng and Kruse[1]; Zheng et al.[2]), but it was chosen as an appropriate rebuilding level primarily on the basis of length-based recruitment models that, in turn, rely on the inferred underlying stock-recruitment relationship for the population (Zheng et al., 1995a, 1995b).

Present assumptions regarding red king crab growth rates suggest that Bristol Bay crabs recruit to the reproductive stock at an age of approximately 6 years after settlement (i.e. ~seven years following egg fertilization; Zheng et al., 1995a, 1995b), which corresponds to mean sizes of 105 mm carapace length (CL) and 97 mm CL for males and females, respectively. These size-at-age values are based on Stevens and Munk (1990), Weber (1967), and Balsiger (1974). However, none of these sources represents a comprehensive treatment of growth from settlement through maturity based entirely on field-collected data from the Bristol Bay region. Balsiger's (1974) growth

[1] Zheng, J., and G. H. Kruse. 1999. Status of king crab stocks in the eastern Bering Sea in 1999. Alaska Department of Fish and Game, Reg. Inf. Rep. 5J99-09. Division of Commercial Fisheries, Alaska Department of Fish and Game, P.O. Box 25526, Juneau, Alaska, 99801.

[2] Zheng, J., M. C. Murphy, and G. H. Kruse. 996. Overview of population estimation methods and recommended harvest strategy for red king crabs in Bristol Bay. Alaska Department of Fish and Game, Reg. Inf. Rep. 5J96-04. Division of Commercial Fisheries, Alaska Department of Fish and Game, P.O. Box 25526, Juneau, Alaska, 99801.

model was derived from data from Bristol Bay, but it considered only individuals >81 mm carapace length, well above the size at which multiple yearly molts are expected. Weber's (1967) work was based on juvenile crabs collected more than 40 years ago at Unalaska Island (Fig. 1) in the eastern Aleutian Islands, which is located west of Unimak Pass and the coastal shelf break and which is oceanographically separated from Bristol Bay by the southern origins of the Bering Slope current. Stevens and Munk's (1990) growth model was based on data from the Kodiak region (Fig. 1); a separate growth model was developed by Stevens (1990) to consider the eastern Bering Sea, but it relies on Weber's (1967) findings regarding the growth of early juvenile crabs.

Accurate growth rate information is important to properly calibrate the length-based recruitment model and also to determine appropriate time lags between spawning and subsequent recruitment. Given the likelihood of temporal and geographic variability in growth rates, it is likely that the Bristol Bay stock exhibits growth rates different from those observed at different locations by the aforementioned researchers. Environmentally induced changes in molt schedule resulting in considerable variability in size-at-age is a common feature in Crustacea (e.g. Hartnoll, 1982; Hill et al., 1989; Huner and Romaire, 1990; Wainright and Armstrong, 1993; Tremblay and Eagles, 1997) and probably also in red king crab (Stevens, 1990; Stevens and Munk, 1990). Thus, further analysis of the growth of prerecruit crabs within Bristol Bay is needed. In our study, we analyzed growth of early juvenile red king crabs in Bristol Bay by fitting growth equations to length-frequency data collected between 1983 and 1991. We compared the inferred growth rate for Bristol Bay with that determined from identical models fitted to Weber's (1967) data, and with two sets of length-frequency data available for the Kodiak region (Dew, 1990; Donaldson et al., 1992) to assess whether application of these growth rates to Bristol Bay crabs is appropriate. We then expanded our analyses to include older prerecruit crabs in Bristol Bay by analyzing length-frequency distributions from 25 years of southeast Bering Sea trawl survey data in order to identify growth patterns associated with strong year classes and in order to elucidate size-at-age characteristics displayed by the population.

## Materials and methods

### Growth of early juvenile red king crab

Growth of early juvenile crabs, from settlement through approximately 3 years after settlement, was examined by using catch data from targeted sampling in Bristol Bay obtained from three sources: 1) work conducted in 1983 under the auspices of the Outer Continental Shelf Assessment Program (OCSEAP) (documented in McMurray et al.[3]), 2) OCSEAP work conducted in 1985 (previously unpublished), and, 3) work conducted in 1991 by the National Marine Fisheries Service (NMFS; documented in Stevens and MacIntosh[4]). In addition we reviewed the




**Figure 1**

Map of western Alaska, showing the locations of Bristol Bay, Unalaska Island, and Kodiak Island.

results of Weber (1967), who collected early juvenile red king crabs at Unalaska Island in 1958 and 1959, as well as data collected in the Kodiak region from 1987 to 1989, originally published in Dew (1990), and from 1990 to 1991, originally published in Donaldson et al. (1992). Note that, in the present paper, all ages and year-class designations are referenced to approximate settlement date. That is, they refer to the postsettlement age of benthic crabs and do not include the larval phase or the egg incubation period. This postsettlement age should be taken into consideration for applications in which the year when eggs were extruded is important.

**Bristol Bay** Waters of the Bering Sea east of 163.5°W longitude and south of 59°N latitude will be considered "Bristol Bay." This area is larger than the area that is often referred to as Bristol Bay; it is more typically considered the "southeast Bering Sea." However, because we also report information from Unalaska Island, also in the southeastern Bering Sea, we choose to make a distinction between Bristol Bay and Unalaska Island in order to avoid confusion. In 1983, surveys of juvenile red king crab abundance were conducted throughout the Bristol Bay region (Fig. 2A) with surface-deployed try-net otter trawls and a rock dredge, during three sampling periods: 18 April–7 May, 2–17 June, and 9–23 September (McMurray et al.[3]).

[3] McMurray, G., A. H. Vogel, P. A. Fishman, D. A. Armstrong, and S. C. Jewett. 1984. Distribution of larval and juvenile red king crab (*Paralithodes camtschatica*) in Bristol Bay. U.S. Dep. Commer., NOAA, OCSEAP Final Report 53(1986):267–477, Anchorage, Alaska. [Available from D.A. Armstrong at: School of Fisheries and Aquatic Sciences, Univ. Washington, Box 355020, Seattle, WA 98195.]

[4] Stevens, B. G., and R. A. MacIntosh. 1991. Cruise 91-1 Ocean Hope 3: 1991 eastern Bering Sea juvenile red king crab survey, May 24–June 3, 1991. U.S. Dep. Commer., NOAA, NMFS, AFSC, RACE. Seattle, Washington. [Available from B. G. Stevens at AFSC Kodiak Laboratory, National Marine Fisheries Service, 301 Research Court, Kodiak, Alaska, 99615.]




**Figure 2**

Location of survey stations in Bristol Bay (east of 163.5° west longitude) visited during 1983 OCSEAP studies (**A**), 1985 OCSEAP studies (**B**, open squares), and by Stevens and MacIntosh in 1991 (B, closed circles).

Try nets had a headrope length of 5.4 m and either wooden or aluminum doors measuring 0.4 m wide × 0.9 m tall. The rock dredge was constructed of a rigid steel frame with a mouth opening 0.9 m wide × 0.4 m tall. The 1985 survey was conducted from 19 July to 2 August, nearshore (generally inside the 50-m isobath), along the north Aleutian Shelf from Unimak Pass through the Port Moller region (Fig. 2B) also with try net and rock dredge. The 1991 survey (Stevens and MacIntosh[4]) was conducted from 26 May to 1 June along the North Aleutian Shelf from Port Moller to Kvichak Bay (Fig. 2B) with a 3.1-m wide beam trawl. The carapace length (CL) of all red king crab captured in the above surveys was measured and recorded onboard the vessels.

For each sampling period, length-frequency histograms were constructed to identify individual age classes (cohorts) within the data. Male and female crabs were pooled because sex-specific growth rates are not apparent until reproductive age (Weber, 1967; Dew, 1990), and the combined length-frequency data were analyzed by using the FiSAT software package (Gayanilo and Pauly, 1997) to determine the mean size (±1 SD) for each identifiable cohort. FiSAT employs a combination of Bhattacharya's method (Bhattacharya, 1967) and NORMSEP (Hasselblad, 1966; Pauly and Caddy[5]) to decompose complex size-frequency distributions into a series of best-fit normal curves that represent each cohort within the data set. In our study, the "mean size" of a cohort refers to the mean (±1 SD) of its associated best-fit normal curve, as determined by FiSAT size-frequency decomposition.

Postsettlement age, in Julian days, was then calculated for each cohort. For each year's length-frequency histogram, the cohort with the smallest mean CL was assigned age 0, and the subsequent sizes assigned age 1 and age 2 (Fig. 3). The age of each cohort (in days) was calculated as the time from settlement in the cohort's settlement year until the median sampling date for the survey period. Settlement was estimated to be 15 July of each year because numerous 2-mm-CL individuals occurred in late July during the 1983 surveys, and a carapace length of 2 mm is typical of the first benthic instar (Kurata, 1961; Donaldson et al., 1992; Loher and Armstrong, 2000). Postsettlement age was plotted against the associated mean CL and two growth curves were fitted to the data. The first curve was a seasonalized version of the von Bertalanffy growth model obtained from Anastacio and Marques (1995), where growth in carapace length is expressed as

$$L_t = L_{max}\left\{1 - e^{-[K\times D\times(t-t_{min})+C\times(K\times D/2\Pi)\times\sin 2\Pi t\times(t-t_s)]}\right\}^{(1/D)}, \quad (1)$$

The second curve was a seasonalized Gompertz model, where growth in carapace length is expressed as

$$L_t = L_{max}\left\{e^{-e^{-[K\times(t-t_{min})+C\times(K/2\Pi)\times\sin 2\Pi t\times(t-t_s)]}}\right\}, \quad (2)$$

where (in both models) $L_t$ = carapace length at time $t$;
$L_{max}$ = maximum carapace length;
$t$ = given time;
$t_{min}$ = time at which the carapace length is the minimum size for the life-stage of interest (in this case, minimum size for benthic red king crab=2 mm);

[5] Pauly, D., and J. F. Caddy. 1985. A modification of Bhattacharya's method for the analysis of mixtures of normal distributions. FAO Fisheries Circular 781, Sales and Marketing Group, Information Division, FAO, Viale delle Terme di Caracalla, 00100 Rome.




**Figure 3**

Length-frequency histograms of early juvenile red king crabs captured in Bristol Bay from 1983 to 1991. Each age class is denoted by shading: open bars = age 0+; shaded bars = age 1+; closed bars = age 2+. For each sampling date an average time after settlement (ATAS) is reported. The ATAS is the number of days estimated to have elapsed between 15 July of the sampling year (the most recent settlement event) and the median sampling date. For example, the ATAS for crabs sampled on 15 September was 62 days. This value corresponds to the estimated age of age 0+ crabs in that sample, whereas age 1+ crab would have been ~365 days older than the ATAS, and age 2+ crabs ~730 days older than the ATAS. Note that the June samples fell at nearly one calendar year following the previous year's settlement; hence, age 0+ crab in those samples were nearly age 1. Arrows indicate the mean CL of each size cohort, as determined with FiSAT; mean values were used to fit the growth curve.

$t_s$ = lag-time between the start of growth and the first seasonal growth oscillation; oscillations are sinusoidal with a one-year period;
$K$ = intrinsic growth rate;
$C$ = parameter ranging from 0 to 1 that controls the strength of the seasonal growth oscillation; 0 = no seasonal signature; 1 = strong seasonality with a brief period each year during which growth ceases; and
$D$ = parameter expressing metabolic deviation from the von Bertalanffy 2/3 metabolic rule (in our study, $D$ = 1 [no deviation]).

**Unalaska and Kodiak islands** Growth rate of early juvenile red king crab at Unalaska Island was assessed by using data published in Weber (1967). In that study immature crabs were collected, primarily with SCUBA, during four sampling periods in 1958 (22 April–17 May; 30 May–6 June; 13 July; 17 September–1 October) and two periods in 1959 (11–24 February; 24 May–2 June). Two data sources were used that originated in Kodiak Island: 1) Donaldson et al. (1992), who documented growth of red king crab in artificial habitat collectors for approximately one year after settlement between June 1990 and May 1991, and; 2) Dew (1990) who collected data on podding age 1+ to 2+ red king crab using SCUBA observations between 20 November 1987 and 3 June 1989. The data in Weber (1967) and Dew (1990) were used to construct length-frequency histograms (males and females pooled) and the characteris-

**Table 1**

Points used to construct the growth curve for early juvenile red king crab in Bristol Bay. Data originally presented in Stevens and MacIntosh[4] are indicated as such; previously unpublished data are listed as "new."

| Collection date | Source | Estimated age (days after settlement) | *n* | Size range (mm) | Mean size ±1 SD |
|---|---|---|---|---|---|
| 19 Jul–2 Aug 1985 | new | 11.5 | 6 | 2 | 2.00 ±0.000 |
| 10 Sep–19 Sep 1983 | new | 62 | 134 | 3.0–6.0 | 3.97 ±0.581 |
| 21 Apr–6 Jun 1983 | new | 283.5 | 10 | 4.0–6.0 | 4.27 ±0.730 |
| 26 May– Jun 1991 | Stevens and MacIntosh | 319.5 | 35 | 4–14 | 7.70 ±1.645 |
| 5 Jun–16 Jun 1983 | new | 332 | 13 | 3.0–7.2 | 5.19 ±0.982 |
| 18 Jul–2 Aug 1985 | new | 376.5 | 3 | 8.0–8.4 | 8.13 ±0.231 |
| 10 Sep–19 Sep 1983 | new | 427 | 45 | 10–28 | 14.63 ±3.447 |
| 21 Apr–6 May 1983 | new | 648.5 | 30 | 10–32 | 16.70 ±5.389 |
| 26 May–1 Jun 1991 | Stevens and MacIntosh | 684.5 | 16 | 18–29 | 22.53 ±3.562 |
| 5 Jun–16 Jun 1983 | new | 697 | 25 | 12–26 | 17.05 ±4.102 |
| 26 May–1 Jun 1991 | Stevens and MacIntosh | 1049.5 | 7 | 34–47 | 40.36 ±3.681 |

tics of individual cohorts were resolved by using FiSAT. Donaldson et al. (1992) did not provide their original data; therefore we simply used the mean size-at-age values that they reported. Mean size-at-age values for each region were then fitted with seasonalized von Bertalanffy and Gompertz curves, as described previously. Age of crabs at Unalaska Island was calculated by using 1 July as the approximate settlement date, as suggested by Weber (1967); 14 June was used for Kodiak Island because Donaldson et al. (1992) first collected benthic instars on this date.

### Mean size-at-age of late juvenile through early reproductive-age crabs

Growth of late juvenile (~age 2+) through early reproductive-age crabs was examined by using data from the annual NMFS groundfish trawl survey from 1975 to 1999 (please refer to Otto [1986] for a description of the annual trawl survey protocol and its spatial coverage). These data were used to construct annual length-frequency histograms for the Bristol Bay stock; length-frequency distributions were examined to identify strong size modes, and the mean CL (±1 SD) associated with each mode was determined by using FiSAT. Two approaches were then employed to assign discrete mean size-at-age categories to the population, focusing on size modes with mean CL <100 mm. First, each strong year class that recruited to the population was identified, and the growth of individuals in these year classes was determined by examination of progression of their length-frequency modes, from the first appearance of the year classes until they could no longer be resolved. Second, the mean CL associated with every strong size mode that appeared in all the annual length-frequency distributions, over the entire time series, was plotted to identify commonly occurring mean CLs that might represent consecutive size-at-age categories.

### Bottom temperature

Area-averaged near-bottom temperature was calculated for each year from 1975 to 1999, to more fully interpret the growth rates observed in Bristol Bay region with respect to variation over time. The area-averaged near-bottom temperature represented the mean temperature over all trawl-survey stations located within the Bristol Bay region. Because the survey data were sometimes incomplete and sampling stations were not located in precisely the same location each year, the temperature at the center of each trawl survey station was statistically interpolated with known values at surrounding points (i.e. "kriged"; see Cressie, 1993). Kriging was performed with the Surfer 6.04 software package (Keckler, 1994) and a linear variogram model (Cressie, 1993). Area-averaged near-bottom temperatures for the Bristol Bay region were then calculated by using the kriged estimates obtained from the center of trawl survey stations.

## Results

### Early juvenile red king crab

Resolution of age classes from length-frequency data collected in Bristol Bay in 1983, 1985, and 1991 provided 11 estimates of mean size-at-age representing individuals ranging from approximately 12 to 1050 days after settlement (refer to Fig. 3, Table 1). Weber (1967) provided information resulting in 24 mean size-at-age estimates (Table 2) at Unalaska, for crabs estimated to be ~86–964 days after settlement. Donaldson et al. (1992) and Dew (1990) provided 22 estimates of mean size-at-age for crabs ~0–720 days after settlement (Table 3) at Kodiak Island. Plots of early juvenile mean size-at-age are presented in Figure 4 for Bristol Bay, Unalaska Island, and

**Table 2**

Points used to construct the Unalaska early juvenile red king crab growth curve from data of Weber (1967) (his Appendix Table II). The original data come from crabs that were collected at a number of study sites around Unalaska Island; the "sample area" column in this table refers to different study sites as designated by Weber (1967). Refer to the original paper for descriptions of each specific sample area.

| Collection date | Sample area | Estimated age (days after settlement) | Molt status | Size range (mm) | Mean size (mm) ±1 SD |
|---|---|---|---|---|---|
| 17 Sep–1 Oct 1958 | 2 | 86 | not reported | 4–6 | 4.81 ±0.634 |
| 11–24 Feb 1959 | 2 | 232.5 | postmolt | 7–10 | 8.32 ±0.750 |
| 22 Apr–17 May 1958 | 2 | 309 | not reported | 8–17 | 12.62 ±1.799 |
| 24 May–2 Jun 1959 | 2 | 333.5 | pre- and postmolt | 8–16 | 11.26 ±1.406 |
| 30 May–6 Jun 1958 | 2 | 338.5 | not reported | 9–18 | 12.73 ±1.433 |
| 13 Jul 1958 | 2 | 379 | not reported | 13–18 | 16.06 ±0.869 |
| 17 Sep–1 Oct 1958 | 2 | 452 | not reported | 15–34 | 24.08 ±3.994 |
| 17 Sep–1 Oct 1958 | 4 | 452 | not reported | 17–29 | 20.22 ±1.500 |
| 11 –24 Feb 1959 | 2 | 598.5 | premolt | 23–35 | 29.62 ±2.376 |
| 11–24 Feb 1959 | 2 | 598.5 | postmolt | 25–44 | 35.12 ±3.798 |
| 11–24 Feb 1959 | 4 | 598.5 | pre- and postmolt | 22–40 | 30.56 ±4.200 |
| 22 Apr–17 May 1958 | 1 | 674 | not reported | 29–35 | 31.85 ±1.682 |
| 22 Apr–17 May 1958 | 2 | 674 | not reported | 23–48 | 36.68 ±4.392 |
| 24 May–2 Jun 1959 | 4 | 698.5 | premolt | 28–41 | 29.10 ±2.783 |
| 24 May–2 Jun 1959 | 4 | 698.5 | postmolt | 31–41 | 33.73 ±2.919 |
| 24 May–2 Jun 1959 | 2 | 698.5 | premolt | 25–41 | 35.00 ±2.388 |
| 24 May–2 Jun 1959 | 2 | 698.5 | postmolt | 28–51 | 39.65 ±4.238 |
| 30 May–6 Jun 1958 | 2 | 703.5 | not reported | 29–50 | 39.75 ±4.373 |
| 17 Sep–1 Oct 1958 | 4 | 817 | not reported | 41–53 | 45.86 ±3.308 |
| 11–24 Feb 1959 | 1 | 963.5 | premolt | 33–51 | 40.85 ±3.162 |
| 11–24 Feb 1959 | 1 | 963.5 | postmolt | 41–69 | 49.17 ±4.925 |
| 22–24 Feb 1959 | 3 | 963.5 | premolt | 42–74 | 52.70 ±6.000 |
| 11–24 Feb 1959 | 3 | 963.5 | postmolt | 51–82 | 66.23 ±8.030 |
| 11–24 Feb 1959 | 4 | 963.5 | premolt | 43–78 | 57.19 ±7.659 |

Kodiak Island data. Seasonalized Gompertz growth functions were fitted to the data; von Bertalanffy growth curves are not presented because their fits were poorer by comparison (Table 4). Model fits and parameter estimates for Gompertz growth functions are reported in Table 5, along with model estimates of CL at ages 1.0 and 2.0 at all sites, and at age 3.0 for Bristol Bay and Unalaska Island; predicted sizes at ages 0.9, 1.9, and 2.9 are also reported because the Bering Sea trawl survey occurs ~0.1 years prior to the settlement season. No data were available for Kodiak Island crabs beyond age ~2; therefore model predictions are not reported beyond this age.

Intrinsic growth rate was lowest in Bristol Bay and was coupled with a very strong seasonal signature, with no growth expected during mid-winter. The growth rate observed at Kodiak Island was higher than in Bristol Bay, with a moderate seasonal signature, whereas




**Figure 4**

Seasonalized Gompertz growth curves fitted to mean size-at-age data from Bristol Bay, Unalaska Island, and Kodiak Island. Note that no data were available for Kodiak crabs > ~2 years after settlement.

**Table 3**

Points used to construct the Kodiak early juvenile red king crab growth curve. Data from Donaldson et al. (1992) come directly from their Table 1; data from Dew (1990) are compiled from his Figures 7 and 8.

| Collection date | Estimated age (days after settlement) | Size range (mm) | Mean size (mm) ±SD |
|---|---|---|---|
| Donaldson et al. (1992) | | | |
| 14 Jun 1990 | 0 | 1.9–2.4 | 2.18 ±0.118 |
| 28 Jun 1990 | 14 | 1.8–2.6 | 2.18 ±0.171 |
| 13 Jul 1990 | 29 | 2.0–3.3 | 2.61 ±0.363 |
| 26 Jul 1990 | 42 | 2.5–3.4 | 2.84 ±0.152 |
| 10 Aug 1990 | 57 | 2.4–4.2 | 3.64 ±0.408 |
| 23 Aug 1990 | 70 | 2.6–5.4 | 3.78 ±0.362 |
| 7 Sep 1990 | 85 | 3.4–5.1 | 4.47 ±0.421 |
| 21 Sep 1990 | 109 | 4.2–6.4 | 4.98 ±0.605 |
| 16 Oct 1990 | 124 | 3.4–6.1 | 5.29 ±0.491 |
| 4 Dec 1990 | 173 | 4.6–7.9 | 6.58 ±0.703 |
| 11 Feb 1991 | 212 | 4.6–9.0 | 7.71 ±1.201 |
| 27 Mar 1991 | 257 | 6.1–10.3 | 7.95 ±0.769 |
| 14 May 1991 | 335 | 8.7–13.0 | 10.46 ±1.079 |
| 29 May 1991 | 350 | 8.6–9.9 | 9.33 ±0.474 |
| Dew (1990) | | | |
| 6 Oct 1988 | 480 | 17.5–34.0 | 25.67 ±2.446 |
| 18 Nov 1988 | 523 | 19.0–35.5 | 27.87 ±2.795 |
| 20 Nov 1987 | 525 | 22.0–32.5 | 26.93 ±2.572 |
| 18 Dec 1987 | 553 | 22.0–27.4 | 28.10 ±2.726 |
| 24 Feb 1989 | 621 | 25.0–43.0 | 35.27 ±3.693 |
| 8 Mar 1988 | 641 | 22.0–41.5 | 34.09 ±3.688 |
| 19 Apr 1988 | 675 | 26.5–44.5 | 36.49 ±3.176 |
| 3 Jun 1988 | 720 | 31.0–49.0 | 40.18 ±3.361 |

growth rate observed at Unalaska Island was most similar to that at Kodiak Island, and very little seasonal signature could be detected. Estimated mean size at age 2 at Unalaska Island (37.6 mm CL) was much closer to the value estimated for Kodiak Island (42.2 mm CL) than for Bristol Bay (22.7 mm CL, Table 5); at age 3, Bristol Bay crabs were expected to average ~20 mm smaller than crabs observed at Unalaska Island. It is important to note that the choice of settlement dates for the models had negligible effect on size-at-age estimates, except where such changes caused cohorts to be re-assigned to younger age classes than seemed reasonable (i.e. if early spring settlement dates had been chosen, very small crabs would have been assigned to late age 0, instead of being considered immediately postsettlement crabs). Changes in settlement date primarily affected estimated size-at-age over the first few months but had little impact on estimation of size at 1, 2, or 3 years after settlement.

**Table 4**

Comparison of coefficient of determination ($r^2$) goodness-of-fit values associated with seasonalized Gompertz and von Bertalanffy growth curves fitted to early juvenile red king crab size-at-age data.

| Region | Gompertz | von Bertalanffy |
|---|---|---|
| Bristol Bay | 0.968 | 0.842 |
| Unalaska Island | 0.922 | 0.863 |
| Kodiak Island | 0.995 | 0.919 |

## Late juvenile through early reproductive-age crabs

Male red king crabs collected in NMFS trawl surveys from 1975 to 1999 ranged from 6 to 201 mm CL, and all sizes between 23 and 198 mm CL were observed. Female red king crabs ranged from 7 to 192 mm CL; all sizes between 24 and 165 mm CL. Over the entire time series, two particularly strong year classes recruited to the Bristol Bay population whose growth and size-at-age characteristics could be tracked by modal size progression. The first year class was evident from 1979 to 1984, first appearing as size modes with mean CL = ~32 mm and

~30 mm for males and females, respectively (Fig. 5, Table 6); these crabs were most likely ~2.9 years of age and represented the year class that settled in 1976. Males of this year class grew to a mean CL = ~88 mm by 1983 and the mode became indistinguishable in 1984; females grew to a mean CL = ~84 mm by 1983, and ~92 mm by 1984 (Fig. 5, Table 6). The second strong year class was evident from 1994 to 1999 (Fig. 6, Table 6) and first appeared at larger sizes than did the 1976 year class. Males first appeared at a mean CL = ~54 mm and grew to a mean CL = ~137 mm in 1999; females first appeared at ~53 mm CL in 1994 and grew to ~109 mm in 1999. This year class first appeared at sizes that were essentially equivalent to the sizes that individuals from the 1976 year class had attained at age 3.9. Thus, this second year class most likely represented crabs that settled in 1990.

For both year classes, growth of males and females was similar up to ~85 mm CL, after which females grew more slowly than males. However, the growth rates of the two year classes were not equivalent: the 1976 year class grew slower than the 1990 year class (Fig. 7) because the 1976 year class required two years (from 1980 to 1982) to progress from a mean CL = ~50 mm to mean CL = ~70 mm, whereas the 1990 year class achieved this level of growth within a single year (from 1994 to 1995). Mean sizes during the following two years of growth (from 1982 to 1984 for the 1976 year class; 1995 to 1997 for the 1990 year class, Table 6) were similar between year classes.

Within all of the annual length-frequency distributions, considering only size modes with mean CL <100 mm, 24 modes were identified for both males and for females. These included the modes presented previously for the 1976 and 1990 year classes, and an additional 16 modes for males, and 14 modes for females. The additional modes represent other year classes whose modal progression could not be tracked for a substantial period of time. The mean CL (±1 SD) of all 24 male and female modes is presented in Figure 8, plotted sequentially by increas-

**Table 5**

Characteristics of seasonalized Gompertz growth curves fitted to size-at-age data from Bristol Bay, Unalaska Island, and Kodiak Island. Values are reported for ages 0.9, 1.9, and 2.9 because these ages roughly correspond to trawl survey data: the trawl survey typically occurs in late May, ~0.9 years following the previous year's settlement. For all curves, $L_{max}$ = 200 mm CL. See "Materials and methods" section for definitions of model parameters.

| Model parameters | Bristol Bay | Unalaska Island | Kodiak Island |
|---|---|---|---|
| $r^2$ (fit) | 0.968 | 0.922 | 0.995 |
| $K$ | 0.415 | 0.421 | 0.634 |
| $C$ | 1.000 | 0.275 | 0.599 |
| $t_s$ (year) | 0.553 | 0.535 | 0.680 |
| $t_{min}$ (year) | –3.922 | –3.286 | –2.659 |
| Length estimates from the models | | | |
| age 0.0 | 3.1 | 3.6 | 2.7 |
| age 0.9 | 7.0 | 14.5 | 9.9 |
| age 1.0 | 8.5 | 16.4 | 11.7 |
| age 1.9 | 19.5 | 34.5 | 38.0 |
| age 2.0 | 22.7 | 37.6 | 42.2 |
| age 2.9 | 41.9 | 62.7 | — |
| age 3.0 | 46.7 | 66.4 | — |
| Slowest growth in | January | January | February |

**Table 6**

Summary of mean size (±1 SD) of the cohorts depicted in size-frequency progression plots (Figs. 5 and 6), as determined with FiSAT.

| Year class | Year | Male mean size (mm) ±1 SD | Female mean size (mm) ±1 SD | Age (years after settlement) |
|---|---|---|---|---|
| 1976 | 1979 | 31.9 ±2.79 | 29.6 ±2.04 | 2.9 |
| | 1980 | 50.7 ±4.10 | 50.2 ±3.40 | 3.9 |
| | 1981 | 63.2 ±5.06 | 64.0 ±5.73 | 4.9 |
| | 1982 | 70.1 ±7.50 | 71.4 ±8.20 | 5.9 |
| | 1983 | 88.4 ±14.74 | 83.5 ±11.23 | 6.9 |
| | 1984 | cohort indistinct | 92.3 ±4.84 | 7.9 |
| 1990 | 1994 | 54.1 ±4.67 | 52.7 ±3.75 | 3.9 |
| | 1995 | 73.4 ±4.90 | 71.9 ±7.28 | 4.9 |
| | 1996 | 86.5 ±8.01 | 83.5 ±5.89 | 5.9 |
| | 1997 | 104.2 ±8.44 | 97.4 ±3.62 | 6.9 |
| | 1998 | 117.8 ±11.40 | 105.3 ±8.01 | 7.9 |
| | 1999 | 136.5 ±11.92 | 109.3 ±7.79 | 8.9 |




**Figure 5**

Yearly length-frequency histograms for male (left) and female (right) Bristol Bay region red king crabs <130 mm CL, 1979–1984, revealing mean size-at-age of the 1976 year class by modal progression. Arrows indicate the cohort's mean CL each year, as determined with FiSAT; dark shading indicates CLs within 1 SD of the mean. The vertical lines located at 105 mm CL and 97 mm CL indicate size at reproductive recruitment for males and females, respectively, as defined by Zheng et al. (1995a, 1995b). Note that the 1979 histograms have been truncated at $n = 80$ observations in order to make the relevant size modes more visible. The 1984 male size mode was not resolved with FiSAT; we considered this size mode to be too indistinguishable from the surrounding data to yield an accurate result.

ing mean CL. This figure represents an attempt to identify commonly occurring sizes that may represent the expected CLs of consecutive age classes. For male crabs, a cluster of three observations occurred at ~35 mm CL, separated from the remaining observations, and likely representing a single age class; for females, two observations occurred at the same approximate size, separated from the remaining modes. Considering the estimated age




**Figure 6**

Yearly length-frequency histograms for male (left) and female (right) Bristol Bay region red king crabs <155 mm CL, 1994–1999, revealing mean size-at-age of the 1990 year class with modal progression. Arrows indicate the cohort's mean CL each year, as determined with FiSAT, and dark shading indicates CLs within 1 SD of that mean. The vertical lines located at 105 mm CL and 97 mm CL indicate size-at-reproductive recruitment for males and females, respectively, as defined by Zheng et al. (1995a, 1995b).

at size 2.9 of ~40 mm obtained from the growth curve presented earlier (Fig. 4, Table 5), these small crab are likely age 2.9. At larger sizes, there was considerable overlap between modes, suggesting that the mean CL of older age classes is quite variable and difficult to distinguish among years.

## Bottom temperature

From 1975 to 1999, area-averaged near-bottom temperatures in Bristol Bay ranged between a low of 0.7°C in 1976 and a high of 5.2 °C in 1981 (Fig. 9). In general, the coldest temperatures were observed in the 1970s, warmest tem-




**Figure 7**

Best-fit normal curves, as determined with FiSAT, of the length-frequency modes associated with male and female red king crabs of the 1976 (solid curves) and 1990 (dashed curves) year classes, from age 3.9 to 6.9, demonstrating the divergence in size-at-age between the two year classes. In particular, note the large difference in apparent molt increment displayed by the year classes between ages 3.9 and 4.9. All curves have been standardized to a uniform height in order to facilitate their comparison, because the number of crabs that were observed differed between years.

peratures during the early 1980s, and moderate temperatures from the mid-1980s through the late 1990s.

## Discussion

Our analyses demonstrate that Bristol Bay red king crab grow slower than previously assumed. The present stock-recruitment relationship used to manage this population is based on growth models (Weber, 1967; Balsiger, 1974; Stevens and Munk, 1990) that suggest a time lag of six years between settlement and subsequent recruitment (i.e. seven years after fertilization), where reproductive recruitment is defined to occur at ~97 mm and ~105 mm CL for females and males, respectively (Zheng et al., 1995a, 1995b). Our results indicate that mean age at full reproductive recruitment is likely 8–9 years after settlement. Crabs that settled in 1990 began to reach reproductive size in 1997, at 7 years after settlement, and at this age only ~50% of the individuals were at or above the reproductive size cut-off; the mode did not become fully recruited for an additional 1 to 2 years. Individuals of the 1976 year class grew even slower and, on average, were still slightly smaller than reproductive size at 8 years after settlement. This year class would not have begun to contribute substantially to the reproductive stock until at least 9 years after settlement.

The discrepancy between our estimates of mean age-at-recruitment and presently accepted values is due in part to incorrect assumptions of the latter regarding the growth of early juveniles in Bristol Bay. Weber (1967) conducted one of the few comprehensive *in situ* studies of early juvenile size-at-age in Alaskan waters and clearly demonstrated that red king crab should be expected to reach a mean CL = ~66 mm CL three years after settlement. This conclusion contrasted sharply with earlier data from Bristol Bay that showed strong size modes with means of 4 mm, 9 mm, and 17 mm CL in early summer samples in 1956, 1957, and 1958, respectively (Fisheries Agency of Japan[6]), apparently representing the modal progression of the 1956 year class, and suggesting a much slower growth rate. More recent studies conducted in Kodiak (Dew, 1990; Donaldson et al., 1992) have supported Weber's (1967) conclusions; thus, the observations made by the Fisheries Agency of Japan[6] have been largely ignored. Our results suggest that the data from Kodiak is in good accord with Weber's (1967) conclusions, but that mean CLs of 4 mm,

[6] Fisheries Agency of Japan. 1959. Report of research on king crab in the eastern Bering Sea. Int. N. Pac. Fish. Comm. Annu. Rep., p. 71–78. [Available from Secretariat, North Pacific Anadromous Fish Commission, Suite 502, 889 West Pender Street, Vancouver, British Columbia, V6C 3BC.]




**Figure 8**

Mean CL ±1 SD of all strong size modes of Bristol Bay red king crabs appearing in the annual length-frequency data from NMFS Bering Sea trawl surveys, 1975–1999; only modes with mean CL <100 mm are plotted. Size modes are plotted in length-sequential order along the *x*-axis, starting with the smallest observed mean CL. The numbers adjacent to each mode indicate the year during which that mode was observed. A grouping of modes separate from the remaining data, indicated by the dashed line and the arrow, likely represents a discrete age class; these small crabs are most likely age 2.9 (refer to Fig. 4). The remaining observations are continuous; substantial size overlap between consecutive modes suggest that discrete size-at-age categories are lacking at >~40 mm mean CL.




**Figure 9**

Area-averaged June bottom temperatures in the Bristol Bay region from 1975 to 1999. The five-year running averages represent a five-year period that includes the plotted date and the four years preceding it; for example, the five-year average temperature plotted at 1981 (3.0°C) represents mean area-averaged temperature from 1977 to 1981.

9 mm, and 17 mm are consistent with sizes at age 0, 1, and 2, respectively, within Bristol Bay. This may be due, in part, to reduced molt frequency in Bristol Bay: the data obtained from Weber (1967) indicate that mid-winter molting was not uncommon during his study, whereas our data suggest that this may not be typical in Bristol Bay. It is difficult to determine whether the discrepancies represent actual regional differences, or simply differences between different studies conducted at different sites and at different times, but our results indicate that it is probably inappropriate to apply the early juvenile growth rate obtained by Weber (1967) to the Bristol Bay stock; use of the faster growth rate likely results in lower estimates of age-at-recruitment than the population displays.

Variable growth of older juveniles (age 3+) may further delay age-at-recruitment. This variability is evident when comparing growth of the 1976 year class to that of the 1990 year class; age-at-recruitment was 1–2 years greater in the former, due to slow growth of prerecruits. In particular, note that females of the 1976 year class, averaging 83.5 mm CL in 1983, displayed a mean increase in carapace length during molting (molt increment [MI]) of ~9 mm during the 1984 spring molt, whereas females of the 1990 year class, also averaging 83.5 mm CL in 1996, exhibited a mean MI of ~14 mm during the 1997 spring molt. As a result, females from the 1976 year class required four years to grow from a mean CL = ~50 mm to a mean CL = ~92 mm CL and at ~8 years after settlement were still slightly smaller than the estimated size for full reproductive recruitment (Zheng et al., 1995a, 1995b). Females from the 1990 year class were able to accomplish slightly more mean growth, from ~53 mm to ~97 mm CL, in only three years.

The large difference in growth rate between the 1976 and 1990 year classes may have been caused by water temperature differences during the two time periods. Molt schedules and growth rates can be strongly influenced by ambient temperature (Kurata, 1960, 1961; Nakanishi, 1985), and considerable variability in size at maturity has been observed over the species' geographic range in both males (Paul et al., 1991) and females (see review in Blau, 1990; Otto et al., 1990). Though a number of factors may contribute to the observed variability, reduced growth associated with colder bottom temperatures has been invoked to explain the smaller size-at-maturity observed in the Norton Sound population as compared with other Bering Sea stocks (Blau, 1990; Otto et al., 1990), and modeling suggests that regional and temporal variation in temperature can have broad effects on age-at-recruitment (Stevens, 1990; Stevens and Munk, 1991). June bottom temperature profiles in Bristol Bay suggest that the 1976 year class was subjected to lower temperatures than the 1990 year class, primarily at early juvenile ages. Although a detailed analysis of temperature-dependent growth would require year-round temperature records, which are not available for this region, June temperatures in Bristol Bay may serve as a proxy for thermal conditions throughout the year. Bottom temperatures in Bristol Bay are linked to seasonal sea ice, that in some years covers much of Bristol Bay (NIC, 1994; Wyllie-Echeverria, 1995; Neibauer, 1998), and the development of sea ice can have strong effects on bottom temperature conditions throughout the year. "Cold pool" bottom waters (<1°C) produced in the winter during ice formation may persist well into the summer, and potentially into the following winter, once insulated from surface heating by the development of the summer thermocline (Azumaya and Ohtani, 1995).

In many Crustacea, temperature primarily affects the molt schedule and has little influence on the magnitude of the MI (Hartnoll, 1982; Wainright and Armstrong, 1993), but laboratory studies conducted with red king crab indicate substantial variability in MI-at-age, across ranges of temperatures, as well as under stable environmental conditions. Rearing crabs ~6.3 mm CL under constant temperatures of ~10°C, Molyneaux and Shirley (1988) reported changes in CL at molt that ranged from –4.4% to 52.2%; similarly, for juvenile premolt crabs 33–36 mm CL, reared at ~5.0°C, Gharrett (1986) observed MIs ranging from 3 to 8 mm. At reproductive age, Weber and Miyahara (1962) observed that MI varied between 5 and 23 mm CL per molt in adult males, and large variability in MI associated with water temperatures between 0° and 12°C has been demonstrated for ovigerous females (Shirley et al., 1990).

Because the changes in mean CL the we observed for the Bristol Bay stock were determined through modal analysis of the entire population, it is reasonable to suspect that apparent differences in growth between years and cohorts do not represent MI variability but may be explained as variability in the number of molted versus unmolted crab within particular survey years. This is reasonable to assume, considering that red king crab may skip molting so that the annual molt schedule is replaced by a biennial or triennial cycle (Weber and Miyahara, 1962; McCaughran and Powell, 1977; Balsiger, 1974). However, closer examination of the trawl survey data indicates that, of the 23 size modes of crab observed, none comprised less than 94% new-shelled crabs that had recently molted (senior author, unpubl. data). A high proportion of newly molted crabs was characteristic of nearly all the identifiable size modes (senior author, unpubl. data): of the 53 modes identified, 35 comprised entirely new-shelled crabs, 17 comprised 94–99% new-shelled crabs, and only one comprised >10% old-shell individuals (the female cohort with mean CL=~94mm CL in 1982; population=11.4% old-shell). Thus, the difference in growth rates observed between the two year classes of late juveniles cannot be explained by variations in molt frequency; the trawl survey data support the conclusion that variability in MI is a characteristic shared by both sexes across a range of ages.

Substantial variability in MI is an important life history characteristic that confounds attempts to assign discrete size-at-age categories to a population. The greater the variability in MI among individuals and over time, the greater will be the range of sizes associated with crabs in a given year class, making modes more difficult to resolve from one another. Size-at-age values presently used for Bristol Bay red king crabs were derived from studies consisting of 1–4 years of data (Weber and Miyahara, 1962; Weber, 1967; McCaughran and Powell, 1977; Incze et al., 1986), but our analyses show that a strong tenden-

cy toward specific mean size-at-age is not apparent if longer time scales are considered. Size-at-age characteristics may be different depending on which year class is considered, and among crabs >~40 mm CL, mean CLs of identifiable cohorts displayed a fairly continuous distribution with considerable overlap between adjacent size modes; we could not identify specific mean CLs that could be consistently assigned to various age classes. In addition to confounding age estimates, variable MI may cause different year classes to recruit at different ages, over different time spans, and increase the number of year classes that constitute each year's new recruitment. These issues have been treated elsewhere with respect to variable intermolt period (Stevens, 1990); variability in MI will produce the same effects.

We explored the possibility that overlap between size modes might be attributable to changes in growth rate within the population over time. That is, because bottom temperatures in Bristol Bay were colder during the 1970s than they have been more recently, growth was expected to be slower in the 1970s than later in the time series (Stevens, 1990). Thus, one might expect size modes to fall closer together early in the time series and to be spaced farther apart later. However, we were unable to resolve a clear temporal component in the data; even consecutive year classes sometimes had different mean size-at-age and growth increment characteristics. For example, the 1976 year class displayed a mean CL = ~51 mm (range: 44–57 mm) and ~63 mm CL (range: 56–71mm) in 1980 and 1981, respectively. These represented clear and well-separated size modes at ages 3.9 and 4.9, respectively. However, note the occurrence of a strong mode in the 1979 data, with a mean CL = ~58mm CL (range=50–65 mm); this mode probably represents a single year class, settled in 1975, which would be expected to have mean size-at-age characteristics similar to those of the 1976 year class. Yet, this mode of the 1975 year class fell almost precisely between, and its range encompassed the mean sizes of both age 3.9 and age 4.9 crabs from the 1976 year class. Such features were not uncommon in the length frequencies that were presented. For applications requiring accurate size-at-age information, the onerous task of year-by-year and year-class-by-year-class assessments may be necessary.

In summary, our results demonstrate that both male and female red king crab in Bristol Bay reach maturity at least one year later than presently assumed (i.e. at ~7 years after settlement) due to slower growth from settlement through age 3. Furthermore, variability in MI in late juveniles can result in further reduction in growth rate such that reproductive recruitment is delayed by an additional 1–2 years (i.e. reproductive recruitment at ~8–9 years after settlement). From a management perspective, variable MI is a life-history characteristic that should be considered in growth- and length-based models of recruitment. Present models attempt to simulate MI variability (Zheng et al., 1995a), but little information exists on the magnitude of that variability and its changes over time and space. Such information will be valuable to managers to calibrate recruitment models with respect to lag times between spawning and subsequent recruitment, as well as to predict how year classes enter the spawning population and the fishery; more research in this area is warranted. Inappropriate growth rate and lag-time assumptions have resulted in assigning the wrong year's spawning stock biomass to subsequent recruitment levels in the Bristol Bay stock-recruitment curve; spawning stock abundances are presently offset ~1–2 years from the recruitment levels that they generated. This offset may affect the precise shape of the stock-recruitment curve and alter some of the models associated with it. Such changes may prove negligible with respect to actual harvest strategies, but it may be prudent to make the appropriate adjustments given knowledge of greater age at reproductive recruitment.

The Bristol Bay red king crab stock has been typified by large fluctuations in fishable abundance and by relatively rare, strong recruitment pulses generating the bulk of the fishery. Most recently, relatively strong catches from 1997–99, yielding a combined landed catch of ~35 million pounds of crabs with an exvessel value estimated at over $137 million (Morrison et al.[7]), were supported almost entirely by the 1990 year class (i.e. were spawned by the 1989 reproductive stock). Former assumptions would have led us to assign this pulse to the 1990 spawning stock and assume that the planktonic larval phase and settlement occurred during 1991. From a management standpoint, the ramifications of such an error may be negligible because estimated effective spawning biomass was similar in both 1989 and 1990 (Zheng and Kruse[1]). However, as we try to elucidate the mechanisms that generated this strong year class, it is crucial that we accurately determine when those crabs were larvae, early benthic individuals, and later stage crabs. Physical forcing, for example, has been shown to play a large role in determining recruitment variability in a number of commercially important crustacean species worldwide (e.g. Polovina and Mitchum, 1992; Polovina et al., 1993; McConnaughey et al., 1994; Rothlisberg et al., 1994; Jones and Epifanio, 1995; McConnaughey and Armstrong, 1995; Rozenkranz et al., 1998). Similar correlations between recruitment and physical parameters have been attempted for king crabs (Zheng and Kruse, 2000), but our ability to identify causes of recruitment variability relies upon associating the correct life history stages with physical forcing events. Even seemingly minor errors in growth-rate assumptions can have serious impacts on our understanding of population dynamics.

## Acknowledgments

Funding for preparation of this manuscript was provided by the University of Washington, Seattle, through a Victor and Tamara Loosanoff Fellowship. We wish to thank Julie

[7] Morrison, R., F. Bowers, R. Gish, E. Wilson, W. Jones, and B. Palach. 2000. Annual management report for shellfish fisheries of the Bering Sea. *In* Annual management report for shellfish fisheries of the westward region, 1999, p. 147–261. Regional Information Report 4KOO-55, Kodiak. Division of Commercial Fisheries, Alaska Department of Fish and Game, P.O. Box 25526, Juneau, Alaska, 99801.

Keister, P. Sean MacDonald, and our anonymous reviewers for thoughtful suggestions that improved the quality of this manuscript, and Steven Hare (International Pacific Halibut Commission) and Gary Walters (NMFS, AFSC) for providing near-bottom temperature data.

## Literature cited

Anastacio, P. M., and J. C. Marques.
1995. Population biology and production of the red swamp crayfish *Procambarus clarkii* (Girard) in the lower Mendego River Valley, Portugal. J. Crustacean Biol. 15(1):156–168.

Azumaya, T., and K. Ohtani.
1995. Effect of winter meteorological conditions on the formation of the cold bottom water in the eastern Bering Sea Shelf. J. Oceanogr. 51:665–680.

Balsiger, J. W.
1974. A computer simulation model for the eastern Bering Sea king crab population. Ph.D. diss., School of Fisheries, Univ. Washington, Seattle, WA, 197 p.

Bhattacharya, C. G.
1967. A simple method of resolution of a distribution into Gaussian components. Biometrics 23:115–135.

Blau, S. F.
1990. Size at maturity of female red king crabs (*Paralithodes camtschatica*) in the Adak management area, Alaska. *In* Proceedings of the international symposium on king and Tanner crabs, p. 105–116. Univ. Alaska Sea Grant Report 90-04, Anchorage, AK.

Cressie, N. A. C.
1993. Statistics for spatial data. John Wiley and Sons, New York, NY, 900 p.

Dew, B.
1990. Behavioral ecology of podding red king crab, *Paralithodes camtschatica*. Can. J. Fish. Aquat. Sci. 47:1944–1958.

Donaldson, W. E., S. C. Beyersdorfer, D. Pengilly, and S. F. Blau.
1992. Growth of red king crab, *Paralithodes camtschaticus* (Tilesius, 1815), in artificial habitat collectors at Kodiak, Alaska. J. Shellfish Res. 11(1):85–89.

Gayanilo, F. C., Jr., and D. E. Pauly.
1997. FAO-ICLARM stock assessment tools (FiSAT): reference manual. FAO Computerized Information Series (Fisheries) 8, FAO, Rome, 262 p.

Gharrett, J. A.
1986. Effects of a hydrocarbon-contaminated diet on survival, feeding, growth, and molting of juvenile red king crab, *Paralithodes camtschatica* (Tilesius). M.S. thesis. Univ. Alaska, Juneau, AK, 178 p.

Hartnoll, R. G.
1982. Growth. *In* The biology of Crustacea, volume 2: embryology, morphology, and genetics (L. B. Ebele, ed.), p. 111–196. Academic Press. New York, NY.

Hasselblad, V.
1966. Estimation of parameters for a mixture of normal distributions. Technometrics 8:431–444.

Hill, J., D. L. Fowler, and M. J. VanDenAvyle
1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (mid-Atlantic): blue crab. Biol. Rep. US Fish Wildl. Serv., 27 p.

Huner, J. V., and R. P. Romaire.
1990. Crawfish culture in the southeastern USA. World Aquaculture 21:58–66.

Incze, L. S., R. S. Otto, and M. K. McDowell.
1986. Recruitment variability of juvenile red king crab, *Paralithodes camtschatica*, in the southeast Bering Sea. *In* North Pacific workshop on stock assessment and management of invertebrates (G. S. Jamieson, and N. Bourne, eds.), p. 370–378. Can. Spec. Pub. Fish. Aquat. Sci. 92.

Jones, M. B., and C. E. Epifanio.
1995. Settlement of brachyuran megalopae in Delaware Bay: an analysis of time series data. Mar. Ecol. Prog. Ser. 125:67–76.

Keckler, D.
1994. Surfer for Windows: user's manual. Golden Software, Golden, CO, 428 p.

Kurata, H.
1960. Studies on the larvae and postlarvae of *Paralithodes camtschatica*. III. The influence of temperature and salinity on growth of the larvae. Bull. Hokkaido Reg. Fish. Res. Lab. 17:9–14.
1961. Studies on the larvae and postlarvae of *Paralithodes camtschatica*. IV. Growth of the postlarvae. Bull. Hokkaido Reg. Fish. Res. Lab. 18:1–9.

Loher, T., and D. Armstrong.
2000. Effects of habitat complexity and relative larval supply on the establishment of early benthic phase red king crab (*Paralithodes camtschaticus* Tilesius, 1815) populations in Auke Bay, Alaska. J. Exp. Mar. Biol. Ecol. 245(1):83–109.

Loher, T., P. S. Hill, G. A. Harrington, and E. Cassano.
1998. Management of Bristol Bay red king crab: a critical intersections approach to fisheries management. Rev. Fish. Sci. 6(3):169–251.

McCaughran, D. A., and G. C. Powell.
1977. Growth model for Alaskan king crab (*Paralithodes camtschatica*). J. Fish. Res. Board Can. 34: 989–995.

McConnaughey, R. A., and D. A. Armstrong.
1995. Potential effects of global climate change on Dungeness crab (*Cancer magister*) populations in the northeast Pacific Ocean. *In* Climate change and northern fish populations (R. J. Beamish, ed.), p. 291–306. Can. Spec. Pub. Fish. Aquat. Sci. 121.

McConnaughey, R. A., D. A. Armstrong, B. M. Hickey, and D. R. Gunderson.
1994. Interannual variability in coastal Washington Dungeness (*Cancer magister*) populations, larval advection and the coastal landing strip. Fish. Oceanogr. 3:22–38.

Molyneaux, D. B., and T. C. Shirley.
1988. Molting and growth of eyestalk-ablated juvenile red king crabs, *Paralithodes camtschatica* (Crustacea: Lithodidae). Comp. Biochem. Physiol. 91:245–251.

Nakanishi, T.
1985. The effect of the environment on the survival rate, growth and respiration of eggs, larvae and post-larvae of king crab (*Paralithodes camtschatica*). *In* Proceedings of the international king crab symposium, p. 167–185. Univ. Alaska Sea Grant Report 85-12, Anchorage, AK.

Neibauer, H. J.
1998. Variability on Bering Sea ice cover as affected by a regime shift in the north Pacific in the period 1947–1996. J. Geophys. Res. 103(C12):27717–27737.

NIC (National Ice Center).
1994. Navy/NOAA National Ice Center (NIC) weekly sea ice concentrations and extents, 1972–1994. National Ice Center, Washington, D.C. [Compact disc.]

Otto, R. S.
1986. Management and assessment of Bering sea king crab stocks. *In* North Pacific workshop on stock assessment and management of invertebrates (G. S. Jamieson and N. Bourne, eds.), p. 195–227. Can. Spec. Pub. Fish. Aquat. Sci. 92.

Otto, R. S., R. A. MacIntosh, and P. A. Cumminskey.
1990. Fecundity and other reproductive parameters of female red king crab (*Paralithodes camtschatica*) in Bristol Bay and Norton Sound. *In* Proceedings of the international symposium on king and Tanner crabs, p. 65–90. Univ. Alaska Sea Grant Report 90-04, Anchorage, AK.

Paul, J. M., A. J. Paul, R. S. Otto, and R. A. MacIntosh.
1991. Spermatophore presence in relation to carapace length for eastern Bering Sea blue king crab (*Paralithodes platypus*, Brandt, 1850) and red king crab (*Paralithodes camtschaticus* (Tilesius, 1815)). J. Shellfish Res. 10:157–163.

Polovina, J. J., W. R. Haight, R. B. Moffit, and F. A. Parrish.
1993. The role of benthic habitat, oceanography, and fishing on the population dynamics of the spiny lobster, *Panulirus marginatus* (Decapoda, Palinuridae), in the Hawaiian Achipelago. *In* Proceedings of the fourth annual workshop on Lobster Biology and Management. Crustaceana 68:203–212.

Polovina, J. J., and G. T. Mitchum
1992. Variability in spiny lobster recruitment and sea level in the northwestern Hawaiian Islands. Fish. Bull. 90:483–493.

Rothlisberg, P. C., J. A. Church, and C. B. Fandry.
1994. A mechanism for near-shore concentration and estuarine recruitment of post-larval *Penaeus plebejus* Hess (Decapoda, Penaeidae). Estuarine Coastal Shelf Sci. 40:115–138.

Rozenkranz, G. E., A. V. Tyler, G. H. Kruse, and H. J. Neibauer.
1998. Relationship between wind and year class strength of Tanner crabs in the southeast Bering Sea. Alaska Fish. Res. Bull. 5:18–24.

Shirley, T. C., S. M. Shirley, and S. Korn.
1990. Incubation period, molting and growth of female red king crabs: effects of temperature. *In* Proceedings of the international symposium on king and Tanner crabs, p. 51–63. Univ. Alaska Sea Grant Report 90-04, Anchorage, AK.

Stevens, B. G.
1990. Temperature-dependent growth of juvenile red king crab (*Paralithodes camtschatica*) and its effect on size-at-age and subsequent recruitment in the eastern Bering Sea. Can. J. Fish. Aquat. Sci. 47:1307–1317.

Stevens, B. G., and J. E. Munk.
1990. A temperature-dependent growth model for juvenile red king crab, *Paralithodes camtschatica*, in Kodiak, Alaska. *In* Proceedings of the international symposium on king and Tanner crabs, p. 293–304. Univ. Alaska Sea Grant Report 90-04, Anchorage, AK.

Tremblay, M. J., and M. D. Eagles.
1997. Molt timing and growth of the lobster, *Homarus americanus*, off northeastern Cape Breton Island, Nova Scotia. J. Shellfish Res. 16:383–394.

Wainwright, T. C., and D. A. Armstrong.
1993. Growth pattern in Dungeness crab (*Cancer magister* Dana): synthesis of data and comparison of models. J. Crustacean Biol. 13:36–50.

Weber, D. D.
1967. Growth of the immature king crab *Paralithodes camtschatica* (Tilesius). Bull. Int. N. Pac. Fish. Comm. 21:21–53.

Weber, D. D., and T. Miyahara.
1962. Growth of the adult male king crab *Paralithodes camtschatica* (Tilesius). Fish. Bull. 62:53–75.

Wyllie-Echeverria, T.
1995. Sea ice conditions and the distribution of walleye pollock (*Theragra chalcogramma*) on the Bering and Chukchi Sea Shelf. *In* Climate change and northern fish populations (R. J. Beamish, ed.), p. 131–136. Can. Spec. Publ. Fish. Aquat. Sci. 121.

Zheng, J. M., and G. H. Kruse.
2000. Recruitment patterns of Alaskan crabs in relation to decadal shifts in climate and physical oceanography. ICES J. Mar. Sci. 57:438–451.

Zheng, J., M.C. Murphy, and G. H. Kruse.
1995a. A length-based population model and stock-recruitment relationships for red king crab, *Paralithodes camtschaticus*, in Bristol Bay. Can. J. Fish. Aquat. Sci. 52:1229–1246.
1995b. Updated length-based population model and stock-recruitment relationship for red king crab in Bristol Bay, Alaska. Alsk. Fish. Res. Bull. 2:114–124.

# Age, growth, and mortality of California halibut, *Paralichthys californicus*, along southern and central California

**Leslie S. MacNair**
**Michael L. Domeier**
**Calvin S. Y. Chun**

California Department of Fish and Game
4665 Lampson Avenue, Suite C
Los Alamitos, California 90720

E-mail address (for C. S. Y. Chun, contact author): cchun@dfg.ca.gov

**Abstract**–California halibut, *Paralichthys californicus*, collected by a 400-mesh eastern trawl in southern (Mexican border to Point Conception) and central (Point Conception to Tomales Bay) California were aged by using whole and sectioned otoliths (sagittae) to determine age, growth, and mortality. Males represented 69% of the sample from southern California and 53% of the sample from central California. A higher proportion of California halibut were older in southern California than in central California. Although California halibut can live as long as 30 years, the oldest fish found in our study was 13 years old. For southern California, the von Bertalanffy growth function (VBGF) was $L_t=925.3(1-e^{-0.08(t+2.2)})$ for males and $L_t=1367.7(1-e^{-0.08(t+1.2)})$ for females. For central California, the VBGF was $L_t=956.7(1-e^{-0.10(t+2.1)})$ for males and $L_t=1477.1(1-e^{-0.10(t+0.2)})$ for females. The VBGF showed that at the same age, females on average were larger than males in both southern and central California. The VGBF also showed that both male and female halibut in central areas on average were larger than halibut from southern California. Instantaneous mortality rates of halibut in southern California were estimated at 0.91 for males and 0.68 for females. Mortality estimates for central California could not be calculated because of small sample sizes.

Manuscript accepted 9 March 2001.
Fish. Bull. 99:588–600 (2001).

California halibut, *Paralichthys californicus*, is an important flatfish species for sport and commercial fisheries in nearshore waters off central and southern California (Frey, 1971). The gear most commonly used commercially to harvest California halibut are otter trawls, gill nets, and trammel nets, whereas hook-and-line or spear are used mostly in the recreational fishery (Kramer and Sunada, 1992).

Information on age, growth, and mortality of California halibut are important to the management of both the sport and commercial halibut fisheries. Previous studies have addressed these issues to a limited extent. Pattison and McAllister (1990) studied techniques to age halibut and determined length-at-age of halibut along the California coast, but most of the fish were taken south of Point Conception and they did not do separate analyses for southern and central California. Sunada et al. (1990) analyzed age, size, and sex composition of commercial landings in southern California. Haaker (1975) studied the biology of California halibut in Anaheim Bay, including age and growth, but the fish he used were generally less than three years of age, immature, and included fish only to 510 mm total length. In contrast to other studies, our study provides information on age and growth for central and southern California separately. Mortality for southern California was also calculated.

## Methods

Southern California was defined as the area between the U.S.-Mexico border and Point Conception, and central California as the area between Point Conception and Tomales Bay. Bottom trawl surveys were conducted off central California from 8 July through 3 August 1993 and off southern California from 14 February through 18 March 1994. California halibut were caught with a 400-mesh eastern trawl (15 m wide × 1.5 m high; 9.8 cm mesh body, 8.5 cm mesh codend). For both areas, sampling effort was stratified by depth: 0–20 fathoms (fm), 21–40 fm, and 41–60 fm. Fifty stations within each depth stratum were randomly selected for sampling by trawling parallel to isobaths at each station. The sample sizes were not proportional to the estimated area of the strata (Table 1).

An additional ten California halibut were collected with the same 400-mesh eastern trawl gear in southern California from July 1994 to June 1995. These were included in the age and growth determinations. However, they were not used for mortality estimates because they were collected at a different time period from the other halibut collected off southern California.

Each halibut was measured to the nearest millimeter for total length (TL) and standard length (SL). Gonads were examined visually to determine the sex of the fish. Sagittal otoliths were removed and stored dry in vials for age determination. Pattison and McAllister (1990) determined that when compared with other fish structures, otoliths provided the most reliable ages for California halibut. They found that in an otolith, an opaque band was formed during spring and summer (from April to October) and a translucent band was

**Table 1**

Number of trawls and number of California halibut used in age and growth analyses by depth strata and region. Asterisk (*) indicates that these fish were not used in the mortality estimates because they were caught at a later date.

| | Southern California | | | | Central California | | |
|---|---|---|---|---|---|---|---|
| Stratum | Depth (fm) | No. of trawls | No. of halibut used | Area (nmi$^2$) | No. of trawls | No. of halibut used | Area (nmi$^2$) |
| 1 | 0–20 | 58 | 913 +3* | 575 | 47 | 254 | 587 |
| 2 | 21–40 | 46 | 165 +2* | 495 | 39 | 22 | 905 |
| 3 | 41–60 | 45 | 13 +1* | 425 | 40 | 0 | 1124 |
| Total | | | 1091 +6* | | | 276 | |

deposited in winter (from November to March). A combination of an inner opaque and adjacent outer translucent zone represents an annular growth ring or annulus (Casselman, 1983). This growth ring was validated by Pattison and McAllister (1990) with oxytetracycline-marked otoliths; they found that only one complete ring (opaque and translucent) was formed each year.

Pattison and McAllister suggested that age estimates of older fish (age 11 or older) may be underestimated when whole otoliths, rather than sectioned otoliths, are read. In the preliminary analysis in our study, to test if a significant difference existed between estimating age with whole otoliths and estimating age with sectioned otoliths, an otolith from each of 80 randomly selected fish (10 fish from each 100-mm size class from 200 to 900 mm TL) was viewed both whole and sectioned to estimate age. A mutually determined age (an age agreed upon by the two readers) was established for the whole otolith, as well as for the sectioned version of the otolith. If the age readings were identical for each reader, then that age was considered the correct age. Otherwise, the two readers discussed the readings and agreed upon an age. The nonparametric, paired Wilcoxon test was applied to the pairs of whole and sectioned otolith ages for each fish. In addition, a paired Wilcoxon test was applied to otoliths for fish restricted to 8 years and older ($n$=32).

In addition to determining whether to use whole or sectioned otoliths, the preliminary analysis also involved determining how halibut age was to be estimated: either by using the mutually determined age, as already described, or an averaged age. The averaged age was a single value calculated as the mean of the four independent readings (whole or sectioned version of an otolith read by two readers on two different occasions) and rounded to the nearest integer. Sample means generally have the same type of sampling distribution as individual observations, but with a smaller variance. Hence, we applied the nonparametric paired Wilcoxon test to the pairs of averaged and mutually determined ages for each otolith ($n$=80). The results from the preliminary analysis determined how we conducted the process of age determination in the main part of the study.



**Figure 1**

An example of an otolith viewed whole. Age was determined to be 3 years.

We acknowledge that using sectioned otoliths is the recommended procedure when analyzing long-lived fish. Although individual California halibut may live as long as 30 years (Frey, 1971; Pattison and McAllister, 1990), all the fish in our study were relatively young (less than 14 years old). Because of our preliminary analysis and because sectioning a large number of otoliths (potentially over 1300) would have entailed considerable costs, all otoliths were initially read whole. We did use sectioned otoliths when the variability in the age readings for that otolith seemed to be unacceptably high and when the whole otolith was judged unreadable because of the anise oil residue.

Before recording ages, two readers consulted with each other to standardize their technique for counting annuli. Each otolith was initially read whole, that is, the surface of the uncut otolith was read for the number of annuli present. The whole otolith was submerged in water, illuminated with fiber optic light transmitted from the sides, and viewed against a dark background with a dissecting microscope (16× to 25×). Figure 1 is an example of an otolith

**Figure 2**

An example of a whole otolith that was unreadable (**A**) according to our criteria, but readable sectioned (**B**). Age was determined to be 9 years.

viewed whole. Some whole otoliths from central California were soaked in anise oil during initial handling to facilitate viewing annuli. The anise oil, however, left a residue that later caused problems in reading the otoliths. Therefore, these otoliths had to be sectioned before they could be read. For each otolith (either whole or sectioned), the two readers counted the annuli twice at different times, for a total of four independent readings. The birth date for a halibut was assigned to 1 January (Williams and Bedford, 1974).

The mean and standard deviation of the four age determinations for each whole otolith were calculated. Because of our preliminary analysis, if the standard deviation was less than 1.5 years, the mean, rounded to the nearest integer, was the estimated age for that halibut. A standard deviation of 1.5 years was arbitrarily chosen as the upper limit for acceptance of an age determination. This deviation represented a difference in ages of 3 years or more among readings. If the standard deviation was 1.5 years or larger the whole otolith was judged unreadable. These otoliths were cut laterally through the nucleus with a diamond blade on a Buehler low-speed Isomet saw. Three sections, approximately 0.38 mm thick, were cut from each otolith and mounted on a glass slide with Eukitt clear mounting medium. These sections were viewed with a compound microscope (25× to 100×) with transmitted light. The best of three sections on a slide was chosen for age determination. Figure 2 is an example of a whole otolith that was unreadable, but readable when sectioned.

The mean and standard deviation of the four age determinations for each sectioned otolith were calculated. In accordance with our preliminary analysis, if the standard deviation was less than 1.5 years, the rounded mean was considered the estimated age. If the standard deviation was 1.5 years or larger, the otolith was not used in the analysis because it was felt to be unreadable. The index of average percent error was calculated to compare within-reader precision (Beamish and Fournier, 1981) for both whole and sectioned otoliths.

Several analyses were made on total lengths and ages by sex and region. Comparisons of both length and age distributions between sexes for each region were made by using the two-sample Kolmogorov-Smirnov test (Hollander and Wolfe, 1973). Length-at-age data for females were compared with those for males sampled from the same region by using the nonparametric Mann-Whitney test. The Mann-Whitney test was also used to compare the length-at-age data between regions for each sex separately.

The von Bertalanffy growth function was fitted to the length-at-age data for individual fish (Ricker, 1975); parameters were estimated by nonlinear least squares by us-

ing the Gauss-Newton method in the program NLIN from SAS (1990). Outliers were removed when the von Bertalanffy growth function was fitted to the data. Outliers were data points that clearly stood apart in scatter plots of individual total length versus age. Means and standard deviations of observed length-at-age, and the theoretical von Bertalanffy growth function (VBGF) were plotted for females and males by region.

Hotelling's $T^2$ test (Bernard, 1981) compared growth parameters between male and female halibut in each region. This test was also used to determine if growth in southern California was different from growth in central California, for each sex separately.

The estimated annual survival rate, $S$, was calculated for males and females separately by region by using Robson and Chapman's method (1961). Total annual mortality, $A$, was calculated by using the complement of annual survival, $A = 1 - S$. The instantaneous mortality rate, $Z$, was estimated by using $-\ln(S)$ (Ricker, 1975). All statistical analyses were performed at the 0.05 level of significance.

## Results

In our preliminary analysis, a subsample of whole otoliths from 80 fish were read and then sectioned to determine the best method for reading otoliths for age determination. In Figure 3 is a plot of the mutually determined ages from sectioned and whole otoliths, along with a 45 degree line and a linear regression. No significant difference was found between ages from whole and sectioned otoliths when using all 80 fish (paired Wilcoxon test, $P$=0.21), and when using only older fish (8 years and older) (paired Wilcoxon test, $P$=0.11).

Also, as part of the preliminary analysis, in the comparison of averaged ages with mutually determined ages for the estimation of halibut age, we found no significant differences for both whole (paired Wilcoxon test, mean difference=0.15, $P$=0.23) and sectioned ototliths (paired Wilcoxon test, mean difference=0.16, $P$=0.06). Therefore, ages were estimated for each individual by calculating an average of four independent readings from whole or sectioned otoliths because an average of several readings is considered a more reliable measurement than a single reading (Fleiss, 1986).

The total number of trawls conducted by strata and region are listed in Table 1. The original intent was to perform 50 trawls at each depth stratum for both southern and central California. However, substrate obstructions and other problems limited the number of successful tows at most strata. A total of 58 trawls were successfully completed in stratum 1 in southern California.

Because most of the halibut were collected in a single stratum (stratum 1), comparisons among strata were not made (Table 1). In southern California, a total of 916 individuals were collected from stratum 1 (84% of total), 167 individuals from stratum 2, and only 14 individuals from stratum 3. In central California, a total of 254 individuals were collected from stratum 1 (92% of total), 22 individuals from stratum 2, and none from stratum 3.




**Figure 3**

A scatter plot of mutually determined ages for sectioned otoliths and whole otoliths. The dotted line is the linear regression through the origin (slope is 0.947); solid line is the 45 degree line.

For southern California, pairs of whole otoliths from 1109 individuals were initially examined to estimate age. Twenty-two pairs of whole otoliths were unreadable, six pairs had no discernible opaque and translucent zones, and sixteen pairs had large variations in the four age readings (SD ≥1.5). These 22 pairs of whole otoliths were sectioned to determine age. Of these, two were eliminated because of large variations in the four age readings (SD ≥1.5). An additional ten were eliminated as outliers because lengths were anomalously small or large for the estimated age when the von Bertalanffy growth function was fitted to the data. Thus, a total of 1097 individuals were used to evaluate age and growth for halibut in southern California. The sample consisted of 69% males (761 individuals) and 31% females (336 individuals). Tables 2 and 3 provide counts of total length versus age for males and females, respectively, in southern California.

In central California, pairs of whole otoliths from 292 individuals were initially examined to estimate age. Otoliths from 28 halibut had no discernible opaque and translucent zones and 49 had large variations in the four age readings (SD ≥1.5), resulting in a total of 77 otoliths that were sectioned to estimate age. Of these, fourteen otolith pairs were eliminated because of large variations in readings (SD ≥1.5). An additional two became outliers that were removed when the von Bertalanffy growth function was fitted to the data. Thus, a total of 276 individuals were used to evaluate age and growth for halibut in central California. The sample consisted of 53% (147 individuals) males and 47% (129 individuals) females. Counts of total length versus age for males and females, respectively, in central California are shown in Tables 4 and 5.

Within-reader index of average percent error for whole otoliths ranged from 8.1% to 13.2%. For sectioned otoliths,

**Table 2**

Male California halibut total length versus age (mostly whole otoliths, some sectioned) in southern California.

| | Age (yr) | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| TL (mm) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | Mean age |
| 201–225 | | 2 | 1 | | | | | | | | | | 2.3 |
| 226–250 | 2 | 5 | 2 | | | | | | | | | | 2.0 |
| 251–275 | 1 | 20 | 14 | | | | | | | | | | 2.4 |
| 276–300 | 1 | 51 | 59 | 4 | 1 | | | | | | | | 2.6 |
| 301–325 | | 14 | 33 | 4 | 2 | | | | | | | | 2.9 |
| 326–350 | | 7 | 21 | 18 | 5 | | | | | | | | 3.4 |
| 351–375 | | 4 | 32 | 34 | 17 | 3 | | | | | | | 3.8 |
| 376–400 | | | 11 | 25 | 16 | 6 | 1 | | | | | | 4.3 |
| 401–425 | | | 5 | 10 | 25 | 8 | 12 | 2 | | | | | 5.3 |
| 426–450 | | | | 8 | 15 | 15 | 11 | 2 | | | | | 5.7 |
| 451–500 | | | | 6 | 14 | 26 | 21 | 6 | 6 | | | | 6.3 |
| 501–525 | | | | 1 | 7 | 14 | 17 | 5 | 2 | | | | 6.5 |
| 526–550 | | | | | 1 | 5 | 9 | 8 | 4 | 2 | | | 7.5 |
| 551–575 | | | | | 2 | 9 | 9 | 9 | 3 | 1 | | | 7.2 |
| 576–600 | | | | | 1 | 4 | 7 | 1 | 3 | 0 | 1 | 1 | 7.6 |
| 601–625 | | | | | | 1 | 6 | 5 | 2 | 3 | | | 8.0 |
| 626–650 | | | | | | 1 | 4 | 1 | 2 | | | | 7.5 |
| 651–675 | | | | | | | | | | | | | 0.0 |
| 676–700 | | | | | | | | | | | | | 0.0 |
| 701–725 | | | | | | | | 1 | 1 | | | | 8.5 |
| Mean length | 259 | 289 | 319 | 377 | 417 | 482 | 506 | 538 | 552 | 578 | 577 | 586 | |

Median length of distribution: 387 mm TL

within-reader average percent error ranged from 4.7% to 8.9%.

In southern California, significant differences between males and females were found in comparisons of length distributions (Kolmogorov-Smirnov: $D$=0.42, $P$=0.0001). A higher proportion of males were smaller than females (Fig. 4). The median length for males was 387 mm TL (range: 210–707 mm TL); the median length for females was 544 mm TL (range: 241–987 mm TL) (Tables 2, 3, and 6). Females on average were significantly larger than males for ages 3 through 10 (Mann-Whitney tests, Table 6).

Some differences in lengths were also found between males and females in central California. Most of the larger fish were females (Fig. 4). However, the length distributions of males and females in central California were not significantly different (Kolmogorov-Smirnov: $D$=0.14, $P$=0.12). The median length for males was 437 mm TL (range: 285–781 mm TL); the median length for females was 444 mm TL (range: 262–1039 mm TL) (Tables 4, 5, and 7). Females on average were significantly larger than males for ages 3 and 5 through 8 (Mann-Whitney tests, Table 7).

In addition, comparisons were made between regions for each sex separately. In comparisons of mean length at ages, the males in central California on average were significantly larger than those in southern California for ages 2 through 9, except age 8 (Mann-Whitney tests, Table 8). The females in central California on average were significantly larger than those in southern California for only ages 5, 7, and 8 (Mann-Whitney tests, Table 8). However, the median length of females in southern California (544 mm) was greater than the median length for those in central California (444 mm).

In southern California, age distributions were significantly different for males and females (Kolmogorov-Smirnov: $D$=0.18, $P$=0.0001). A higher proportion of females were older fish compared with males. The primary age mode for females was age 6 and for males it was age 3 (Fig. 5). In central California, age distributions were also different for males and females (Kolmogorov: $D$=0.20, $P$=0.008). Males had a higher percentage in the 6 to 8 year range and a lower percentage in the 3 to 5 year range compared with females (Fig. 5). In central California, the primary mode for both males and females was age 3. With both regions combined, females were found up to age 13 and males up to age 12. In comparisons of age distributions between regions, we found that a higher proportion of halibut in southern California was older than halibut in central California (Fig. 5).

Because differences in length-at-age were found between sexes, von Bertalanffy growth parameters were cal-

**Table 3**

Female California halibut total length versus age (mostly whole otoliths, some sectioned) in southern California.

| | Age (yr) | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| TL (mm) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | Mean age |
| 201–225 | | | | | | | | | | | | | | 0.0 |
| 226–250 | | 3 | | | | | | | | | | | | 2.0 |
| 251–275 | 1 | 5 | 4 | | | | | | | | | | | 2.3 |
| 276–300 | | 7 | 7 | | | | | | | | | | | 2.5 |
| 301–325 | | 1 | 2 | | | | | | | | | | | 2.7 |
| 326–350 | | 3 | 6 | 5 | | | | | | | | | | 3.1 |
| 351–375 | | 2 | 4 | 2 | | | | | | | | | | 3.0 |
| 376–400 | | | 4 | 5 | 2 | | | | | | | | | 3.8 |
| 401–425 | | | 6 | 12 | 3 | 1 | | | | | | | | 4.0 |
| 426–450 | | | 5 | 5 | 3 | 1 | | | | | | | | 4.0 |
| 451–500 | | | 4 | 9 | 10 | 4 | 3 | 1 | | | | | | 4.9 |
| 501–525 | | | | 1 | 17 | 5 | 1 | | | | | | | 5.3 |
| 526–550 | | | | 3 | 9 | 8 | 1 | 1 | 1 | | | | | 5.6 |
| 551–575 | | | | 4 | 3 | 10 | 6 | 2 | | | | | | 6.0 |
| 576–600 | | | | 4 | 3 | 8 | 4 | 4 | 1 | | | | | 6.2 |
| 601–625 | | | | 2 | 4 | 8 | 6 | 2 | | | | | | 6.1 |
| 626–650 | | | | | 3 | 10 | 9 | 4 | 1 | | | | | 6.6 |
| 651–675 | | | | 2 | 1 | 10 | 6 | 3 | 2 | | | | | 6.5 |
| 676–700 | | | | | | 3 | 1 | 1 | 0 | 0 | 1 | | | 7.3 |
| 701–725 | | | | | | | 3 | 1 | 1 | 1 | | | | 8.0 |
| 726–750 | | | | | | 2 | 0 | 2 | | | | | | 7.0 |
| 751–775 | | | | | | | 1 | | | | | | | 7.0 |
| 776–800 | | | | | | | | 1 | 1 | 3 | | | | 9.4 |
| 801–825 | | | | | | | | 1 | 1 | 1 | | | | 9.0 |
| 826–850 | | | | | | | | 1 | 1 | 1 | | | | 9.0 |
| 851–875 | | | | | | | | 2 | 0 | 1 | | | | 8.7 |
| 876–900 | | | | | | | | | | | | | | 0.0 |
| 901–925 | | | | | | | | | 1 | 1 | | | | 9.5 |
| 926–950 | | | | | | | | | 1 | 1 | | | | 9.5 |
| 951–975 | | | | | | | | | | | | | 1 | 13.0 |
| 976–1000 | | | | | | | | | | | 1 | | | 11.0 |
| Mean length | 252 | 292 | 364 | 463 | 520 | 591 | 616 | 666 | 736 | 828 | 836 | 000 | 964 | |

Median length of distribution: 544 mm TL

culated separately for males and females for both regions (Table 9). Outliers were removed when the von Bertalanffy growth function was fitted to the data. With outliers removed, the standard errors of the asymptotic mean length, $L_\infty$, became considerably smaller. Graphs of the von Bertalanffy growth function are shown in Figure 6 (where females and males are compared by region) and Figure 7 (where southern and central California are compared by sex).

Significant differences between males and females in southern California were found for all three growth parameters (Hotelling's $T^2$ tests, $P<0.0001$). For central California, two of the three growth parameters were significantly different between males and females (Hotelling's $T^2$ tests, $P<0.0001$). Females grew faster and on average were larger than males at the same age in both regions (Fig. 6).

Similar comparisons were made between regions for each sex separately (Fig. 7). For females, all three growth parameters were significantly different between southern and central California (Hotelling's $T^2$ tests, $P\leq0.003$). In contrast, the only significant difference between regions for males was in $K$, the Brody growth coefficient (Hotelling's $T^2$ tests, $P<0.0001$).

In southern California, survival rates were based on ages 7 to 12 years for males and ages 6 to 11 years for females. The estimated annual survival rates, $S$, for halibut from south-

**Table 4**

Male California halibut total length versus age (mostly whole otoliths, some sectioned) in central California.

| | Age (yr) | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| TL (mm) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | Mean age |
| 201–225 | | | | | | | | | | | | | 0.0 |
| 226–250 | | | | | | | | | | | | | 0.0 |
| 251–275 | | | | | | | | | | | | | 0.0 |
| 276–300 | | 4 | 1 | | | | | | | | | | 2.2 |
| 301–325 | | 4 | 3 | | | | | | | | | | 2.4 |
| 326–350 | | 5 | 4 | | | | | | | | | | 2.4 |
| 351–375 | | | 9 | | | | | | | | | | 3.0 |
| 376–400 | | 2 | 15 | 8 | | | | | | | | | 3.2 |
| 401–425 | | | 9 | 1 | 1 | | | | | | | | 3.3 |
| 426–450 | | | 2 | 7 | 2 | | | | | | | | 4.0 |
| 451–500 | | | 3 | 6 | 2 | 4 | 1 | 1 | | | | | 4.8 |
| 501–525 | | | | 2 | 1 | 5 | 1 | 3 | | | | | 6.2 |
| 526–550 | | | | 1 | 1 | 7 | 1 | 1 | | | | | 6.0 |
| 551–575 | | | | | 1 | 2 | 4 | 1 | | | | | 6.6 |
| 576–600 | | | | | 1 | 0 | 0 | 2 | | | | | 7.0 |
| 601–625 | | | | | | | 1 | 1 | 1 | | | | 8.0 |
| 626–650 | | | | | | 1 | 2 | 2 | 1 | | | | 7.5 |
| 651–675 | | | | 1 | 0 | 0 | 2 | | | | | | 6.0 |
| 676–700 | | | | | | 1 | 0 | 1 | 1 | | | | 7.7 |
| 701–725 | | | | | | | | 1 | | | | | 8.0 |
| 726–750 | | | | | | | 1 | 0 | 0 | 0 | 1 | | 9.0 |
| 776–800 | | | | | | | | | | 1 | | | 10.0 |
| Mean length | 000 | 326 | 385 | 448 | 493 | 539 | 596 | 582 | 649 | 781 | 737 | 000 | |

Median length of distribution: 437 mm TL




**Figure 4**

Length frequencies for male and female California halibut sampled off southern California (So. CA) and central California (Central CA).

**Table 5**

Female California halibut total length versus age (mostly whole otoliths, some sectioned) in central California.

| | Age (yr) | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| TL (mm) | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | Mean age |
| 201–225 | | | | | | | | | | | | | | 0.0 |
| 226–250 | | | | | | | | | | | | | | 0.0 |
| 251–275 | | 1 | | | | | | | | | | | | 2.0 |
| 276–300 | | 3 | 4 | | | | | | | | | | | 2.6 |
| 301–325 | | 5 | 7 | | | | | | | | | | | 2.6 |
| 326–350 | | 2 | 10 | 2 | | | | | | | | | | 3.0 |
| 351–375 | | 1 | 7 | | | | | | | | | | | 2.9 |
| 376–400 | | | 8 | 5 | | | | | | | | | | 3.4 |
| 401–425 | | | 3 | 2 | | | | | | | | | | 3.4 |
| 426–450 | | | 2 | 6 | | | | | | | | | | 3.8 |
| 451–500 | | | 1 | 3 | 2 | | | | | | | | | 4.2 |
| 501–525 | | | | 7 | 2 | | | | | | | | | 4.2 |
| 526–550 | | | 2 | 1 | 3 | | | | | | | | | 4.2 |
| 551–575 | | | 1 | 3 | 1 | | | | | | | | | 4.0 |
| 576–600 | | | 1 | 2 | 2 | 1 | | | | | | | | 4.5 |
| 601–625 | | | | 1 | | | | | | | | | | 4.0 |
| 626–650 | | | | | 1 | 1 | 0 | 0 | 1 | | | | | 6.7 |
| 651–675 | | | | 1 | 1 | | | | | | | | | 4.5 |
| 676–700 | | | | 1 | 0 | 1 | 1 | 1 | | | | | | 6.3 |
| 701–725 | | | | | 1 | 0 | 0 | 1 | | | | | | 6.5 |
| 726–750 | | | | | 3 | | | | | | | | | 5.0 |
| 751–775 | | | | | 1 | 0 | 1 | | | | | | | 6.0 |
| 776–800 | | | | | | 1 | | | | | | | | 6.0 |
| 801–825 | | | | 1 | | | | | | | | | | 4.0 |
| 826–850 | | | | | | | | 1 | | | | | | 8.0 |
| 851–875 | | | | | | | | | 1 | | | | | 9.0 |
| 876–900 | | | | | | | | 1 | 1 | | | | | 8.5 |
| 901–925 | | | | | | | | 1 | 2 | | | | | 8.7 |
| 926–950 | | | | | | | | | | | | | | 0.0 |
| 951–975 | | | | | | | | | | 1 | | | | 10.0 |
| 976–1000 | | | | | | | | | | | | | | 0.0 |
| 1001–1025 | | | | | | | | | | | | 1 | | 12.0 |
| 1026–1050 | | | | | | | | | | | | 1 | | 12.0 |
| Mean length | 000 | 313 | 373 | 493 | 606 | 672 | 726 | 810 | 842 | 970 | 000 | 1027 | 000 | |

Median length of distribution: 444 mm TL

ern California were 0.40 (95% confidence interval=0.34, 0.46) for males and 0.51 (95% confidence interval=0.45, 0.57) for females. Estimated annual mortality rates, $A$, were 0.60 for males and 0.49 for females. Instantaneous mortalities, $Z$, were estimated at 0.91 for males and 0.68 for females. We were unable to calculate survival rates and mortalities for central California halibut because the sample sizes were too small.

## Discussion

The California halibut can be a long-lived species, living as long as 30 years (Frey, 1971; Pattison and McAllister, 1990). However, none of the fish in our study was older than 13 years, and most of the fish were less than 11 years old (Fig. 5). We feel that the low average percent error in our study showed that both whole and sectioned otoliths were reliable methods for aging California halibut for the age range of our study.

Pattison and McAllister (1990) suggested that because of the asymmetrical growth of the sagittae in larger fish, ages assigned to older fish (age 11 or older) may be underestimated if whole otoliths, rather than sectioned otoliths, were read. Manooch and Potts (1997) in a growth study of greater amberjack also did not find whole otoliths useful in aging fish. In contrast, we found that age estimates from whole and sectioned otoliths were not sig-

nificantly different, even when the analysis was limited to ages 8 to 13 years. However, our conclusions are limited to primarily younger fish because most of the fish in our study were less than 11 years old. Pattison and McAllister recommended sectioned otoliths for fish aged 11 years or older. Nonetheless, we concur that use of sectioned otoliths in older individuals may reduce the variation in assigned ages.

We feel that the larger percentage of unreadable whole otoliths in central California versus southern California was due primarily to the preparation techniques used for age determination. The anise oil used in the initial handling of some central California otoliths created a white film that precluded the otoliths from being read at a later time. Because of problems encountered with anise oil, we do not recommend applying it to otoliths until one is ready to age them.

**Table 6**

Mean total length (mm) and standard deviation for male and female California halibut by age group in southern California. Mann-Whitney comparisons of male and female halibut for mean length at age. Asterisk (*) indicates statistical significance.

| Age (yr) | Male | | | | | Female | | | | | Mann-Whitney *P*-value |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | *n* | Mean | SD | Min | Max | *n* | Mean | SD | Min | Max | |
| 1 | 4 | 259.3 | 25.0 | 232 | 290 | 1 | 252.0 | — | 252 | 252 | — |
| 2 | 103 | 289.4 | 28.4 | 210 | 369 | 21 | 291.9 | 38.4 | 241 | 368 | 0.77 |
| 3 | 178 | 319.3 | 39.3 | 220 | 411 | 42 | 364.4 | 63.9 | 254 | 477 | 0.0001* |
| 4 | 110 | 377.3 | 42.2 | 291 | 501 | 54 | 462.6 | 86.7 | 342 | 660 | 0.0001* |
| 5 | 106 | 417.2 | 55.0 | 294 | 561 | 58 | 520.5 | 64.3 | 389 | 671 | 0.0001* |
| 6 | 92 | 481.7 | 61.9 | 362 | 646 | 70 | 591.2 | 67.7 | 401 | 735 | 0.0001* |
| 7 | 97 | 505.6 | 65.1 | 399 | 645 | 41 | 616.1 | 60.5 | 493 | 758 | 0.0001* |
| 8 | 40 | 538.3 | 62.6 | 403 | 707 | 26 | 666.0 | 101.4 | 492 | 859 | 0.0001* |
| 9 | 23 | 551.5 | 68.0 | 451 | 707 | 11 | 736.2 | 131.9 | 534 | 934 | 0.0002* |
| 10 | 6 | 578.2 | 34.5 | 542 | 619 | 9 | 828.3 | 67.4 | 720 | 937 | 0.002* |
| 11 | 1 | 577.0 | — | 577 | 577 | 2 | 836.0 | 213.5 | 685 | 987 | — |
| 12 | 1 | 586.0 | — | 586 | 586 | — | — | — | — | — | — |
| 13 | — | — | — | — | — | 1 | 964.0 | — | 964 | 964 | — |
| Total | 761 | | | | | 336 | | | | | |

**Table 7**

Mean total length (mm) and standard deviation for male and female California halibut by age-group in central California. Mann-Whitney comparisons of male and female halibut for mean length at age. Asterisk (*) indicates statistical significance.

| Age (yr) | Male | | | | | Female | | | | | Mann-Whitney *P*-value |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | *n* | Mean | SD | Min | Max | *n* | Mean | SD | Min | Max | |
| 2 | 15 | 325.9 | 29.8 | 285 | 381 | 12 | 312.9 | 28.7 | 262 | 367 | 0.29 |
| 3 | 46 | 384.8 | 38.1 | 293 | 472 | 46 | 373.4 | 70.0 | 279 | 593 | 0.02* |
| 4 | 26 | 447.8 | 60.5 | 380 | 660 | 35 | 493.5 | 102.3 | 328 | 820 | 0.07 |
| 5 | 9 | 492.8 | 59.0 | 424 | 582 | 17 | 606.2 | 101.4 | 474 | 758 | 0.007* |
| 6 | 20 | 538.6 | 49.4 | 479 | 680 | 4 | 672.0 | 87.0 | 582 | 782 | 0.006* |
| 7 | 13 | 595.6 | 72.7 | 490 | 748 | 2 | 725.5 | 43.1 | 695 | 756 | 0.05* |
| 8 | 13 | 582.5 | 74.4 | 475 | 719 | 5 | 810.4 | 99.3 | 694 | 925 | 0.002* |
| 9 | 3 | 648.7 | 29.5 | 619 | 678 | 5 | 842.0 | 112.2 | 547 | 923 | — |
| 10 | 1 | 781.0 | — | 781 | 781 | 1 | 970.0 | — | 970 | 970 | — |
| 11 | 1 | 737.0 | — | 737 | 737 | — | — | — | — | — | — |
| 12 | — | — | — | — | — | 2 | 1026.5 | 17.7 | 1014 | 1039 | — |
| Total | 147 | | | | | 129 | | | | | |

The data in our study confirmed that females grow faster than males, as found in previous studies (Pattison and McAllister, 1990; Sunada et al., 1990), and that growth was different between males and females for each region. Females in both regions had larger asymptotic mean lengths, $L_\infty$, than males (Table 9).

Growth curves for males and females crossed between ages 1 and 2 for southern California and at age 3 for central California (Fig. 6). This finding suggests that young male and female halibut are of similar size until about ages 2 and 3 in southern and central California, respectively. However, Haaker (1975) found that juvenile female halibut in Anaheim Bay grew faster than males. The crossover in growth curves may be due to the small sample sizes for ages 1 and 2 in our study. The growth curves also involved some extrapolation for very young ages.

Our study showed lower mean length-at-age than that obtained by Pattison and McAllister (1990); however, the differences for females were slight. These differences may partly be attributed to different environmental conditions in the years prior to each study. Pattison and McAllister used otoliths collected from 1955–66 and 1984–88, whereas we used otoliths collected from 1993 to 1995. Manooch and Potts (1997) in a study of greater amberjack in the Gulf of Mexico also cited temporal changes as one of the factors that may have contributed to differences in growth between greater amberjack in southern Florida and northern Gulf of Mexico. The geographic range of the samples may also have affected the results because we found growth differences between southern and central California. Pattison and McAllister's study encompassed samples from central and southern California, but they did not separate the two regions in their analysis. In addition, gear selectivity may have contributed to the differences found between our two studies. Pattison and McAllister used halibut that were collected with several gear types (trawl, gill net, beach seine, hook and line, and spear), whereas our study sampled halibut with only trawl gear of a specific mesh size. Potts et al. (1998) found while studying vermillion snapper from the southeastern United States that the different gear types used in sample collection may bias growth results.

Sunada et al. (1990) found that for commercial halibut landings in southern California the asymptotic mean length, $L_\infty$, was 909 mm TL for males and 1445 mm TL for females. In comparison, our study showed for southern California a slightly higher $L_\infty$ for males at 925 mm TL, and a lower $L_\infty$ for females at 1368 mm TL. The asymptotic mean length for females may be lower because fewer large females were collected in our study. Although our study had different growth parameters than those of Sunada et al. (1990), the differences were only slight. The differences may have been due to gear vulnerability. Sunada et al. (1990) collected halibut from both trawl nets and gill nets and found that fewer females were collected in trawl nets, which may be attributed to trawl gear being restricted to offshore areas during spawning season. Also

**Table 8**

Mann-Whitney comparisons of California halibut from southern and central California for mean length at age. Asterisk (*) indicates statistical significance.

| Age (yr) | Mean observed total length (mm) | | Sample size $n$ | | Mann-Whitney $P$-value |
|---|---|---|---|---|---|
| | Southern | Central | Southern | Central | |
| Males | | | | | |
| 2 | 289.4 | 325.9 | 103 | 15 | 0.0001* |
| 3 | 319.3 | 384.8 | 178 | 46 | 0.0001* |
| 4 | 377.3 | 447.8 | 110 | 26 | 0.0001* |
| 5 | 417.2 | 492.8 | 106 | 9 | 0.0006* |
| 6 | 481.7 | 538.6 | 92 | 20 | 0.0002* |
| 7 | 505.6 | 595.6 | 97 | 13 | 0.0002* |
| 8 | 538.3 | 582.5 | 40 | 13 | 0.08 |
| 9 | 551.5 | 648.7 | 23 | 3 | 0.03* |
| Females | | | | | |
| 2 | 291.9 | 312.9 | 21 | 12 | 0.10 |
| 3 | 364.4 | 373.4 | 42 | 46 | 0.92 |
| 4 | 462.6 | 493.5 | 54 | 35 | 0.21 |
| 5 | 520.5 | 606.2 | 58 | 17 | 0.002* |
| 6 | 591.2 | 672.0 | 70 | 4 | 0.07 |
| 7 | 616.1 | 725.5 | 41 | 2 | 0.04* |
| 8 | 666.0 | 810.4 | 26 | 5 | 0.01* |
| 9 | 736.2 | 842.0 | 11 | 5 | 0.17 |

**Table 9**

Von Bertalanffy growth parameters with asymptotic standard errors (SE) for California halibut by sex in southern and central California. Length is given in mm TL.

| Region | Sex | $n$ | $L_\infty$ | SE | $K$ | SE | $t_0$ | SE |
|---|---|---|---|---|---|---|---|---|
| Southern | male | 761 | 925.3 | 121.4 | 0.08 | 0.02 | −2.2 | 0.41 |
| | female | 336 | 1367.7 | 273.4 | 0.08 | 0.03 | −1.2 | 0.48 |
| Central | male | 147 | 956.7 | 211.9 | 0.10 | 0.05 | −2.1 | 0.91 |
| | female | 129 | 1477.1 | 308.1 | 0.10 | 0.04 | −0.2 | 0.43 |




**Figure 5**

Age frequencies in percent for male and female California halibut sampled off southern and central California (CA).




**Figure 6**

Von Bertalanffy growth curves for comparing growth between sexes by region.




**Figure 7**

Von Bertalanffy growth curves for comparing growth between regions by sex.

females, which are generally larger, may be more capable of escaping a trawl net than males. Other factors that may have affected the growth parameters include the following: method of collection, time periods of sampling, and different environmental conditions in the years prior to each study.

Many factors can affect growth rates of fish including differences in the seasonality of spawning, environmental factors, amount and size of food, and genetics (Weatherly and Gill, 1987; Moyle and Cech, 1988). Southern and central California are biogeographically different. Differences between central and southern California coastal waters include temperature, water circulation patterns, bottom topography, and substrate. In particular, Jow (1990) noted differences in bottom topography and substrate between the trawl areas for California halibut in southern and northern (San Francisco Bay area) California. The biogeographical differences among these regions could possibly cause growth rates of the same species to differ. Studies of other fish species have shown differences in growth between geographical regions. For example, Parrish et al. (1985) found latitudinal differences in the growth of northern anchovy, *Engraulis mordax*. They found that the growth rate of juvenile northern anchovy in central California was greater than that of juvenile northern anchovy in southern California. Butler et al. (1996) found differences in maturation and length at age of Pacific sardine between latitudinal regions. Sardine appeared to mature at a younger age off both southern and Baja California than off Monterey; one-, two-, and three-year old fish were smaller at age off Baja and larger off Monterey. Deriso et al. (1996) confirmed the findings of sardine growth differences by Butler et al. (1996). The exact mechanism for latitudinal differences in growth rates of California halibut is still unclear and further research is needed to clarify this phenomenon.

## Acknowledgments

We wish to express our appreciation to Larry Jacobson with the National Marine Fisheries Service, for his detailed review of an earlier draft of the paper. We appreciate the efforts of Steve Wertz, Sandra Owen, and Paul Gregory who helped with various aspects of this study. We also thank the reviewers for their comments and suggestions that improved the paper.

## Literature cited

Beamish, R. J., and D. A. Fournier.
1981. A method for comparing the precision of a set of age determinations. Can. J. Fish. Aquat. Sci. 38: 982–983.

Bernard, D. R.
1981. Multivariate analysis as a means of comparing growth in fish. Can. J. Fish. Aquat. Sci. 38:233–236.

Butler, J. L., M. L. Granados G., J. T. Barnes, M. Yaremko, and B. J. Macewicz..
1996. Age composition, growth, and maturation of the Pacific sardine (*Sardinops sagax*) during 1994. Calif. Coop. Oceanic Fish Invest. Rep. 37:152–159.

Casselman, J. M.
1983. Age and growth assessment of fish from their calcified structures-techniques and tools. *In* Proceedings of the international workshop on age determination of oceanic pelagic fishes: tunas, billfishes, sharks (E. Prince and L. Pulos, eds.) p. 1–17. U.S. Dep. Commer, NOAA Tech. Report 8.

Deriso, R. B., J. T. Barnes, L. D. Jacobson, and P. R. Arenas.
1996. Catch-at-age analysis for Pacific sardine (*Sardinops sagax*) 1983–1995. Calif. Coop. Oceanic Fish Invest. Rep. 37:175–187.

Fleiss, J. L.
1986. The design and analysis of clinical experiments. John Wiley and Sons, New York, NY, 432 p.

Frey, H. W., ed.
1971. California's living marine resources and their utilization. Calif. Dep. Fish Game, Sacramento, CA, 148 p.

Haaker, P. L.
1975. The biology of the California halibut, *Paralichthys californicus* (Ayres), in Anaheim Bay, California. *In* The marine resources of Anaheim Bay (E. D. Lane and C. W. Hill, eds), p. 137–151. Calif. Dep. Fish Game, Fish Bull. 165.

Hollander, M., and D. A. Wolfe.
1973. Nonparametric statistical methods. John Wiley and Sons, New York, NY, 503 p.

Jow, T.
1990. The California halibut trawl fishery. *In* The California halibut, *Paralichthys californicus*, resources and fisheries (C. W. Haugen, ed.), p. 229–241. Calif. Dep. Fish Game, Fish Bull. 174.

Kramer, S. H., and J. S. Sunada.
1992. California halibut. *In* California's living marine resources and their utilization (W. S. Leet, C. M. Dewess, and C. W. Haugen, eds.), p. 94–97. California Sea Grant, Univ. Calif., Davis, CA.

Manooch, C. S. III, and J. C. Potts.
1997. Age, growth, and mortality of greater amberjack, *Seriola dumerili*, from the U.S. Gulf of Mexico headboat fishery. Bull. Mar. Sci. 61:671–683.

Moyle, P. B., and J. J. Cech Jr.
1988. Fishes: an introduction to ichthyology, 2nd ed. Prentice-Hall, Inc., Englewood Cliffs, NJ, p. 593

Parrish, R. H., D. L. Mallicoate, and K. F. Mais.
1985. Regional variations in the growth and age composition of northern anchovy, *Engraulis mordax*. Fish. Bull. 83:483–496.

Pattison, C. A. and R. D. McAllister.
1990. Age determination of California halibut, *Paralichthys californicus*. *In* The California halibut, *Paralichthys californicus*, resources and fisheries (C. W. Haugen, ed.), p. 207–216. Calif. Dep. Fish Game, Fish Bull. 174.

Potts, J. C., C. S. Manooch III, and D. S. Vaughan.
1998. Age and growth of vermillion snapper from the southeastern United States. Trans. Am. Fish. Soc. 127:787–795.

Ricker, W. E.
1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191, 382 p.

Robson, D. S., and D. G. Chapman.
1961. Catch curves and mortality rates. Trans. Am. Fish. Soc. 90:181–189.

SAS.
1990. SAS/STAT user's guide, version 6, fourth ed., vol. 2. SAS Institute, Inc., Cary, NC, 846 p.

Sunada, J. S., P. V. Velez, and C. A. Pattison.
1990. Age, size, and sex composition of California halibut from southern California commercial fishery landings, 1983–1988. *In* The California halibut, *Paralichthys californicus*, resources and fisheries (C.W. Haugen, ed.), p. 303–319. Calif. Dep. Fish Game, Fish Bull. 174.

Weatherly, A. H., and H. S. Gill.
1987. The biology of fish growth. Academic Press, Orlando, FL, 443 p.

Williams, T., and B. C. Bedford.
1974. The use of otoliths for age determination. *In* Aging of fish: proceedings of an international symposium, p. 114–123. Unwin Brothers Ltd., Surrey, England.

# Morphological development and growth of laboratory-reared larval and juvenile *Thunnus thynnus* (Pisces: Scombridae)

**Shigeru Miyashita**
**Yoshifumi Sawada**
**Tokihiko Okada**
**Osamu Murata**
**Hidemi Kumai**

Fisheries Laboratory of Kinki University,
3153, Shirahama,
Wakayama 649-2211, Japan
E-mail address (for S. Miyashita): miyasita@cypress.ne.jp

**Abstract**–Morphological development and growth of larval and juvenile Pacific bluefin tuna, *Thunnus thynnus*, were studied from laboratory-reared specimens. Average body length (BL) of newly hatched larvae was 2.83 mm and larvae grew on average 5.80 mm by 10 days, 10.62 mm by 20 days, and 35.74 mm by 30 days after hatching. Growth was especially accelerated after 20 days from hatching. Newly hatched larvae had small melanophores scattered over their bodies except for the finfold. On day 1 after hatching (3.35–3.74 mm BL), a characteristic melanophore pattern appeared and it was partially maintained until day 3 after hatching. At approximately 4 mm BL, larvae had developed melanophore patterns similar to those of preflexion *Thunnus* spp. larvae, such as melanophores on the dorsum of the gut, midlateral trunk, and tail, at the dorsal and ventral midlines of trunk and tail, and on the lower jaw. Erythrophores appeared at 4.63 mm BL in the caudal area. Jaw teeth appeared at 5–6 mm BL. The preopercular angle spine, anterior preopercular spine, and posttemporal spine, developed at approximately 7 mm BL, when erythrophores appeared on the trunk and tail. The notochord flexion occurred between 6 and 8 mm BL. At approximately 8 mm BL, erythrophores disappeared and juvenile coloration appeared on the trunk and tail, consisting of dense patches of melanophores at the dorsal and anal-fin bases, embedded melanophores, and melanophores at the periphery of the eye. The adult complement of fin-ray counts was attained at 10 mm BL, when the juvenile melanophore pattern was attained, although the pattern was not fully developed. Specimens larger than 20 mm BL did not have erythrophores. Squamation began at 27 mm BL and head spines disappeared by 38 mm BL.

Manuscript accepted 10 April 2001.
Fish. Bull. 99:601–616 (2001).

The Pacific bluefin tuna is a large scombrid that migrates transoceanically between the western and eastern Pacific Ocean (Orange and Fink, 1963; Fujita, 1998). The Pacific bluefin tuna, *Thunnus thynnus orientalis* (Temminck and Schlegel), is considered a separate subspecies from the Atlantic bluefin tuna, *T. thynnus thynnus* Linnaeus (Gibbs and Collette, 1967; Collette and Nauen, 1983). Recently, Cho and Inoue (1993) showed reproductive isolation between the two subspecies. Most recently, Collette (1999) considered these subspecies to be separate species on the basis of morphology and molecular data.

Our knowledge of the early life history of oceanic species, such as tunas, is incomplete. Morphological characteristics common to scombrid larvae are large head, gape, and eyes; development of head spination; and posterior migration of the anus (Collette et al., 1984). In studies of early-stage bluefin tuna (e.g. Yabe et al., 1966; Matsumoto et al., 1972; Richards and Potthoff, 1974; Kohno et al., 1982), melanophore patterns were used to identify tuna larvae. Erythrophore patterns were proposed as another characteristic by Ueyanagi (1966) and Matsumoto et al. (1972). However, ontogenetic changes of melanophore and erythrophore patterns and morphological characteristics have not sufficiently been investigated because of the difficulty in obtaining a complete series of wild specimens in their early developmental stages.

Morphological, physiological, and behavioral information has recently been collected from laboratory-reared Pacific bluefin tuna specimens (Harada et al., 1971a; Kaji et al., 1996; Takii et al., 1997; Kumai, 1998; Miyashita et al., 1998; Miyashita et al., 2000, in press; Sawada et al., 2000). The ability to produce a complete series of early-stage specimens provides an opportunity to enhance our understanding of the early development of bluefin tuna. In this study, we describe the morphological development and pigment patterns of reared larval and juvenile bluefin tuna.

## Materials and methods

Eggs were obtained on 10 July 1994 from bluefin tuna that had been caught off Ohshima when 20–40 cm in total length [0-year class] and that had been raised for seven years naturally in a net cage at the Ohshima Experiment Station of the Kinki University Fisheries Laboratory. The net cage was equipped with a polyvinyl chloride sheet surrounding the top 2 m of the net side wall to prevent eggs from flowing out of the net. Fertilized eggs were collected at the surface with a 330-μm mesh net. Eggs were transferred to a tank in a land-based nursery center and kept in a fine mesh net at 24.5°C for approximately 12 hours. The following morning, eggs were transferred to a 20-$m^3$ rearing tank and larvae hatched at

approximately 2200 h on 11 July. The first 24 hours after hatching were counted as day 0 (day-0) after hatching. Water temperature ranged from 24.5 to 27.7°C during the rearing period ($\bar{x}$=25.5°C, Fig. 1). The feeding scheme for larvae and juveniles was as follows: rotifers, *Brachionus rotundiformis,* from day 2 to day 22 after hatching, *Artemia* nauplii from day 10 to day 25, and other live fish larvae (*Oplegnathus fasciatus*) from day 12 to day 30.

At 1000 h daily from day 0 (12 July) to day 30 (11 August), 20 tuna were sampled. Before preservation in 5% formalin, individual computer-captured video images were made with a video camera attached to a binocular microscope while all specimens were under MS-222 anesthesia. Ten body parts were measured from these images: total length (TL), body length (BL), preanal length, body height at the base of the first dorsal spine, head length, head height, snout length, eye diameter, caudal peduncle depth, and upper jaw length. Video image measurements to $1/10^2$ mm with an accuracy of ±1% were obtained from NIH Image (version 1.61) image analysis software (NIH, 1997). Before and during notochord flexion, BL was measured from the tip of the upper jaw to the end of the notochord. After notochord flexion, BL was measured from the tip of the upper jaw to the posterior margin of the hypurals. All specimens ($n$=620, 3.49–37.78 mm BL) were used to determine growth estimates and to record pigmentation patterns and spine and fin development.

Sixty-nine specimens (5.26–33.68 mm BL) were stained with alizarin red-S to describe spine development, and squamation. Pigment patterns, especially the position of body melanophores in relation to myomeres, were determined for specimens from 3 to 16 days after hatching ($n$=180, 3.63–9.96 mm BL). At irregular intervals from day 0 to day 22, one to five fish (a total of 53 specimens, 3.38–26.76 mm BL) were examined while anesthetized with MS-222 to determine erythrophore distribution. Additional samples were examined at irregular intervals from hatching to day 2.

A representative series of specimens from this study was deposited in the Aquatic Natural History Museum of the Fisheries Research Station, Kyoto University (FAKU 129041–129075).




**Figure 1**

Body length, diet, and water temperature (W.T.) during the rearing of *Thunnus thynnus*. Vertical lines indicate ±SD ($n$=20).

## Results

### Structural characteristics and growth

Eggs, having a mean diameter of 1.02 mm (SD=0.02; $n$=20), were spherical and pelagic and had a single oil globule of 0.26 ±0.05 (mean ±SD) mm in diameter.

Newly hatched larvae measured 2.83 ±0.16 mm in BL (mean ±SD, $n$=20; Fig. 2A). At 3 days after hatching (3.81 ±0.14 mm BL), the mouth was developed and larvae began feeding. Yolk was present until day 5 (4.32 ±0.13 mm BL). Larvae grew to 5.58 ±0.35 mm by day 10, 9.34 ±1.36 mm by day 20 and 30.36 ±4.34 mm by day 30 after hatching (Fig.1). From hatching to day 20, growth rate was 0.33 mm/day, then increased to 2.10 mm/day from day 20 to day 30.

Notochord flexion began as early as 5.76 mm BL, and the largest preflexion larva was 5.83 mm BL. The smallest postflexion larva was 7.46 mm BL; the largest flexion larva, 8.65 mm BL. Thus, the flexion stage occurred from 6 to 8 mm BL (Fig. 2E).

The relative values of the nine body parts, except preanal length, increased during larval development, then subsequently declined (Fig. 3). The ratio of head length, head height, eye diameter, snout length, and upper jaw length to BL reached a maximum at 11–13 mm BL. Head length increased rapidly from 10% BL at hatching to 40% at 13 mm BL and then decreased slowly to 34 % BL at 40 mm BL. Head height also increased rapidly from 10% BL at hatching to 27% at 11 mm BL and decreased slowly to 23% BL at 40 mm BL. Eye diameter increased from 10% BL at hatching to 15% at 11 mm BL and then decreased slowly to 9% BL at 40 mm BL. Snout length rapidly increased from 2% BL at hatching to 14% at 12 mm BL and then decreased slowly to 9% BL at 40 mm BL. Upper jaw length increased rapidly from 5% BL at hatching to 20–30% at 12 mm BL and then decreased slowly to 14% BL at 40 mm BL. The




**Figure 2**

Development of *Thunnus thynnus* (**A**) newly hatched larva, 2.91 mm BL; (**B**) 1-day-old yolksac larva, 3.54 mm BL; (**C**) 4-day-old preflexion larva, 4.02 mm BL; (**D**) 7-day-old flexion larva, 5.34 mm BL; (**E**) 15-day-old postflexion larva, 7.23 mm BL; (**F**) 20-day-old postflexion larva, 8.64 mm BL; (**G**) 25-day-old juvenile, 13.55 mm BL.

relative depth of the caudal peduncle attained a maximum (ca. 7% BL) at 8 mm BL. In contrast, body height and total length were maximum in relation to BL at approximately 16 and 20 mm BL, respectively. Preanal length increased from 35–44% BL at 3–7 mm BL to about 65% BL at approximately 15 mm BL.




**Figure 3**

Measurements of *Thunnus thynnus*, shown as a percentage of body length (BL).

**Table 1**

Numbers of fin spines and rays of *Thunnus thynnus* larvae and juveniles. Roman and Arabic numerals show numbers of spines and soft rays, respectively.

| Size range (BL; mm) | First dorsal fin | Second dorsal fin | Dorsal finlet | Anal fin | Pectoral fin | Pelvic fin | Number of specimens examined |
|---|---|---|---|---|---|---|---|
| BL ≤5.99 | 0 | 0 | 0 | 0 | 0 | 0 | 10 |
| 6.00–6.99 | I–X | 0 | 0 | 0–6 | 0–13 | 0–I+1 | 17 |
| 7.00–7.99 | 0–XIV | 0–13 | 0–8 | 0–12 | 0–31 | 0–I+5 | 13 |
| 8.00–8.99 | IX–XIV | 12–14 | 6–9 | 7–14 | 11–31 | I+4–I+5 | 11 |
| 9.00–9.99 | XI–XV | 14 | 8–9 | 13–14 | 16–31 | I+5 | 8 |
| 10.00–19.99 | XIV–XV | 14 | 8–9 | 14 | 31–34 | I+5 | 14 |
| 20.00–29.99 | XV | 14 | 8–9 | 13–14 | 31–36 | I+5 | 19 |
| 30.00–37.78 | XV | 14 | 8–9 | 14 | 31–35 | I+5 | 8 |

## Fin development

The number of fin rays increased between 6 and 10 mm BL (Table 1). The smallest specimen possessing dorsal spines was 6.32 mm BL. Caudal-fin rays, second dorsal-, pelvic-, anal-, and pectoral-fin rays appeared at 7.10, 7.76, 8.00, 8.47, and 8.67 mm BL, respectively. The smallest specimen with an adult complement of fin-ray counts (Iwai et al., 1965; Collette and Nauen, 1983) was 9.68 mm BL, whereas the largest specimen with an incomplete fin-ray complement was 10.02 mm BL. The number of pectoral-fin rays for specimens >10 mm BL varied among individuals.

## Pigmentation

Newly hatched larvae (2.62–3.05 mm BL) had small melanophores scattered over the body, head, notochord, yolk, and oil globule, but not on the finfold (Fig. 2A).

On day 1 after hatching (3.35–3.74 mm BL) (Fig. 2B), dendritic melanophores were visible from snout to fore- and midbrain and on the dorsum of the gut; three clusters of melanophores occurred in the dorsal midline along with punctate melanophores; melanophores occurred along the ventral midline (punctate melanophores formed a line at the anterior caudal area and a cluster of melanophores occurred in the posterior caudal area); and melanophores appeared along the lateral midline of the trunk. The eyes were pigmented.

On day 2 after hatching (3.40–4.18 mm BL), the pattern of melanophores was transitional between those of 1-day and older preflexion larvae. Melanophores on the snout disappeared, whereas clusters of melanophores on the anteriodorsum of the trunk (second myomere), anteriordorsum of the caudal area (17th to 21st myomere) and posteriodorsum of the caudal area (30th to 39th myomere), as well as on the anterio- and midventral edge (14th to 26th myomere), posteroventral edge (28th to 39th myomere), and lateral midline (posterior to the 13th myomere) of the caudal area began to shrink (although they were still large [Tables 2–5]).

On day 3 after hatching, forebrain melanophores disappeared in all specimens >3.76 mm BL, whereas the smallest specimen lacking these melanophores was 3.63 mm BL. All specimens >3.82 mm BL had melanophores scattered on the dorsum of the gut, which formed a cap of melanophores over the gut in later stages. Clusters of melanophores on the dorsal and ventral midline, and lateral midline of the body now appeared punctate and additional punctate melanophores appeared in these areas (Tables 2–5). Melanophores were present on the midbrain, on the anteriodorsal side and on the base of the hindbrain; and several melanophores appeared on the lower side of the lower jaw. The smallest specimen with melanophores on the lower jaw was 3.41 mm BL, whereas the largest lacking them was 9.46 mm BL (day 16). Most specimens >5.5 mm BL had these lower jaw melanophores. On and after day 4 (3.92–4.43 mm BL), all specimens had the melanophore pattern of preflexion larvae (Fig. 2C).

Melanophores began to develop on several other areas in larvae larger than about 4.5 mm BL (Fig. 2, D–F). Embedded melanophores appeared in the lateral muscle at 4.49 mm BL (Table 6), upper jaw tip at 5.47 mm BL, operculum and preoperculum at 6.18 mm BL, the membrane of the first dorsal fin at 6.32 mm BL, forebrain at 6.83 mm BL, and the cleithral symphysis at 10.79 mm BL. The largest specimens lacking melanophores in these areas were those that were 6.44 mm BL (for upper jaw tip), 6.72 mm BL (for lower jaw tip), 6.94 mm BL (for operculum), 9.69 mm BL (for forebrain), 10.28 mm BL (for cleithral symphysis), and 10.49 mm BL (for preoperculum).

Melanophores forming a dorsal cap of the gut enlarged at about 6.5 mm BL and the rim reached the ventral surface of the gut in some specimens >7.5 mm BL. The body melanophore pattern changed with growth. Dorsal midline melanophores disappeared from the trunk from 4.5 mm BL to 7 mm BL, and ventral and lateral midline melanophores disappeared from the trunk (Tables 2–5).

Larvae showed partial juvenile pigment patterns from about 8 mm BL (Fig. 2F). Dorsolateral and dorsal midline melanophores appeared at the trunk and increased in frequency in the caudal area from 7.5 mm BL (Table 2). Embedded melanophores frequently appeared at the posterior trunk and caudal area (Table 6). Dense patches of melanophores appeared at the fin base of the first and second dorsal, and anal fin bases (Tables 7 and 8; these melanophores occurred on the myosepta), as well as at the periphery of the eye, especially below and behind the eye from about 8.0 mm BL. Embedded melanophores appeared near the notochord and neural and haemal spines, and melanophores at the lateral midline of body extended internally from about 8.2 mm BL. Generally, melanophores on the dorsal and ventral midline, lateral midline, and those embedded increased with BL (Tables 2–6).

The smallest specimen having juvenile pigmentation (i.e. densely pigmented patches appearing on the body) was 13.55 mm BL (Fig. 2G). More distinct body patches appeared and increased in number as juveniles grew.

In many larvae, melanophores were found in the area of the hypural bones and on the caudal finfold or fin rays. The smallest specimens having melanophores in the area of hypural bones and on the caudal finfold were 6.42 mm BL and 6.81 mm BL, respectively.

Erythrophores first appeared at 4.63 mm BL (6 days after hatching) on the tail: thirty-four erythrophores at the posteroventral edge, five at the posterolateral midline, and two at the posterodorsal edge (Table 9). As larvae grew, erythrophores appeared at the caudal finfold at 4.75 mm BL, on the lower jaw at 6.10 mm BL, and at the hypural plate at 6.96 mm BL; at the hypural plate and caudal finfold, erythrophores were observed only on specimens 6.96–7.20 and 4.75–7.20 mm BL, respectively. At the dorsal and ventral edge of the trunk and tail, erythrophores became larger and decreased in number in larvae >7 mm. At the ventral edge, adjacent erythrophores were united. At the lateral midline of the body, the number decreased at 8.5 mm BL. Erythrophores disappeared from the lower jaw at 11.26 mm BL, from the posterolateral midline and dorsal edge of the tail at 15.85 mm BL, and from the ventral edge of the tail at 19.72 mm BL. Erythrophores were not observed on the upper jaw. The number of erythrophores, especially the ventral edge erythrophores, de-

**Table 2**

Incidence of dorsal midline melanophores at the trunk and tail (%) in *Thunnus thynnus*. Each melanophore is

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 |
| 3.50–3.99 | | 14 | | | | | 7 | | | | | 14 | 7 | 7 | 7 | 7 | 36 | 29 | 36 |
| 4.00–4.49 | | 4 | 4 | | | | | | | 4 | 15 | | 4 | 4 | 7 | 7 | 15 | 41 | 22 |
| 4.50–4.99 | | | | | 4 | 4 | | | | | | | | | 4 | 4 | 7 | 15 | 7 |
| 5.00–5.49 | | | | | | | | | | | 8 | | | | 8 | 8 | 8 | | 17 |
| 5.50–5.99 | | 10 | 10 | | | | | | | | | | | 10 | 10 | 10 | | | |
| 6.00–6.49 | | | | | | | | | | | | | | | | | 20 | 30 | 30 |
| 6.50–6.99 | | | | | | | | | | | | | | | | | | | 17 |
| 7.00–7.49 | | | | | | | | | | | 10 | 10 | 10 | 10 | 20 | 20 | | 30 | |
| 7.50–7.99 | | | | | | | | | | 10 | 10 | 10 | 20 | 40 | 20 | 40 | 20 | 10 | 20 |
| 8.00–8.49 | | | | | | | 8 | 15 | 31 | 23 | 62 | 77 | 77 | 85 | 85 | 85 | 69 | 69 | 62 |
| 8.50–8.99 | 23 | 54 | 38 | 46 | 46 | 54 | 54 | 54 | 62 | 85 | 85 | 85 | 85 | 92 | 85 | 92 | 92 | 92 | 92 |
| 9.00–9.49 | 25 | 67 | 17 | 25 | 33 | 33 | 83 | 58 | 50 | 58 | 58 | 67 | 67 | 67 | 58 | 75 | 83 | 67 | 67 |
| 9.50–9.99 | 30 | 80 | 20 | 30 | 40 | 40 | 100 | 70 | 60 | 70 | 70 | 80 | 80 | 80 | 70 | 90 | 100 | 80 | 80 |

**Table 3**

Incidence of the ventral midline and ventrolateral melanophores of the trunk and tail (%) in *Thunnus thynnus*. Each melanophore not observed on the first to sixth myomeres.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 |
| 3.50–3.99 | 7 | 7 | | | | 7 | 7 | 7 | 14 | 14 | 29 | 7 | 7 | 50 | 7 | 21 | 21 |
| 4.00–4.49 | | | | 4 | 4 | | | | 7 | 7 | 22 | 22 | 11 | 15 | | 7 | 19 |
| 4.50–4.99 | | | | | | 10 | 5 | 5 | 0 | 10 | 10 | 29 | 24 | 19 | 19 | 24 | 33 |
| 5.00–5.49 | | | | | | | | | | | 8 | | | | 17 | 17 | 17 |
| 5.50–5.99 | | | | | | | | | | | 10 | 10 | | | | 30 | |
| 6.00–6.49 | | | | | | | | | | | | | 10 | 10 | | 30 | 10 |
| 6.50–6.99 | | | | | | | | | | | | | | | 6 | 11 | 6 |
| 7.00–7.49 | | | | | | | | | | | | | | 20 | | | 10 |
| 7.50–7.99 | | | | | | | | | | | | | | | | | |
| 8.00–8.49 | | | | | | | | | | | | | | | | | |
| 8.50–8.99 | | | | | | | | | | | | | | | 8 | 8 | 8 |
| 9.00–9.49 | | | | | | | | | | | | | | | | | |
| 9.50–9.99 | | | | | | | | | | | | | | | | | |

creased as the fish grew; no erythrophores were observed in specimens >20 mm BL.

## Head spination

Major head spines first appeared in specimens at 4–7 mm BL. The posterior preopercular angle spine and anterior preopercular spine appeared at 4.23 mm BL (Fig. 2D), and the posttemporal spine at 6.80 mm BL. The largest specimen lacking anterior and posterior preopercular spines was 4.48 mm BL, and the largest specimen lacking posttemporal spines, 7.21 mm BL (Fig. 2E). No other spines or spinelets of the head appeared in later developmental stages. The number of anterior and posterior preopercular spines increased 2–3 (nine maximum) and 5–7 (nine maximum), respectively, in the size range of 5–16 mm BL (Fig. 2, F

**Table 2**

recorded for the myomere on which it occurs. *n* indicates the number of specimens examined at each size range.

| Myomere | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 29 | 29 | 7 | | | 7 | | | 7 | 7 | 14 | 14 | 21 | 21 | 36 | 43 | 14 | 14 | 7 | 14 | 14 |
| | 7 | 22 | 7 | | 7 | 19 | 15 | 4 | 11 | 19 | 4 | 19 | 19 | 33 | 30 | 26 | 30 | 15 | 19 | 27 |
| 7 | 11 | 4 | 4 | 4 | 7 | 11 | 7 | 11 | 15 | 4 | 11 | 15 | 19 | 15 | 19 | 22 | 7 | 7 | | 21 |
| 8 | 8 | 17 | | | 25 | 8 | 17 | 8 | 17 | 8 | 8 | 8 | 17 | 17 | 17 | 25 | 25 | | 8 | 12 |
| 10 | | | | | 10 | | | 10 | 10 | 10 | | 30 | 10 | 20 | 10 | 10 | | | | 10 |
| | 20 | 10 | 20 | | 20 | | 30 | 10 | 20 | 10 | 10 | 50 | 40 | 50 | 10 | 10 | | | 10 | 10 |
| 22 | | 11 | | 6 | | 6 | | 6 | 6 | 6 | 11 | 6 | 22 | 22 | 6 | 11 | 22 | 6 | | 18 |
| 10 | | | | | 10 | | | 10 | 20 | | 10 | | | 20 | 10 | 30 | 70 | 30 | 20 | 10 |
| 20 | 20 | 20 | 10 | | | 20 | 10 | 20 | 0 | 10 | 30 | 20 | 20 | 10 | 30 | 40 | 10 | 40 | 10 | 10 |
| 54 | 38 | 38 | 31 | 23 | 38 | 31 | 23 | 23 | 15 | 31 | 8 | 31 | 15 | 23 | 8 | 15 | 15 | 8 | 38 | 13 |
| 92 | 77 | 77 | 77 | 69 | 69 | 69 | 69 | 69 | 62 | 54 | 69 | 69 | 69 | 54 | 54 | 62 | 46 | 38 | 23 | 13 |
| 83 | 67 | 58 | 75 | 58 | 83 | 58 | 58 | 67 | 58 | 75 | 50 | 50 | 33 | 33 | 25 | 58 | 67 | 33 | 17 | 12 |
| 100 | 80 | 70 | 90 | 70 | 100 | 70 | 70 | 80 | 70 | 90 | 60 | 60 | 40 | 40 | 30 | 70 | 80 | 40 | 20 | 10 |

**Table 3**

is recorded for the myomere on which it occurs. *n* indicates the number of specimens examined at each size range. Melanophores were

| Myomere | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 29 | 14 | 7 | 57 | 7 | 64 | 29 | 14 | 21 | 79 | 50 | 36 | 57 | 50 | 36 | 43 | 36 | 14 |
| 15 | 15 | 37 | 11 | 26 | 37 | 41 | 41 | 30 | 44 | 44 | 26 | 26 | 48 | 30 | 33 | 52 | 27 |
| 19 | 33 | 29 | 29 | 24 | 43 | 38 | 38 | 52 | 24 | 57 | 43 | 43 | 19 | 33 | 29 | 57 | 21 |
| 17 | 8 | 33 | 17 | 17 | 42 | 42 | 17 | 25 | 83 | 50 | 50 | 50 | 33 | 58 | 33 | 25 | 12 |
| 20 | 30 | 40 | 60 | 10 | 10 | 60 | 60 | 40 | 70 | 60 | 60 | 30 | 20 | 30 | 20 | 20 | 10 |
| 30 | 10 | 20 | 60 | 30 | 30 | 10 | 30 | 30 | 40 | 50 | 20 | 0 | 10 | 30 | 20 | 0 | 10 |
| 6 | 11 | 22 | 28 | 39 | 39 | 33 | 56 | 28 | 44 | 28 | 28 | 22 | 50 | 22 | 17 | 39 | 18 |
| | 10 | 30 | 50 | 20 | 50 | 60 | 60 | 40 | 60 | 80 | 50 | 40 | 60 | 30 | 30 | 30 | 10 |
| 10 | 10 | 20 | 30 | 20 | 40 | 40 | 50 | 50 | 50 | 90 | 70 | 60 | 50 | 70 | 40 | 60 | 10 |
| | | 8 | 15 | 15 | 23 | 54 | 46 | 62 | 54 | 69 | 62 | 62 | 54 | 62 | 46 | 46 | 13 |
| 8 | | 15 | 8 | 23 | 31 | 54 | 46 | 62 | 46 | 85 | 77 | 54 | 77 | 77 | 85 | 31 | 13 |
| | | | | 8 | 25 | 33 | 42 | 58 | 33 | 58 | 67 | 67 | 67 | 75 | 50 | 42 | 12 |
| | | | | 10 | 40 | 40 | 40 | 60 | 70 | 80 | 70 | 70 | 70 | 70 | 60 | 40 | 10 |

and G). The number of anterior and posterior preopercular spines decreased at 17 mm BL. At >19.23 mm BL, no anterior preopercular spines were observed, except in one specimen (24.00 mm BL) that had two anterior preopercular spines. Juveniles had three posttemporal spines when these spines were fully developed. The largest specimen with a posttemporal spine was 24.23 mm BL, and the largest specimen examined, 37.78 mm BL, had no spines on its head.

## Teeth and squamation

Upper and lower jaw teeth were first observed at 5.35 and 6.32 mm BL, respectively. Palatine teeth appeared at 7.20 mm BL and most specimens >9 mm BL had fully developed palatine teeth. At >18.00 mm BL, specimens did not have these teeth.

Squamation was incomplete in the size range examined in this study. The smallest scaled specimen (27.37 mm BL)

**Table 4**

Incidence of melanophores on the left lateral side of the trunk and tail (%) in *Thunnus thynnus*. Each melanophore is recorded for the the first to eighth myomeres.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 |
| 3.50–3.99 | | | 7 | 7 | 7 | 14 | 7 | 29 | 14 | 21 | 7 | 29 | 14 | 7 | 14 |
| 4.00–4.49 | | 4 | | | 11 | 4 | | | | 7 | 11 | 11 | 11 | 11 | 11 |
| 4.50–4.99 | | | | | 5 | 5 | 5 | | 5 | 10 | 5 | 10 | 5 | 29 | 10 |
| 5.00–5.49 | | 8 | | | | 8 | | | | 8 | | 8 | | 17 | 8 |
| 5.50–5.99 | | | | | 20 | | 20 | 20 | 20 | 10 | | 40 | | | 10 |
| 6.00–6.49 | | | | | | | | | 20 | | | 10 | | 20 | 10 |
| 6.50–6.99 | | | | 6 | 0 | 6 | 6 | 6 | | 28 | 11 | | | 17 | 22 |
| 7.00–7.49 | | | | | | | | | 20 | | | | 20 | 20 | |
| 7.50–7.99 | | | | | | | | | | 20 | 10 | 20 | | | 20 |
| 8.00–8.49 | | | | | | 15 | 8 | | | 15 | | 8 | 8 | 8 | 15 |
| 8.50–8.99 | | | | | | | | | | 8 | | 15 | 8 | 8 | 8 |
| 9.00–9.49 | | | | | | | | | | 8 | | | | 17 | 17 |
| 9.50–9.99 | 10 | | | | | | | | | | | 10 | | 10 | 20 |

**Table 5**

Incidence of melanophores on the right lateral side of the trunk and tail (%) in *Thunnus thynnus*. Each melanophore is recorded for on the first to nineth myomeres.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | |
| 3.50–3.99 | 0 | 7 | 7 | 7 | 14 | 7 | 21 | 14 | 21 | 14 | 29 | 14 | 7 | 14 | |
| 4.00–4.49 | 4 | | | 11 | 4 | 4 | | | 11 | 4 | 15 | 11 | 11 | 11 | |
| 4.50–4.99 | | | | 5 | | | | 5 | 10 | 5 | 10 | 5 | 19 | 10 | |
| 5.00–5.49 | | | | | 17 | 8 | | | 17 | | 8 | | 33 | 8 | |
| 5.50–5.99 | 10 | | | 20 | | 10 | 20 | 10 | | 10 | 40 | | | 10 | |
| 6.00–6.49 | | | | | | | | 20 | 10 | | 10 | | 30 | 10 | |
| 6.50–6.99 | | 6 | 6 | | 6 | 6 | 6 | | 28 | 11 | | | 17 | 22 | |
| 7.00–7.49 | | | | | | | | 20 | 10 | 20 | | 10 | 10 | | |
| 7.50–7.99 | | | | | 10 | | | 10 | | 10 | 20 | 20 | | 10 | |
| 8.00–8.49 | | | | | | | | | 31 | | | | | 15 | |
| 8.50–8.99 | | | | | 15 | 15 | | | | | 31 | 31 | 31 | 15 | |
| 9.00–9.49 | | | | | 8 | | | | 8 | | | | 17 | 17 | |
| 9.50–9.99 | | | | | | | | | | | 10 | 10 | | 30 | |

had anterior lateral line scales. Scales near the anterior part of the lateral line appeared at 30.31 mm BL, and postorbital scales at 30.81 mm BL.

## Discussion

### Development and growth

*Thunnus thynnus* eggs were reported to have a smooth chorion, narrow perivitelline space, homogeneous yolk, and a single oil globule (Miyashita et al., 2000), and this was reconfirmed in our study. These characteristics are similar to those found in other *Thunnus* eggs: *T. thynnus* from the Mediterranean (Podoa, 1956), *T. obesus* (Kikawa, 1953), *T. albacares* (Harada et al., 1971b; Mori et al., 1971), and *T. alalunga* (Yoshida and Otsu, 1963).

The growth strategy of bluefin tuna apparently is to develop foraging structures before other organs and at an early stage to enable feeding on larger organisms. Parts of the head were large from early development (Fig. 3) as mentioned by Collette et al. (1984) and reached maximum

**Table 4**

myomere on which it occurs. *n* indicates the number of specimens examined at each size range. Melanophores were not observed on

| Myomere | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 14 | 36 | 7 | | 21 | 7 | 7 | 14 | 14 | 7 | 14 | 7 | | 14 | 7 | | 14 |
| | 30 | 11 | 7 | 15 | 4 | 4 | 11 | | 11 | 4 | 7 | 11 | 7 | 7 | 7 | 27 |
| 14 | 10 | 19 | 14 | 5 | 5 | | 5 | 5 | 19 | 10 | 14 | 5 | 10 | | 5 | 21 |
| 17 | 8 | 8 | 8 | 33 | | | | 8 | 17 | 25 | 8 | 8 | | | | 12 |
| 20 | | 10 | 10 | 20 | | 20 | 40 | | 10 | 10 | 30 | 10 | | | | 10 |
| | 10 | 10 | | 20 | 10 | 20 | | | 10 | | | | | | | 10 |
| 11 | 11 | 6 | 11 | 17 | 6 | 6 | 11 | 6 | 11 | 6 | 17 | 11 | | 6 | | 18 |
| | | 20 | 10 | 10 | | | 10 | | 10 | 20 | 10 | 10 | 10 | | 10 | 10 |
| | | 20 | 20 | 10 | 20 | 20 | | 30 | | 10 | 10 | 40 | | 20 | 10 | 10 |
| 8 | | 8 | 15 | 8 | | 15 | 31 | 31 | | 8 | | 8 | | 8 | | 13 |
| 15 | 8 | 8 | 31 | 15 | 23 | 15 | | 23 | 15 | 15 | 8 | 8 | 8 | 8 | 8 | 13 |
| 8 | 8 | 17 | 17 | 25 | 25 | 42 | 58 | 33 | 50 | 33 | 50 | 33 | 33 | 50 | 42 | 12 |
| | 10 | 10 | | 20 | 20 | 20 | | 50 | 60 | 40 | 50 | 40 | 30 | 20 | 50 | 10 |

**Table 5**

the myomere on which it occurs. *n* indicates the number of specimens examined at each size range. Melanophores were not observed

| Myomere | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 14 | 36 | 7 | | 21 | 7 | 7 | 14 | 14 | 14 | 7 | 7 | | 14 | 7 | 7 | 14 |
| 4 | 30 | 19 | 7 | 15 | 7 | 4 | | | 15 | 4 | 7 | 15 | 19 | 7 | 7 | 27 |
| 14 | | 10 | 14 | | 5 | | | 5 | 19 | 5 | 5 | | | | 5 | 21 |
| 17 | 25 | 17 | 17 | 42 | | | 25 | 8 | 17 | 8 | 17 | 8 | | | | 12 |
| 10 | | | 30 | 20 | | 20 | 10 | | 10 | 50 | 30 | 10 | | | | 10 |
| 10 | 10 | 10 | | 10 | 20 | 10 | | 10 | 10 | 10 | | | | | | 10 |
| 6 | 11 | 11 | 11 | 17 | 6 | 6 | 11 | 6 | 11 | 6 | 17 | 11 | | 6 | | 18 |
| | | 10 | 10 | 30 | 10 | 10 | 10 | 10 | | | 20 | 20 | 20 | | | 10 |
| | 10 | 10 | 30 | 20 | 10 | | | 10 | | 10 | 20 | 30 | | 30 | 20 | 10 |
| 31 | | 8 | 8 | 8 | | 15 | 15 | | | 8 | | 8 | | 8 | | 13 |
| 8 | 15 | 15 | 54 | 23 | 31 | 46 | 46 | 31 | 31 | 31 | 8 | | 8 | 23 | 15 | 13 |
| 8 | 17 | 8 | 8 | 25 | 25 | 42 | 50 | 50 | 42 | 25 | 42 | 25 | 25 | 42 | 25 | 12 |
| | 30 | 20 | | 20 | 20 | 10 | 10 | 30 | 30 | 40 | 50 | 50 | 30 | 20 | 50 | 10 |

ratios in relation to BL at sizes smaller than other body parts, such as total length and body height, except for the caudal peduncle depth (Fig. 4).

Another ontogenetic characteristic of growth in *T. thynnus*, the posterior migration of the anus (Collette et al., 1984), was also confirmed: the anus initially positioned in the anterior part of the body, was located at the center of the body at late flexion, and thereafter in the posterior part of the body (Fig. 3).

Kaji et al. (1996) examined the relative growth of larvae of the Pacific bluefin tuna at 3–14 mm BL and reported that body proportions showed constant relative growth values from 10 mm BL, except for preanal length. Our results showed that at 10 mm BL, constant ratios had not been reached in relation to BL. This difference may be due to an insufficient number of larger specimens in Kaji et al.'s (1996) study. Even at the juvenile stage, ratios in relative BL decreased gradually, except for a few cases (Fig. 3), suggesting that body proportions of juvenile bluefin tuna are different from those of adult fish.

Early growth is rapid in *T. thynnus* compared with other marine fishes cultured in Japan (e.g. red sea bream,

**Table 6**

Incidence of melanophores embedded in the lateral muscle and on the myosepta (%) in *Thunnus thynnus*. Each melanophore is recorded on the first to nineth myomeres.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | |
| 3.50–3.99 | | | | | | | | | | | | | | | |
| 4.00–4.49 | | | | | | | | | | 4 | | | | | |
| 4.50–4.99 | 4 | 4 | 4 | 7 | 4 | | | 4 | 4 | 4 | 4 | | 7 | | |
| 5.00–5.49 | | | | | | 8 | | | 8 | 8 | | | | | |
| 5.50–5.99 | | | 10 | | | | | | | | 10 | | | | |
| 6.00–6.49 | | | | 10 | | 10 | 10 | 10 | 10 | | | | | | |
| 6.50–6.99 | | | | | | | | | | | | | | | |
| 7.00–7.49 | 10 | 10 | | 10 | | | 10 | 20 | 10 | 10 | | 10 | | | |
| 7.50–7.99 | | | 10 | | | 10 | | | | 10 | | 10 | | | |
| 8.00–8.49 | | | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 15 | 23 | 31 | 15 | |
| 8.50–8.99 | | | 8 | 0 | 8 | 8 | 0 | 0 | 15 | 31 | 38 | 46 | 54 | 62 | |
| 9.00–9.49 | | | 8 | 8 | 8 | 8 | 25 | 33 | 33 | 42 | 50 | 50 | 50 | 58 | |
| 9.50–9.99 | | | 10 | 10 | 10 | 10 | 30 | 40 | 40 | 50 | 60 | 60 | 60 | 70 | |

**Table 7**

Incidence of melanophores on the dorsal fin base (%) in *Thunnus thynnus*. Each melanophore is recorded for the nearest myomere. for larvae <7.50 mm BL.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 |
| 7.50–7.99 | 10 | 10 | 10 | 10 | 10 | 20 | 20 | 20 | 30 | 20 | 30 | 20 | 30 | 20 | 20 | 20 | 20 | 10 |
| 8.00–8.49 | 8 | 8 | 8 | 8 | 8 | 15 | 15 | 15 | 23 | 15 | 23 | 15 | 23 | 15 | 15 | 15 | 15 | 8 |
| 8.50–8.99 | | 8 | 15 | 31 | 46 | 46 | 46 | 46 | 46 | 54 | 38 | 54 | 31 | 31 | 31 | 31 | 46 | 46 |
| 9.00–9.49 | 8 | 17 | 25 | 50 | 58 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 |
| 9.50–9.99 | | 20 | 40 | 70 | 80 | 100 | 60 | 60 | 60 | 60 | 100 | 100 | 100 | 100 | 100 | 90 | 100 | 100 |

**Table 8**

Incidence of melanophores on the anal fin base (%) in *Thunnus thynnus*. Each melanophore is recorded for the nearest myomere. *n* indicates the number of specimens examined at each size range. Melanophores were not observed on the first to twenty-first myomeres and for larvae <7.0 mm BL.

| Size range (BL; mm) | Myomere | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 7.00–7.49 | | | 10 | 10 | 20 | 10 | 20 | 20 | 20 | 10 | 10 | 10 | 10 | 10 | | | | | 10 |
| 7.50–7.99 | 10 | 20 | 30 | 30 | 50 | 40 | 60 | 40 | 40 | 40 | 40 | 20 | 30 | 20 | 20 | 20 | 10 | | 10 |
| 8.00–8.49 | | 15 | 8 | 15 | 31 | 38 | 31 | 62 | 69 | 54 | 62 | 62 | 46 | 31 | | 15 | | | 13 |
| 8.50–8.99 | | 8 | 8 | 8 | 8 | 31 | 31 | 38 | 62 | 69 | 69 | 69 | 69 | 69 | 62 | 62 | 38 | 23 | 13 |
| 9.00–9.49 | | | | | 17 | 33 | 42 | 58 | 75 | 83 | 83 | 75 | 67 | 67 | 50 | 42 | 33 | 25 | 12 |
| 9.50–9.99 | | | | | 20 | 40 | 50 | 70 | 90 | 100 | 100 | 90 | 80 | 80 | 60 | 50 | 40 | 30 | 10 |

**Table 6**

for the myomere in which it occurs. *n* indicates the number of specimens examined at each size range. Melanophores were not observed

| Myomere | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| | | | | | | | | | | | | | | | | 14 |
| | | | | | | | | | | | | | | | | 27 |
| | | | | | | | | | | | | | | | | 21 |
| 8 | | | | | | | | | | | | | | | | 12 |
| | | | | | | | | | | | | | | | | 10 |
| | | 10 | | | | | | | | | 10 | | | | | 10 |
| | | 6 | | | | 6 | | | | | | | 11 | 11 | | 18 |
| 10 | 10 | 10 | | | | 10 | 10 | | | | | | 10 | | | 10 |
| | 10 | 10 | 10 | 10 | 30 | 20 | 30 | 20 | | 10 | 30 | 20 | 20 | 20 | 30 | 10 |
| 23 | 23 | 31 | 23 | 38 | 31 | 54 | 38 | 23 | 38 | 15 | 38 | 38 | 31 | 15 | 15 | 13 |
| 69 | 62 | 69 | 77 | 77 | 77 | 77 | 77 | 77 | 69 | 62 | 62 | 62 | 62 | 46 | 23 | 13 |
| 75 | 75 | 83 | 83 | 75 | 83 | 75 | 75 | 75 | 75 | 75 | 75 | 58 | 58 | 67 | 33 | 12 |
| 90 | 90 | 100 | 100 | 90 | 100 | 90 | 90 | 90 | 90 | 90 | 90 | 70 | 70 | 80 | 40 | 10 |

**Table 7**

*n* indicates the number of specimens examined at each size range. Melanophore was not observed on the first to fourth myomeres and

| Myomere | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | *n* |
| 10 | 10 | 20 | 20 | 30 | 20 | 30 | 20 | 20 | 30 | 20 | 10 | 10 | | | | | 10 |
| 8 | 8 | 15 | 15 | 23 | 15 | 23 | 15 | 15 | 23 | 15 | 8 | 8 | | | | | 13 |
| 31 | 46 | 38 | 46 | 46 | 38 | 46 | 46 | 38 | 38 | 31 | 46 | 23 | | 8 | 0 | 15 | 13 |
| 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 58 | 58 | 58 | 42 | 42 | 8 | | 12 |
| 90 | 90 | 100 | 100 | 100 | 100 | 80 | 80 | 90 | 80 | 80 | 70 | 60 | 20 | 30 | 10 | 20 | 10 |

*Pagrus major*, and white trevally, *Pseudocaranx dentex* [Sawada unpubl. data; Fig. 5]). Rapid growth in bluefin tuna is pronounced after day 20 ($\bar{x}$ =9.34 mm BL on day 20) when larvae begin to metamorphose into juveniles. This type of growth phase seems common to scombrids, for example, *T. albacares* (Harada et al, 1971b; Kaji et al., 1999), *Scomber japonicus* (Watanabe, 1970), *Auxis tapeinosoma* (=*A. rocheri*, Harada et al., 1973), and *Sarda orientalis* (Harada et al., 1974) and reaches its height by the time of development of external physical features and the digestive system (Kohno et al., 1984; Miyashita et al., 1998), and by the time of development of their selective feeding characteristics (Sawada et al., 2000).

On day 19 ($\bar{x}$ =10.17 mm TL), the mouth size of *T. thynnus* averaged 2.54 mm according to Shirota's (1970) calculation (mouth size index=upper jaw length $\times 2^{0.5}$). This size is comparable to that of other scombrids, such as *T. albacares* (3.3 mm) and *Katsuwonus pelamis* (3.0 mm) at 10 mm TL, although much larger than that of other species (see Table 2 of Shirota, 1970). The digestive system of *T. thynnus* develops earlier than that of other fishes (Richards and Dove, 1971; Miyashita et al., 1998) and attains an adultlike structure by the juvenile stage, thus allowing bluefin tuna to use food efficiently early in development.

The development of pigment, spines, jaw teeth, and squamation in *T. thynnus* is summarized in Figure 6. The jaw teeth of *T. thynnus* first appeared at 5.35 mm BL (day 10), the palatine teeth at 7 mm BL (flexion stage), and were fully developed at 9 mm BL (postflexion stage). At 7 mm BL, *T. thynnus* larvae fed on other fish larvae and at 8 mm BL (postflexion stage), they were cannibalistic. Teeth appearance and development corresponded to the piscivorous stage during the fast-growth phase after the postflexion stage. Future improved laboratory-rearing techniques

**Table 9**

Number of erythrophores on the body of *Thunnus thynnus* larvae and juveniles.

| Body length (mm) | Dorsal edge | Ventral edge | Lateral midline | Lower jaw | Hypural plate | Caudal fin finfold |
|---|---|---|---|---|---|---|
| 3.38 | | | | | | |
| 3.41 | | | | | | |
| 4.63 | 2 | 34 | 5 | | | |
| 4.75 | 8 | 32 | 4 | | | 1 |
| 5.02 | 4 | 2 | | | | |
| 5.09 | 4 | 36 | | | | |
| 5.11 | 3 | 54 | 4 | | | |
| 5.35 | | 24 | 4 | | | |
| 5.78 | 1 | 2 | 4 | | | |
| 6.10 | 1 | 24 | 5 | 4 | | |
| 6.72 | 3 | 1 | 6 | | | |
| 6.91 | 1 | 1 | 4 | | | |
| 6.96 | 2 | 17 | 6 | 2 | 2 | |
| 6.96 | | 1 | 8 | 3 | | |
| 7.03 | 8 | | 6 | | 1 | |
| 7.06 | 8 | 12 | 8 | 1 | 5 | 2 |
| 7.12 | | 8 | 10 | | 5 | |
| 7.20 | 6 | 8 | 6 | 1 | 3 | 1 |
| 7.92 | 1 | 8 | 6 | 1 | | |
| 8.16 | | 6 | 8 | 3 | | |
| 8.41 | 1 | 11 | 8 | 3 | | |
| 8.74 | 1 | 4 | 5 | | | |
| 8.81 | 4 | 7 | 34 | 2 | | |
| 8.92 | | 3 | 3 | 1 | | |
| 9.02 | | | | 1 | | |
| 9.35 | 3 | 12 | 5 | 3 | | |
| 9.38 | | 3 | 2 | 3 | | |
| 9.56 | | 2 | 2 | 1 | | |
| 9.60 | | 9 | 3 | 1 | | |
| 9.70 | 5 | 1 | 6 | | | |
| 9.93 | 2 | 2 | 2 | | | |
| 10.03 | | 1 | 23 | | | |
| 10.56 | 4 | 1 | 2 | 7 | | |
| 10.97 | 4 | 1 | 3 | | | |
| 11.26 | 1 | 15 | 12 | 2 | | |
| 11.71 | | | | | | |
| 12.20 | | | 4 | | | |
| 13.71 | | 5 | | | | |
| 14.00 | | | | | | |
| 14.79 | | 3 | 1 | | | |
| 14.83 | | 1 | | | | |
| 15.33 | 2 | 4 | 1 | | | |
| 15.49 | | | | | | |
| 15.85 | 1 | 3 | 2 | | | |
| 16.86 | | | | | | |
| 18.73 | | 1 | | | | |
| 19.71 | | | | | | |
| 19.72 | | 2 | | | | |
| 20.40 | | | | | | |
| 21.87 | | | | | | |
| 24.85 | | | | | | |
| 25.89 | | | | | | |




**Figure 4**

Schematic representation of the relative growth (length of body parts in relation to body length) in hatchery-reared bluefin tuna, *Thunnus thynnus*. Flexion larva subdivision (FLS) and numerical complement of fin rays (NCF) are also shown. In relative growth, □ = the peak value of body part proportion (in relation to BL) and ■ = the attainment of a constant value of body part proportion (in relation to BL).

will allow further study of the structure and development of feeding-related bony elements for *T. thynnus*, as has been the case for *S. japonicus* (Kohno et al., 1984) and *Lates calcarifer* (Kohno et al., 1996).

## Pigmentation

Accurate identification of *Thunnus* larvae requires an extensive knowledge of individual, growth-associated, and geographic variations in melanophore and erythrophore patterns. We obtained information on the individual and growth-associated variations of these patterns from laboratory-reared specimens.

The melanophore distribution pattern of *T. thynnus* larvae and juveniles showed four distinct characteristic periods of development: newly hatched, 1–3 days after hatching, preflexion to postflexion, and juvenile.

*T. thynnus thynnus* larvae <3.99 mm BL from the Mediterranean had embedded melanophores (Kohno et al., 1982); but we did not observe these in our specimens until 4.49 mm BL. This difference may be a subspecific difference between *T. thynnus orientalis* and *T. thynnus thynnus*, or it may be due to insufficient numbers of specimens larger than 6.0 mm BL in Kohno et al.'s study, or to the experimental culture product as mentioned later.

Wild-caught *T. thynnus orientalis* larvae (3.15–8.20 mm BL) most frequently had two melanophores both on the dorsal and ventral edges of the trunk and tail (Nishikawa, 1985). This pattern resembles that of *T. thynnus thynnus* larvae from the Mediterranean (2.53–5.25 mm BL, Kohno et al., 1982). But our cultured specimens in the same size range had greater numbers of these melanophores (Tables 2 and 3). Nishikawa (1985) reported the appearance of melanophores at the dorsal fin somewhat earlier than we observed their appearance, but their appearance on the lower and upper jaws occurred in the same size range in both studies where specimens were examined from the same population. Laboratory-reared fish larvae have been shown to have greater pigmentation than wild-caught larvae (e.g. *Pagrus major* [Fukuhara and Kuniyuki, 1978], *Sparus sarba* [Kinoshita, 1986], *Parapristipoma trilineatum* [Kimura and Aritaki, 1985], and *Nibea mitsukurii* [Kinoshita and Fujita, 1988]). Thus, the source of specimens, wild-caught or laboratory-reared, may account for observed differences in larval pigmentation between our study and Nishikawa's study (1985).




**Figure 5**

Growth of bluefin tuna (*Thunnus thynnus*), red sea bream (*Pagrus major*), and white trevally (*Pseudocaranx dentex*). Red sea bream and white trevally were reared at the same temperature range (24–27°C) as bluefin tuna (Sawada, unpubl. data).

A slight difference in the occurrence of lower jaw, trunk, and tail melanophores has been reported among *T. thynnus* from the Mediterranean (Kohno et al., 1982), the Atlantic (Richards and Potthoff, 1974), and the Pacific (Matsumoto et al., 1972). Lower jaw melanophores appeared in all specimens larger than 3.0 mm BL from the Atlantic




**Figure 6**

Schematic representation of the development of pigment, spines, jaw teeth, and squamation in hatchery-reared bluefin tuna (*Thunnus thynnus*). □ = appearance of pigment and spines was subject to individual variation; ■ = pigment and spines were present in all specimens examined.

and larger than 4.0 mm BL from the Pacific. In our specimens, the incidence of these melanophores was 7.1% at 3.50–3.99 mm BL, 29.6% at 4.00–4.49 mm BL, 57.1% at 4.50—4.99 mm BL, 83.3% at 5.00–5.49 mm BL, and more than 90% in specimens >5.50 mm BL. The largest specimen we observed lacking these melanophores was 9.46 mm BL. Our data are similar to those for specimens from the Mediterranean. The difference in lower jaw melanophore distribution between our specimens and those of Matsumoto et al. (1972) might be explained by geographic variation of the melanophore pattern in the Pacific bluefin tuna. Further study is needed to confirm this hypothesis.

Ueyanagi (1966) reported species-specific characteristics of the erythrophore distribution pattern of Thunninae larvae from the Pacific Ocean: *T. alalunga* consistently had more erythrophores on the dorsal edge of the trunk and tail in front of the caudal peduncle than did *T. albacares*, which had only one or two erythrophores at the caudal peduncle; *T. thynnus* and *T. obesus* had patterns transitional between those of *T. alalunga* and *T. albacares*. Our data for *T. thynnus* generally agreed with those of Ueyanagi (1966); however, the number of dorsal erythrophores in one individual ranged more (0–8) than in Ueyanagi's study (1–5). In addition, ventral erythrophores appeared in large numbers (more than 20) at the preflexion stage (4.63–6.10 mm BL). From examinations of tuna larvae taken in Hawaiian waters, Matsumoto et al. (1972) considered the erythrophore pattern useful as a morphological characteristic for identifying tuna larvae. At present, however, information is limited on erythrophore patterns of other Thunninae larvae and juveniles and on the difference of these patterns in wild-caught and laboratory-reared specimens. Thus collection of such information is needed to establish the erythrophore pattern as a species-identifying characteristic of Thunninae larvae and juveniles.

*Thunnus thynnus* did not have xanthophores from hatching to the preflexion stage. However, *T. albacares* (Harada et al., 1971b; Mori et al., 1971) and *T. obesus* (Yasutake et al., 1973) have clusters of xanthophores in the finfolds of both the dorsal and ventral fins, and on the dorsal body, respectively. Thus, the xanthophore pattern can be used for distinguishing these *Thunnus* species at the preflexion stage.

Much additional work on development of tunas under controlled conditions, as well as the study on wild-caught materials, is needed to understand their early life history. We believe recent progress in the technology of rearing tunas will yield important information on the early life history of *Thunnus* spp.

## Acknowledgments

We would like to express our thanks to I. Nakamura (Kyoto University), I. Kinoshita (Kochi University), K. L. Main (Mote Marine Laboratory), anonymous reviewers, and the editorial office of *Fishery Bulletin* for their helpful suggestions and advice. This paper is dedicated to the late T. Harada, who was the previous professor and director of the Fisheries Laboratory of Kinki University and who was one of the leaders of our tuna aquaculture study.

## Literature cited

Cho, S., and S. Inoue.
1970. Intra- and interspecific restriction fragment length polymorphism in mitochondrial genes of *Thunnus* tuna species. Bull. Nat. Res. Inst. Far. Seas Fish. 30:207–225.

Collette, B. B.
1999. Mackerels, molecules, and morphology. Soc. Fr. Ichtyol. 25:149–164.

Collette, B. B., and C. E. Nauen.
1983. FAO species catalogue. Vol. 2: Scombrids of the world: an annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date, 137 p. FAO Fish. Synop. 125, FAO, Rome.

Collette, B. B., T. Potthoff, W. J. Richards, S. Ueyanagi, J. L. Russo, and Y. Nishikawa.
1984. Scombroidei: development and relationships. *In* Ontogeny and systematics of fishes (H. G. Moser, W. J. Richards, D. M. Cohen, M. P. Fahay, A. W. Kendall Jr. and S. L. Richardson, eds.), p. 591–620. Am. Soc. Ichthyol. Herpetol., Spec. Publ. 1.

Fujita, K.
1998. Species and ecology of tunas. *In* From the production to the consumption of tunas (S. Ono, ed.), p. 1–49. Seizando, Tokyo.

Fukuhara, O., and K. Kuniyuki.
1978. Morphological development in the wild larvae of Madai, *Chrysophyrys major*, as compared to that in the laboratory-reared larvae. Bull. Nansei Reg. Fish. Res. Lab. 11:19–25.

Gibbs, R. H., Jr. and B. B. Collette.
1967. Comparative anatomy and systematics of the tunas, genus *Thunnus*. Fish. Bull. 66:65–130.

Harada, T., H. Kumai, K. Mizuno, O. Murata, M. Nakamura, S. Miyashita, and H. Furutani.
1971a. On the rearing of young bluefin tuna. Mem. Fac. Agric. Kinki Univ. 4:153–157. [In Japanese with English summary.]

Harada, T., K. Mizuno, O. Murata, S. Miyashita, and H. Furutani.
1971b. On the artificial rearing of larvae in yellowfin tuna. Mem. Fac. Agri. Kinki Univ., p. 145–152. [In Japanese with English summary.]

Harada, T., O. Murata, and H. Furutani.
1973. On the artificial fertilization and rearing of larvae in marusoda, *Auxis tapeinosoma*. Kinki Univ. Fac. Agric. Bull. 6:113–116. [In Japanese with English summary.]

Harada, T., O. Murata, and S. Miyashita.
1974. On the artificial fertilization and rearing of larvae in bonito. Kinki Univ. Fac. Agric. Bull. 7:1–4. [In Japanese with English summary.]

Iwai, T., I. Nakamura and K. Matsubara.
1965. Taxonomic study of the tunas. Misaki Mar. Biol. Inst. Special Report 2:1–51. [In Japanese with English summary.]

Kaji, T., M. Tanaka, Y. Takahashi, M. Oka, and N. Ishibashi.
1996. Preliminary observations on development of Pacific bluefin tuna *Thunnus thynnus* (Scombridae) larvae reared in the laboratory, with special reference to the digestive system. Mar. Freshwater Res. 47:261–269.

Kaji, T., M. Tanaka, M. Oka, H. Takeuchi, S. Ohsumi, K. Teruya, and J. Hirokawa.
1999. Growth and morphological development of laboratory-reared yellowfin tuna *Thunnus albacares* larvae and juveniles, with special emphasis on the digestive system. Fish. Sci. 65:700–707.

Kikawa, S.
1953. Spawning of bigeye tuna in the sea near the Marshall Islands. Rep. Southern Sea National Research Institute 24:156–158. [In Japanese.]

Kimura, S., and M. Aritaki.
1985. Studies on rearing and development of larval and juvenile threeline grunt (Pisces: Haemulidae). Bull. Fac. Fish. Mie Univ. 12:193–205.

Kinoshita, I.
1986. Postlarve and juveniles of silver sea bream, *Sparus asrba*, occurring in the surf zones of Tosa Bay, Japan. Japan. J. Ichthyol. 33:7–30.

Kinoshita, I., and S. Fujita.
1988. Larvae and juvenile of blue drum, *Nibea mitsukurii*, occurring in the surf zones of Tosa Bay, Japan. Japan. J. Ichthyol. 35:25–30.

Kohno, H., T. Hoshino, F. Yasuda, and Y. Taki.
1982. Larval melanophore patterns of *Thunnus alalunga* and *T. thynnus* from the Mediterranean. Japan. J. Ichthyol. 28:461–465.

Kohno, H., R. Ordonio-Aguilar, A. Ohno, and Y. Taki.
1996. Osteological development of the feeding apparatus in early stage larvae of the seabass, *Lates calcarifer*. Ichthyol. Res. 43:1–9.

Kohno, H., M. Shimizu, and Y. Nose.
1984. Morphological aspects of the development of swimming and feeding functions in larval *Scomber japonicus*. Bull. Japan. Soc. Sci. Fish. 50:1125–1137.

Kumai, H.
1998. Studies on bluefin tuna artificial hatching, rearing and reproduction. Nippon Suisan Gakkaishi 64:601–605.

Matsumoto, W. M., E. H. Ahlstorm, S. Jones, W. L. Klawe, W. J. Richards, and S. Ueyanagi.
1972. On the clarification of larval tuna identification particularly in the genus *Thunnus*. Fish. Bull. 70:1–12.

Miyashita, S., K. Kato, Y. Sawada, O. Murata, Y. Ishitami, K. Shimizu, S. Yamamoto, and H. Kumai.
1998. Development of digestive system and digestive enzyme activities of larval and juvenile bluefin tuna, *Thunnus thynnus*, reared in the laboratory. Suisanzoshoku 46:111–120.

Miyashita, S., Y. Sawada, N. Hattori, H. Nakatsukasa, T. Okada, O. Murata, and H. Kumai.
In press. Mortality of northern bluefin tuna (*Thunnus thynnus*) due to trauma caused by collision during growout culture. J. World Aqua. Soc. 31.

Miyashita, S., Y. Tanaka, Y. Sawada, O. Murata, N. Hattori, K. Takii, Y. Mukai, and H. Kumai.
2000. Embryonic development and effects of water temperature on hatching of the bluefin tuna, *Thunnus thynnus*. Suisanzoshoku 48:199–207.

Mori, K., S. Ueyanagi, and Y. Nishikawa.
1971. The development of artificially fertilized and reared

larvae of yellowfin tuna, *Thunnus albacares*. Bull. Far Seas Fish. Res. Lab. 5:219–232. [In Japanese with English summary.]

NIH (National Institutes of Health).
1997. NIH image analysis software. NIH, Bethesda, MD, 104 p.

Nishikawa, Y.
1985. Identification for larvae of three species of genus *Thunnus* by melanophore patterns. Bull. Far Seas Fish. Res. Lab. 22:119–129. [In Japanese with English summary.]

Orange, C. J., and B. D. Fink.
1963. Migration of a tagged bluefin tuna across the Pacific Ocean. Calif. Fish and Game 49:307–308.

Podoa, E.
1956. Uova, larve e stadi giovanili di Teleostei; Scombriformes. Fauna e Flora del Golfo di Napoli, Monogr. 38:471–507.

Richards, W. J., and G. R. Dove.
1971. Internal development of young tunas of the genera *Katsuwonus*, *Euthynnus*, *Auxis*, and *Thunnus*. Copeia 1:72–78.

Richards, W. J., and T. Potthoff.
1974. Analysis of the taxonomic characters of young scombrid fishes, genus *Thunnus*. *In* The early life history of fish (J. H. S. Blaxter, ed.), p. 623–648. Springer-Verlag, Berlin, Heidelberg, New York.

Sawada, Y., S. Miyashita, M. Aoyama, M. Kurata, Y. Mukai, T. Okada, O. Murata, and H. Kumai.
2000. Rotifer size selectivity and optimal feeding density of bluefin tuna, *Thunnus thunnus*, larvae. Suisanzoshoku 48:169–177.

Shirota, A.
1970. Studies on the mouth size of fish larvae. Bull. Jap. Soc. Sci. Fish. 36:353–368. [In Japanese with English summary.]

Takii, K., S. Miyashita, M. Seoka, Y. Tanaka, Y. Kubo, and H. Kumai.
1997. Changes in chemical contents and enzyme activities during embryonic development of bluefin tuna. Fisheries Sci. 63:1014–1018.

Ueyanagi, S.
1966. On the red pigmentation of larval tuna and its usefulness in species identification. Rep. Nankai. Reg. Fish. Res. Lab. 24:41–48. [In Japanese with English summary.]

Watanabe, T.
1970. Morphology and ecology of early stages of life in Japanese common mackerel, *Scomber japonicus* Houttuyn, with special reference to fluctuation of population. Bull. Tokai Reg. Fish. Res. Lab. 62:1–259. [In Japanese with English summary.]

Yabe, H., S. Ueyanagi, and H. Watanabe.
1966. Studies on the early life history of bluefin tuna *Thunnus thynnus* and on the larva of southern bluefin tuna *T. maccoyii*. Rep. Nankai Reg. Fish. Res. Lab. 23:95–129. [In Japanese with English summary.]

Yasutake H., G. Nishi, and K. Mori.
1973. Artificial fertilization and rearing of bigeye tuna (*Thunnus obesus*) on board, with morphological observations on embryonic through to early post-larval stage. Bull. Far Seas Fish. Res. Lab. 8, June:71–78. [In Japanese with English summary.]

Yoshida, H. O., and T. Otsu.
1963. Synopsis of biological data on albacore *Thunnus germo* (Lacépède) 1880 (Pacific and Indian Oceans). FAO Fish. Rep. 2(6):274–318.

# Age and growth of red snapper, *Lutjanus campechanus*, from an artificial reef area off Alabama in the northern Gulf of Mexico

**William F. Patterson III**
Coastal Fisheries Institute
204 Wetlands Resources Building
Louisiana State University
Baton Rouge, Louisiana 70803
E-mail address: wpatte2@LSU.edu

**James H. Cowan Jr**
Department of Marine Sciences
University of South Alabama
Mobile, Alabama 36688

**Charles A. Wilson**
Department of Oceanography and Coastal Studies
Louisiana State University
Baton Rouge, Louisiana 70803-7503

**Robert L. Shipp**
Department of Marine Sciences
University of South Alabama
Mobile, Alabama 36688;

**Abstract**—Age and growth were examined of red snapper, *Lutjanus campechanus*, captured in an extensive (3100 km$^2$) artificial reef area off Alabama in the northern Gulf of Mexico. Sagittal otoliths were removed from individuals ($n$=1755) sampled from recreational catches and tournament landings. Marginal increment analysis of sectioned otoliths revealed that a single opaque zone formed annually in sagittae from January through May. Fish ages were estimated from the number of opaque zones in otoliths, timing of opaque zone formation, sampling date, and a presumed birthdate of 1 July. Estimated growth of recaptured red snapper ($n$=288) from a tagging experiment was similar to growth estimated from otolith-aged fish and corroborated otolith aging methods. The von Bertalanffy growth function fitted to length-at-age data was $TL = 969$ mm $(1-e^{-0.192(t-0.020)})$ ($P<0.001$; $r^2=0.99$), which was similar to reported growth functions for western Gulf of Mexico and Atlantic red snapper. Results of our study are consistent with the single stock hypothesis for Gulf of Mexico red snapper.

Manuscript accepted 15 March 2001.
Fish. Bull. 99:617–627 (2001).

Red snapper, *Lutjanus campechanus*, are large, predatory reef fish belonging to the family Lutjanidae. They generally are found from Cape Hatteras, North Carolina, to the Yucatan Peninsula, including the waters of the Gulf of Mexico (GOM) but not the Caribbean Sea (Hoese and Moore, 1998). Throughout their range red snapper are distributed along the continental shelf out to the shelf's edge and demonstrate affinity for vertical structure. Adults aggregate on or near coral reefs, gravel bottoms, or rock outcrops, as well as on artificial reefs, oil rigs, and wrecks (Moseley, 1966; Szedlmayer and Shipp, 1994; Stanley and Wilson, 1997).

Red snapper support economically valuable recreational and commercial fisheries in U.S. waters of the GOM (GMFMC, 1989). Federal management of GOM red snapper is based on the assumption that fish from Florida to Texas constitute a single stock. Although genetic evidence supports this assumption (Camper et al., 1993; Gold et al., 1997; Heist and Gold, 2000), fish are not distributed uniformly across the northern GOM. Fisheries-dependent data suggest there is a center of red snapper abundance off southwest Louisiana and a second, smaller center of abundance off Alabama (Goodyear, 1995a; Schirripa and Legault, 1999). For example, from 1981 to 1998 estimated Louisiana landings of red snapper (commercial and recreational catch from state and federal waters) accounted for 32.6% (mean) ±1.5% (SE) of the total GOM harvest, whereas Alabama landings accounted for 11.4% ±0.9% of the total catch (Schirripa and Legault, 1999).

Although fewer red snapper are harvested from waters off Alabama than from the northwestern GOM, the red snapper fishery off Alabama is unique in several ways. Given that Alabama's GOM coastline represents only about 3.0% of the coastline from Tampa, Florida, to Brownsville, Texas, a disproportionately high percentage of the GOM red snapper harvest is caught and land-




**Figure 1**
Map of artificial reef permit areas off Alabama. Date of creation is given for each area.

ed there. The productivity of the red snapper fishery off Alabama occurs despite the fact that few high-relief (>1 m) natural reefs exist on the continental shelf in the north central GOM (Parker et al., 1983; Shultz et al., 1987; Schroeder et al., 1989); however, off the coast of Alabama exists a 3100-km$^2$ area designated for artificial reef deployment (Fig. 1). The correlation between high catch rates and the creation of artificial reefs off Alabama has caused some to speculate that artificial reefs have increased the productivity of the GOM red snapper stock in this area (Szedlmayer and Shipp, 1994; Minton and Heath, 1998).

Despite the implied effect of artificial reefs on the red snapper fishery off Alabama, few studies have focused on red snapper in this area. The objective of our study was to estimate growth rates of adult red snapper captured off Alabama with otolith-aging and mark-recapture methods and to compare growth estimates with growth of adult red snapper from the western GOM and the southeastern United States. As a corollary to our primary objective, we also attempted to validate presumed annual growth rings in otoliths with marginal increment analysis and present a comparison of estimated growth of otolith-aged red snapper with estimated growth of tagged individuals.

## Methods

### Otolith aging

Red snapper were sampled from July 1995 to September 1999. All fish were caught over artificial reef sites off Alabama. Red snapper shorter than the legal size limit (380 mm total length [TL] for most of the study) were randomly sampled from undersize fish caught during research cruises to tag red snapper. Fish longer than the legal size limit were either randomly sampled from recreational catches or sampled opportunistically at spearfishing or hook-and-line fishing tournaments. Total length and fork length (FL) were measured to the nearest mm for all fish, and whole weight was measured to the nearest 0.1 g. Sex was determined for most fish by macroscopic examination of the gonads. Both sagittae were removed from each sampled individual, rinsed of any adhering tissue, and stored in paper coin envelopes until processing.

Otoliths were sectioned in a transverse plane following the methods of Cowan et al. (1995) and were read under transmitted light with either a Micro Design® model 925 microfiche projector or an Optimas® image analysis system (Media Cybernetics, 1999). Otoliths were read independently by two readers. Blind counts of opaque zones of each sectioned otolith were made along the ventral margin of the sulcus acousticus from the core to the proximal surface; marginal increments were scored following Beckman et al. (1991) (Table 1, Fig. 2). Otoliths for which counts of opaque zones differed between readers were read a second time. Precision among readers was evaluated with the coefficient of variation (CV) (Chang, 1982), index of precision (D) (Chang, 1982), and average percent error (APE) (Beamish and Fournier, 1981).

Age was estimated from the number of opaque zones in otolith sections, timing of opaque zone formation, assumed birthdate, and sampling date. It was assumed that opaque

**Table 1**

Scale used in determining condition of red snapper otolith marginal increments.

| Margin score | Margin description |
|---|---|
| 1 | Opaque zone begins to form at edge; zone is <1/3 the thickness of the previous opaque zone. |
| 2 | Opaque zone at edge is between 1/3 and 2/3 the thickness of the previous opaque zone. |
| 3 | Opaque zone at edge is >2/3 the thickness of the previous opaque zone. |
| 4 | Translucent zone begins to form at edge; zone is <1/3 the thickness of the previous translucent zone. |
| 5 | Translucent zone at edge is between 1/3 and 2/3 the thickness of the previous translucent zone. |
| 6 | Translucent zone at edge is >2/3 the thickness of the previous translucent zone. |

**Figure 2**

Digital images of (**A**) the proximal view of the left sagitta from a 408-mm-TL female red snapper sampled in August 1996 and (**B**) a thin section of a sagitta from a 683-mm-TL female red snapper sampled in January 1996. White squares on the thin section designate opaque zones ($n$=6); edge score is one.

zones constitute annuli and that annulus formation begins 1 January (see "Results" section). The birthdate for red snapper in the north central GOM was assumed to be 1 July, which follows the convention of Goodyear (1995a) and is based on the peak in red snapper spawning in the north central and northeastern GOM (Collins et al., 1996; Szedlmayer and Conti, 1999). According to these assumptions, young-of-the-year northern GOM red snapper form their first annulus in sagittae beginning in January when fish are approximately 0.5 yr old. Therefore, the opaque zone closest to the otolith core represents only 0.5 yr of life, which was accounted for in the aging algorithm.

Age (in d) was estimated (for most fish) by first subtracting one opaque zone from the total number of opaque zones in a given otolith and multiplying the difference by 365 d. Next, 182 d was added to the product to account for the first 0.5 yr of life. Finally, the day of year (number of days since 1 January) the fish was sampled was added to account for the number of days in the sampling year that the fish was alive. The result was divided by 365 d to estimate age in years. Age was estimated similarly for fish that were sampled in November and December and had already begun forming opaque margins, except two was subtracted from the total number of opaque zones before multiplying by 365 d in order to assign fish to the correct year class. For fish that were sampled in January that did not have an opaque margin, zero was subtracted from the total number of opaque zones.

Von Bertalanffy (1938, 1957) growth functions (VBGFs) were fitted to TL-at-age data with Proc NLIN in SAS (SAS Institute, Inc., 1996). Von Bertalanffy growth functions were fitted separately for males and females and a likelihood ratio test was used to test for difference in growth between sexes (Kimura, 1980; Cerrato, 1990). Von Bertalanffy growth functions also were fitted for the complete data set and for the complete set excluding tournament sampled fish.

Weight-TL relationships were modeled with non-linear regression for females and males following Ricker (1975). The functions were computed with Proc NLIN in SAS (SAS Institute, Inc., 1996). Sex-specific weight-TL relationships were made linear by taking the log of weight and TL, and difference between sexes was tested with an analysis of covariance (ANCOVA) on the log-transformed data (SAS Institute, Inc., 1996). Lastly, a weight-TL nonlinear regression was computed for females and males combined with Proc NLIN in SAS (SAS Institute, Inc., 1996).

## Mark-Recapture

A tagging study of adult red snapper was conducted off Alabama from March 1995 to August 1999. Fish were caught with hook and line over nine artificial reef tagging sites in the Hugh Swingle general permit area for arti-




**Figure 3**

Distribution of total length (TL) of red snapper sampled for age and growth estimation.

ficial reef deployment (labeled 1986 in Fig. 1). Each fish was measured to TL and FL, tagged with a yellow Floy® (Seattle, WA) internal anchor tag, and released. Individuals were recaptured on subsequent tagging trips or were recovered by recreational and commercial fishermen. Red snapper recaptured on tagging trips were measured to TL and FL and recovered fish were measured when fishermen retained carcasses of tagged fish.

To corroborate red snapper age estimates, VBGF parameters estimated from otolith-aged fish were incorporated into Fabens' (1965) length increment model to predict TL at recapture of tagged fish (Labelle et al., 1993; Thompson et al., 1999). Solving for TL at recapture, Fabens' model is computed as

$$TLr_i = Tl_i + (L_\infty - t_i)\,(1 - e^{-k\Delta t_i}), \quad (1)$$

where $TLr_i$ = predicted $TL$ at recapture of individual $i$;
$TL_i$ = $TL$ at release of individual $i$;
$L_\infty$ = $TL$ asymptote from VBGF estimated from otolith-aged fish;
$K$ = growth coefficient from VBGF estimated from otolith-aged fish; and
$t_i$ = time at liberty of individual $i$.

Predicted TL at recapture from Fabens' method was plotted against observed TL at recapture to compare growth model predictions to observed values.

Growth of tagged red snapper was estimated by regressing their change in TL on days at liberty (SAS Institute, Inc., 1996). A linear regression also was computed on TL-at-age data over the size range of recaptured individuals from the tagging study (SAS Institute, Inc., 1996). Slopes of resultant regressions were compared to assess if estimated growth rates were different between tagged and otolith aged fish.

## Results

### Age and growth

Sagittae were collected from 1755 red snapper, including 360 fish shorter than the minimum size limit (380 mm TL) caught on tagging cruises, 289 fish from tournaments, and 1106 fish randomly sampled from recreational catches (Fig. 3). Sex was not determined for 279 fish, of which 61 individuals were immature. Mean TL (±SE) was 518.3 (±5.04) mm for males and 529.5 (±5.96) mm for females. Total length of immature individuals ranged from 208 to 309 mm.

Of the 1755 otoliths sectioned for age determination, reader 1 and reader 2 agreed on the number of opaque zones for 1610 (91.7%) fish after the first reading; opaque zones in 23 otoliths were deemed not interpretable owing to sample preparation. Count disagreement between readers was the following: one opaque zone for 123 fish, two zones for 18 fish, and three zones for four fish. Otoliths were read a second time if reader counts were not in agreement and second otoliths were sectioned and read of the 23 fish for which otoliths were rejected after the first reading. After the second reading, agreement was reached for 1676 (95.6%) otoliths, including all 23 otoliths that were second sections. Disagreement between readers after the second reading was as follows: one opaque zone for 71 fish, two zones for 7 fish, and three zones for one fish; fish for

which agreement was not reached after the second reading were not assigned ages or included in the growth estimation. Precision estimates were 1.25% for APE, 0.90% for CV, and 0.64% for D after the second reading.

A clear pattern exists in the marginal increment scores, demonstrating that one opaque zone is formed annually in winter (Fig. 4). Most otoliths had opaque margins by January and had translucent margins by June, thus, timing of opaque zone formation appears to be from January through May for most fish. Eight fish (of 118) sampled in November (1997) and six fish (of 66) sampled in December (1996) had opaque margins. Four fish (of 47) sampled in January (1997) had translucent margins.




**Figure 4**

Plot of monthly margin edge scores of red snapper sagittae. Symbol size is relative to the percentage of fish with the corresponding margin edge score. Monthly sample size is given.

The oldest female sampled was 34.1 yr old and the oldest male sampled was 33.2 yr old. The female to male ratio was 1.1:1 overall but was 1.5:1 for fish between 10 and 20 yr old and 3.4:1 for fish greater than 20 yr old (Fig 5, A and B). Von Bertalanffy growth functions for females and males modeled separately were

$$\text{Females: } TL = 976(1-e^{-0.191(t-0.051)}) \quad (F_{3;735} = 21{,}952;\ P<0.001;\ r^2 = 0.99)$$

$$\text{Males: } TL = 956(1-e^{-0.194(t+0.054)}) \quad (F_{3;726} = 19{,}058;\ P<0.001;\ r^2 = 0.99).$$

Computed VBGFs were not significantly different between males and females (likelihood ratio test; $P>0^2_{df=1}=0.2879$), therefore, sexes were modeled together. Von Bertalanffy growth functions computed for all fish and excluding tournament sampled fish were

$$\text{All fish: } TL = 969(1-e^{-0.192(t-0.020)}) \quad (F_{3;1,672}=47{,}690;\ P<0.001;\ r^2=0.99) \text{ (Fig. 6)}$$

$$\text{Excluding tournament fish: } TL = 1181(1-e^{-0.120(t+0.652)}) \quad (F_{3;1,461}=43{,}550;\ P<0.001;\ r^2=0.99).$$

The growth model including all fish was similar to other VBGFs estimated for GOM and southeast U.S. Atlantic red snapper (Fig. 7, Table 2).

Weight-TL nonlinear regression models for females and males modeled separately were

$$\text{Females: } Weight = (4.46 \times 10^{-9})TL^{3.18} \quad (F_{1;73}= 42{,}577;\ P<0.001;\ r^2=0.98)$$

$$\text{Males: } Weight = (5.18 \times 10^{-9})TL^{3.16} \quad (F_{1;728}= 43{,}956;\ P<0.001;\ r^2=0.99).$$

Log-transformed weight-TL relationships were not significantly different between males and females (ANCOVA test for equal slopes; $F_{1;1,465}=1.54$, $P=0.2145$; ANCOVA test for equal intercepts; $F_{1;1,465}=1.49$, $P=0.2226$), therefore, sexes were modeled together. The resultant nonlinear regression model was




**Figure 5**

Distributions of age for (**A**) all sampled red snapper and (**B**) all fish greater than 10 yr old. Legend in section A is the same for both distributions.




**Figure 6**

Scatterplot of red snapper TL-at-age data for all fish assigned ages. Plotted line is the von Bertalanffy growth function fitted to the data.

$$Weight = (4.68 \times 10^{-9})TL^{3.17}$$

($F_{1;1,467}$=85,961; $P$<0.001; $r^2$=0.98).

## Mark-Recapture

A total of 2932 adult red snapper were tagged. Total length at recapture was measured for 288 of 519 recaptured individuals. Mean TL (±SE) of recaptures was 344 (±4.1) mm (range: 183–660 mm) at release and 423 (±5.6) mm (range: 253–726 mm) at recapture, and mean time at liberty (±SE) was 334 (±16.3) d (range: 12–1501 d). Predicted TL from Faben's method corresponded well to observed TL at recapture for tagged fish (Fig. 8). The linear regression of change in TL on days at liberty was statistically significant ($F_{1;286}$=631.0, $P$<0.001, $r^2$=0.76) and had a slope of 0.238 mm/d (Fig. 9A). The range in estimated ages of otolith-aged fish for the linear regression of TL on age was 356–2599 d (approximately one to seven years). The model was statistically significant ($F_{1;1,541}$=6309, $P$<0.001, $r^2$=0.80) and had a slope of 0.240 mm/d (Fig. 9B).

## Discussion

Validating the periodicity of opaque zone formation in otoliths as annual is imperative for age and growth studies where otoliths are used as aging structures (Beamish and McFarlane, 1983, 1987). Annual formation of opaque zones in otoliths has been validated for several tropical and subtropical lutjanids (Manooch, 1987; Fowler, 1995; Cappo et al., 2000), and annual formation of opaque zones in red snapper sagittae has been reported from the northwestern GOM and the southeastern U.S. Atlantic (Render, 1995; Manooch and Potts, 1997; Wilson and Nieland, 2001). Our marginal increment analysis of red snapper otoliths demonstrates that opaque zones in adult red snapper sagittae also are formed annually in the north central GOM. The pattern of monthly marginal increment scores reveals that some fish begin opaque zone formation as early as November and some do not have translucent margins until midsummer; however, the general pattern of opaque zone formation is from January through May. Render (1995) and Wilson and Nieland (2001) have reported a similar pattern for adult red snapper in the northwestern GOM, and opaque zone formation in winter–spring has been shown for sagittae of several other teleosts in the northern GOM (Maceina et al., 1987; Beckman et al., 1989, 1990, 1991; Thompson et al., 1999).

Because relatively few old red snapper were sampled in our study, and samples of old fish came mostly from tournaments in summer months, we were unable to validate ages beyond 8 years. Baker (1999) compared age estimates of otolith-aged GOM red snapper with age estimates from radiometric dating of their otolith cores and reported that age estimates from the two methods were highly correlated for fish up to 37 yr old, thus corroborating otolith-based estimates of age for older fish. Therefore, despite lack of annulus validation for older red snapper ages in our study, we are confident that otolith-based age estimates of older individuals are accurate.

### Age and growth

Red snapper are long-lived reef fish; we observed a maximum age of 34 yr for females and 33 yr for males. Other

**Table 2**

Results from previous studies of the age and growth of Gulf of Mexico and southeastern U.S. Atlantic red snapper.

| Study and location | Aging structure | Maximum age (yr) | $L_\infty$ mm (TL) | $K$ | $t_0$ |
|---|---|---|---|---|---|
| Szedlmayer and Shipp (1994); north central GOM | sectioned otoliths | 42 | 1025 | 0.150 | not reported |
| Wilson and Nieland, 2001; northwest GOM | sectioned otoliths | 53 | 938 | 0.175 | –0.530 |
| Nelson and Manooch (1982); GOM | scales and sectioned otoliths | 15 | 941 | 0.170 | –0.10 |
| Nelson and Manooch (1982); southeast U.S. Atlantic | scales and sectioned otoliths | 15 | 975 | 0.160 | 0.00 |
| Manooch and Potts (1997); southeast U.S. Atlantic | sectioned otoliths | 25 | 955 | 0.146 | 0.182 |

authors have reported fish over 40 yr old (Szedlmayer and Shipp, 1994; Render, 1995; Wilson and Nieland, 2001) and the oldest fish aged to date is 53 yr old (Render, 1995; Wilson and Nieland, 2001). Among western Atlantic lutjanids for which maximum age has been reported, GOM red snapper has the greatest longevity (Acosta and Appledoorn, 1992; Manickchand-Heileman and Phillip, 1996; Potts et al., 1998; Hood and Johnson, 1999). Maximum ages of 30+ and 40+ yr have been reported for several species of Pacific lutjanids (reviewed in Rocha-Olivares, 1998).




**Figure 7**

Von Bertalanffy growth functions estimated from TL-at-age data for Gulf of Mexico red snapper. Legend indicates source of each function.

Overall, the numbers of sampled males and females were nearly equal, but females were predominant in samples greater than 10 yr old. A similar pattern was observed in Pacific red snapper, *Lutjanus peru*, where the female-to-male ratio was essentially equal (1.1:1) for fish less than 10 yr old and 2.4:1 for fish older than 10 yr (Rocha-Olivares, 1998, Fig. 6). Rocha-Olivares (1998) concluded from catch curve analyses that differences in sex-specific numbers at age for *L. peru* resulted from males experiencing a higher mortality rate. Differential mortality between sexes is not unusual in lutjanids (Grimes, 1987) and the predominance of female GOM red snapper in older age classes may result from a higher mortality rate for males.

Fish sampled from tournaments were included in growth estimation because few large, old individuals were sampled randomly from the recreational catch. The inclusion of tournament sampled fish could bias growth estimates because tournament anglers target large fish; thus the potential exists for them to catch or spear the fastest growing individuals at a given age (Ottera, 1992; Vaughan and Burton, 1993; Goodyear, 1995b). Without the fish sampled from tournaments, however, the VBGF did not reach an asymptote; therefore growth parameters were poorly estimated (Hirschhorn, 1974). We feel that excluding the tournament fish from growth function estimation introduced far greater bias than including them.

## Mark-Recapture

Comparisons between estimated growth of tagged red snapper and otolith-aged fish corroborate otolith aging methods. Predicted TL of tagged individuals obtained with Fabens' (1965) method and VBGF parameters estimated from otolith-aged fish are coincident with observed TL at recapture because predicted and observed values corresponded well to the line of 1:1 agreement. Although ours




**Figure 8**

Predicted total length (TL) at recapture computed for tagged red snapper ($n$=288) plotted versus observed TL at recapture. Diagonal line is the line of one-to-one agreement between predicted and observed values.

was a graphical rather than a statistical exercise (Labelle et al., 1993; Thompson et al., 1999), results corroborate otolith-based estimates of growth.

Direct comparison of linear growth functions computed for tagged fish and otolith-aged fish provides further support for otolith-based estimates of growth (slopes from the two equations differed by only 0.002 mm/d). Francis (1988) cautioned against direct comparison of age-based growth estimates with growth estimates from tagging data because different information results from the different data types. The linear model fitted to length-at-age data predicts TL of fish for a given age (in d), whereas the linear model fitted to tagging data predicts the increment of growth expected of a fish at liberty for a given number of days. However, because the relationship between TL and age is linear over the range of TL from the tagging data (Szedlmayer and Shipp, 1994), we submit that the slopes can be compared as estimates of mean growth. The slope of the age-based linear model is an estimate of the population growth rate of young fish, whereas the slope of the length-based linear model is an estimate of the mean growth of tagged individuals. That growth estimates for otolith-aged and tagged red snapper are very similar corroborates the aging method with otoliths and indicates that on average tagging did not affect fish growth.

## Red snapper off Alabama

The red snapper fishery off Alabama is unique in several ways. Despite the reported lack of high-relief natural reef structures on the continental shelf off Alabama (Parker et al., 1983; Shultz et al., 1987; Schroeder et al., 1989), fishermen land a disproportionately high percentage of the annual GOM red snapper harvest from Alabama coastal waters (Schirripa and Legault, 1999). Although Alabama red snapper landings have accounted for approximately 11% of total GOM landings from 1981 to 1998, the recreational fishery off Alabama accounted for mean (±SE) 21.1 (±2.25)% of GOM recreational landings from 1981 to 1998 and 26.5 (±1.83)% of recreational landings from 1990 to 1998 (Schirripa and Legault, 1999). Total landings of red snapper from the U.S. GOM averaged $3.5 \times 10^3$ metric tons from 1981 to 1998, with commercial and recreational landings essentially equal; however, 94% of Alabama red snapper landings from 1990 to 1998 came from the recreational sector of the fishery (Schirripa and Legault, 1999).

The red snapper fishery off Alabama is also unique because it is prosecuted almost entirely over artificial reefs (Minton and Heath, 1998; Shipp, 1999). Artificial reef construction began in the 1950s when charter boat operators gained permission from the Marine Resources Division of the Alabama Department of Conservation to place 250 car bodies on the sea floor off Alabama (Hosking and Swingle, 1989; Minton and Heath, 1998). Since the 1950s, artificial reefs have been constructed of a variety of materials including car bodies, liberty ships, army tanks, kitchen appliances, newspaper bins, and most recently, prefabricated concrete structures. Reef building increased dramatically in the 1980s with the creation (by permit) of artificial reef areas that now encompass a total of 3100 km$^2$,

and it is estimated that over 15,000 artificial reefs currently exist off Alabama (Hosking and Swingle, 1989; Szedlmayer and Shipp, 1994; Minton and Health, 1998; Havard[1]).

The correlation of catch rates with artificial reef construction has caused some to conclude that artificial reefs off Alabama have increased the production of red snapper, as opposed to merely aggregating fish from surrounding areas (Szedlmayer and Shipp, 1994; Minton and Health, 1998). Among the evidence cited in support of this conclusion was that red snapper grew faster, attained larger sizes, and lived longer over artificial reefs off Alabama than other places throughout their range (Szedlmayer and Shipp, 1994). More recent data, including those from the current study, indicate that GOM red snapper off Alabama grow similarly to and reach similar sizes as fish from the northwestern GOM and the Atlantic (Render, 1995; Manooch and Potts, 1997; Wilson and Nieland, 2001). It is also apparent that the maximum longevity of fish off Alabama is no greater than that of fish of other locations in the GOM (Szedlmayer and Shipp, 1994; Render,1995; Wilson and Nieland, 2001).

These results have important implications for management of GOM red snapper. That growth of fish off Alabama is similar to growth of fish in the northwestern GOM is consistent with the management paradigm that northern GOM red snapper constitute a single stock (Goodyear, 1995a) and contrary to the hypothesis that fish off Alabama are unique (Szedlmayer and Shipp, 1994). Moreover, if fishing mortality rates are higher off Alabama than other places in the northern GOM but growth is the same, production of red snapper may be lower off Alabama than in other areas. Furthermore, if red snapper recruit to artificial reefs off Alabama from other areas in the northern GOM, Alabama's red snapper fishery may serve as a net sink for stock-specific production. Future research is needed to compare mortality as well as growth of fish from different areas in the northern GOM and to estimate the source of recruits to Alabama's artificial reefs.




**Figure 9**

Linear regressions fitted to (**A**) change in total length (TL) versus days at liberty for recaptured red snapper ($n$=288) from the tagging study and (**B**) TL-at-age data for otolith-aged red snapper in the size range of fish from the tagging study ($n$=1543).

## Acknowledgments

Funding for this project was provided by NOAA through MaRFIN (grant numbers NA57FF0054 and NA87FF0424). We thank Scott Baker, Forrest Davis, Andy Fischer, Walter Ingram, Jessica McCawley, Dave Nieland, and Melissa Woods for help in sampling red snapper and sectioning and reading otoliths. We thank Mike and Ann Thierry for allowing us to tag red snapper onboard their charter fishing vessel *Lady Ann*. We thank volunteer fishermen who helped on tagging cruises and commercial and recreational fishermen who reported recaptured fish. Lastly, we thank Amada Gonzales for collecting data from fishermen who reported recaptured fish.

## Literature cited

Acosta, A., and R. S. Appeldoorn.
1992. Estimation of growth, mortality, and yield per recruit for *Lutjanus synagris* (Linnaeus) in Puerto Rico. Bull. Mar. Sci.. 50:282–291.

Baker, M. S., Jr.
1999. Radiometric age validation of red snapper, *Lutjanus campechanus*, and red drum, *Scianenops ocellatus*, from

[1] Havard, R. 1999. Personal commun. Alabama Department of Conservation, Marine Resources Division, P.O. Box 189, Dauphin Island, AL 36528.

the northern Gulf of Mexico. Master's thesis, Louisiana State Univ., Baton Rouge, LA, 59 p.

Beamish, R. J., and D. A. Fournier.
1981. A method for comparing the precision of a set of age determinations. Can. J. Fish. Aquat. Sci. 38:982–983.

Beamish, R. J., and G. A. McFarlane.
1983. The forgotten requirement for age validation in fisheries biology. Trans. Am. Fish. Soc. 112:735–743.
1987. Current trends in age determination methodology. *In* Age and growth of fish (R. C. Summerfelt and G. E. Hall, eds.), p. 15–42. Iowa State Univ. Press, Ames, IA.

Beckman, D. W., A. L. Stanley, J. H. Render, and C. A. Wilson.
1990. Age and growth of black drum in Louisiana waters of the Gulf of Mexico. Trans. Am. Fish. Soc. 119:537–544.
1991. Age and growth of sheepshead *Archosargus probatocephalus* in Louisiana waters using otoliths. Fish. Bull. 89:1–8.

Beckman, D. W., C. A. Wilson, and A. L. Stanley.
1989. Age and growth of red drum, *Sciaenops ocellatus*, from offshore waters of the northern Gulf of Mexico. Fish. Bull. 87:17–28.

Camper, J. D., R. C. Barber, L. R. Richardson, and J. R. Gold.
1993. Mitochondrial DNA variation among red snapper (*Lutjanus compechanus*) from the Gulf of Mexico. Mol. Mar. Biol. Biotech. 2:154–161.

Cappo, M., P. Eden, S. J. Newman, and S. Robertson.
2000. A new approach to validation of periodicity and timing of otolith opaque zone formation in the otoliths of eleven species of *Lutjanus* from the central Great Barrier Reef. Fish. Bull. 98:474–488.

Cerrato, R. M.
1990. Interpretable statistical tests for growth comparisons using parameters in the von Bertalanffy equation. Can. J. Fish. Aquat. Sci. 47:1416–1426.

Chang, W. B.
1982. A statistical method for evaluating the reproducibility of age determination. Can. J. Fish. Aquat. Sci. 39:1208–1210.

Collins, L. A., A. G. Johnson, and C. P. Kiem.
1996. Spawning and annual fecundity of the red snapper (*Lutjanus campechanus*) from the northeastern Gulf of Mexico. ICLARM Conf. Proc. 48:174–188.

Cowan, J. H., Jr, R. L. Shipp, H. K. Bailey IV, and D. W. Haywick.
1995. Procedure for rapid processing of large otoliths. Trans. Am. Fish. Soc. 124:280–282.

Fabens, A. J.
1965. Properties and fitting of the von Bertalanffy growth curve. Growth 29:265–289.

Fowler, A. J.
1995. Annulus formation in otoliths of coral reef fish—a review. *In* Recent developments in fish otolith research (D. H. Secor, J. M. Dean, and S. E. Campana, eds.), p. 45–63. Univ. South Carolina Press, Columbia, SC.

Francis, R. I. C. C.
1988. Are growth parameters estimated from tagging and age-length data comparable? Can. J. Fish. Aquat. Sci. 45:936–942.

Gold, J. R., F. Sun, and L. R. Richardson.
1997. Population structure of red snapper from the Gulf of Mexico as inferred from analysis of mitochondrial DNA. Trans. Am. Fish. Soc. 123:386–396.

Goodyear, C .P.
1995a. Red snapper in U.S. waters of the Gulf of Mexico. Stock assessment report MIA-95/96-05. Miami Laboratory, Southeast Fisheries Science Center, Miami, FL.
1995b. Mean size at age: an evaluation of sampling strategies using simulated red grouper data. Trans. Am. Fish. Soc. 124:746–755.

Grimes, C. B.
1987. Reproductive biology of Lutjanidae: a review. *In* Tropical snappers and groupers: biology and fisheries management (J. J. Polivina and S. Ralston, eds.), p. 239–294. Westview Press, Inc., Boulder, CO.

GMFMC (Gulf of Mexico Fishery Management Council).
1989. Amendment number 1 to the reef fish fishery management plan. GMFMC, Tampa, FL. 356 p.

Heist, E. J., and J. R. Gold.
2000. DNA microsatellite loci and genetic structure of red snapper in the Gulf of Mexico. Trans. Am. Fish. Soc. 129:469–475.

Hirschhorn, G.
1974. Effect of different age ranges on von Bertalanffy parameters in three fishes and one mollusk of the northeastern Pacific Ocean. *In* The aging of fish (T. Bagenal, ed.), p.192–199. Unwin Brothers, Ltd., London.

Hoese, H. D., and R. H. Moore.
1998. Fishes of the Gulf of Mexico, 2nd ed. Texas A&M Press, College Station, TX, 422 p.

Hood, P. B., and A. K. Johnson.
1999. Age, growth, mortality, and reproduction of vermillion snapper, *Rhomboplites aurorubens*, from the eastern Gulf of Mexico. Fish. Bull. 97:828–841.

Hosking, W., and H. A. Swingle.
1989. Alabama's artificial reefs. MS-AL Sea Grant Publ. 9-019, Mobile, AL, 6 p.

Kimura, D. K.
1980. Likelihood methods for the von Bertalanffy growth curve. Fish. Bull. 77:765–776.

Labelle, M., J. Hampton, K. Bailey, T. Murray, D. A. Fournier, and J. R. Sibert.
1993. Determination of age and growth of south Pacific albacore (*Thunnus alalunga*) using three methodologies. Fish. Bull. 91:649–663.

Maceina, M. J., D. N. Hata, T.L. Linton, and A. M. Landry Jr.
1987. Age and growth analysis of spotted seatrout from Galveston Bay, Texas. Trans. Am. Fish. Soc. 116:54–59.

Manickchand-Heileman, S. C., and D. A. T. Phillip.
1996. Reproduction, age, and growth of Caribbean red snapper (*Lutjanus purpureus*) in waters off Trinidad and Tobago. ICLARM Conf. Proc. 48:137–149

Manooch, C. S., III.
1987. Age and growth of snappers and groupers. *In* Tropical snappers and groupers: biology and fisheries management (J. J. Polivina and S. Ralston, eds.), p. 329–373. Westview Press, Inc., Boulder, CO.

Manooch, C. S., III, and J. C. Potts.
1997. Age and growth of red snapper, *Lutjanus campechanus*, Lutjanidae, collected along the southeastern United States from North Carolina through the east coast of Florida. J. Elisha Mitchell Sci. Soc. 113:111–122.

Media Cybernetics.
1999. Optimas, version 6.5. Media Cybernetics, Silver Spring, MD.

Minton, R. V., and S. R. Health.
1998. Alabama's artificial reef program: building oases in the desert. Gulf of Mex. Sci. 16:105–106.

Moseley, F. N.
1966. Biology of the red snapper, *Lutjanus aya* Bloch, of the northwestern Gulf of Mexico. Publ. Inst., Mar. Sci. Univ. Texas 11:90–101.

Nelson, R. S., and C. S. Manooch.
1982. Growth and mortality of red snappers in the west-central Atlantic ocean and northern Gulf of Mexico. Trans. Am. Fish. Soc. 111:465–475.

Ottera, H.
1992. Bias in calculating growth rates in cod (*Gadus morhua* L.) due to size selective growth and mortality. J. Fish Biol. 40:465–467.
Parker, R. O., D. R. Colby, and T. D. Willis.
1983. Estimated amount of reef habitat on a portion of the U.S. south Atlantic and Gulf of Mexico continental shelf. Bull. Mar. Sci. 33:67–86.
Potts, J. C., C. S. Manooch III, and D. S. Vaughan.
1998. Age and growth of vermillion snapper from the southeastern United States. Trans. Am. Fish. Soc. 127:787–795.
Render, J. H.
1995. The life history (age, growth, and reproduction) of red snapper (*Lutjanus campechanus*) and its affinity for oil and gas platforms. Ph. D. diss., Louisiana State Univ., Baton Rouge, LA, 76 p.
Ricker, W. E.
1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can., Ottawa, 382 p.
Rocha-Olivares, A.
1998. Age, growth, mortality, and population characteristics of the Pacific red snapper, *Lutjanus peru*, off the southeast coast of Baja California, Mexico. Fish. Bull. 96:562–574.
SAS Institute, Inc.
1996. Statistics, version 6.11. SAS Institute Inc., Box 8000, Cary, NC.
Schirripa, M. J., and C. M. Legault.
1999. Status of red snapper in U.S. waters of the Gulf of Mexico updated through 1998. Report SFD-99/00-75, Sustainable Fisheries Division, Miami Laboratory, National Marine Fisheries Service, Miami, FL.
Schroeder, W. W., A. W. Shultz, and J. J. Dindo.
1989. Inner-shelf hardbottom areas, northeastern Gulf of Mexico. Trans. Gulf Coast Asso. Geol. Soc. 38:535–541.
Schultz, A. W., W. W. Schroeder, and J. R. Abston.
1987. Along-shore and offshore variations in Alabama inner-shelf sediments. *In* Coastal sediments and processes, 9th symposium of coastal sedimentology, p. 141–152. Florida Sate Univ., Tallahassee, FL.
Shipp, R. L.
1999. The artificial reef debate: are we asking the right questions? Gulf of Mex. Sci. 17:51–55.
Stanley, D. R., and C. A. Wilson.
1997. Seasonal and spatial variations in the abundance and size distribution of fishes associated with a petroleum platform in the northern Gulf of Mexico. Can. J. Fish. Aquat. Sci. 54:1166–1176.
Szedlmayer, S. T., and J. Conti.
1999. Nursery habitats, growth rates, and seasonality of age-0 red snapper, *Lutjanus campechanus*, in the northeast Gulf of Mexico. Fish. Bull. 97:626–635.
Szedlmayer, S. T., and R. L. Shipp.
1994. Movement and growth of red snapper, *Lutjanus campechanus*, from an artificial reef area in the northeastern Gulf of Mexico. Bull. Mar. Sci. 55:887–896.
Thompson, B. A., M. Beasley, and C. A. Wilson.
1999. Age distribution and growth of greater amberjack, *Seriola dumerili*, from the north-central Gulf of Mexico. Fish. Bull. 97:362–371.
Vaughan, D. S., and M. L. Burton.
1993. Estimation of von Bertalanffy growth parameters in the presence of size-selective mortality: a simulated example with red grouper. Trans. Am. Fish. Soc. 123:1–8.
von Bertalanffy, L.
1938. A quantitative theory of organic growth. II. Inquiries on growth laws. Human Biol. 10:181–213.
1957. Quantitative laws in metabolism and growth. Quart. Rev. Biol. 32:217–231.
Wilson, C. A., and D. L. Nieland.
2001. Age and growth of red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico off Louisiana. Fish Bull. 99:653–664.

# A comparison of calcified structures for aging summer flounder, *Paralichthys dentatus**

**Ann M. Sipe**
Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, Virginia 23062
E-mail address: amsipe@vims.edu

**Mark E. Chittenden Jr.**
Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, Virginia 23062

**Abstract**—Calcified structures of summer flounder, *Paralichthys dentatus*, were evaluated to identify the best age determination method. Scales, the currently preferred structure, were compared with opercular bones and to right and left whole and sectioned otoliths for ages 0 to 10. All structures showed concentric rings that were interpreted as annual; however structures differed greatly in the clarity of their presumed annual marks. Right and left otoliths generally gave the same age, although they differed in the clarity of marks. Sectioned otoliths, particularly right ones, were the best aging structure. Right sectioned otoliths consistently showed the clearest marks and had the highest confidence scores, lowest reading times, and highest agreement within and between readers, 97% and 96%, respectively. Left sectioned otoliths took twice as long to prepare and were more difficult to interpret than right sectioned otoliths. Whole otoliths were the second best structure and were adequate to age 4 or 5, after which sectioning greatly improved the clarity of marks. Scales were inferior to, and often did not give the same age readings as, whole and sectioned otoliths. Compared with otoliths, scales tended to overage at younger ages and to underage at older ages. Opercular bones were undesirable for aging summer flounder. They were often unclear and inconsistent, and they had the lowest confidence scores, the highest reading times, and only 46% within-reader agreement. A major source of disagreement in scale and otolith age readings was the presence of an early, presumably false, mark on some structures. We compare the formation of this early mark in summer flounder with early mark formation on otoliths of Atlantic croaker, a species with similar life history traits.

Manuscript accepted 21 March 2001.
Fish. Bull. 99:628–640 (2001).

The summer flounder, *Paralichthys dentatus*, ranges from Nova Scotia to Florida, although it is most abundant from Massachusetts to North Carolina (Ginsburg, 1952; Leim and Scott, 1966; Gutherz, 1967). In regions of high abundance, it is one of the most important commercial and recreational fishes on the Atlantic coast (MAFMC, 1987). In the Chesapeake Bay region, for example, summer flounder support an extensive recreational fishery from about March to November, when they are present in the lower portions of the Chesapeake Bay and in coastal waters (Hildebrand and Schroeder, 1928; MAFMC, 1987; Desfosse, 1995). They then support a strong commercial fishery during the fall and winter, when they move offshore to the continental shelf (Ginsburg, 1952; Bigelow and Schroeder, 1953; Poole, 1962; MAFMC, 1987).

Many studies have reported difficulties with the structures used for age determination of summer flounder. Prior to about 1980, whole left otoliths were the most commonly used structure (Poole, 1961; Eldridge, 1962; Smith and Daiber, 1977; Powell, 1982). However, there were disagreements over the location and interpretation of the first presumed annual mark (Poole, 1961; Eldridge, 1962; Smith and Daiber, 1977), largely a result of uncertainties about first year growth rates. This and other problems with whole otoliths (summarized in Smith et al., 1981) prompted a comparison of age determination structures by Shepherd (1980), who reported that presumed annual marks were more distinct on scales than on whole otoliths. Consequently, scales became the preferred structure for aging summer flounder (Smith et al., 1981; Dery, 1988; Almeida et al., 1992). More recently, Szedlmayer et al. (1992) examined first year growth rates to resolve the location and interpretation of the first mark on whole otoliths, but scales have remained the preferred structure (Bolz et al., 2000).

Difficulties have also been reported in using summer flounder scales (Dery, 1988; Desfosse, 1995; Bolz et al., 2000). Desfosse (1995) used marginal increment analysis to validate scales for ages 1 to 3. He reported only 46% within-reader agreement past age 4, however, indicating that marks on scales are not very distinct at older ages. He attributed disagreements to false or indistinct annuli and to crowding of annuli at the scale edge in older fish. Most recently, Bolz et al. (2000) reported only 53% agreement for ages 1 to 5 in a between-agency exchange of scales, with agreement increasing to only 83% after they resolved as many disagreements as possible. They attributed most of the remaining disagreements to the choice of a first annual mark and to differing opinions on what constituted a false mark on scales.

A reexamination of calcified structures for aging summer flounder is needed, given their economic importance and the reported difficulties in

* Contribution 2380 from the Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, VA 23062

age determination with whole otoliths and scales. Previous studies have never evaluated sectioned otoliths in summer flounder, even though sectioned otoliths have often proven a superior structure in other species, especially at older ages when scales and other structures can underage fish (Beamish and McFarlane, 1983). Further study is especially needed because the location of the first mark on otoliths has recently been determined (Szedlmayer et al., 1992). In addition, no work has been done to determine if right-left differences in the location of the focus result in differences in age determination.

The main objective of our study was therefore to evaluate and compare whole otoliths, sectioned otoliths, scales, and opercular bones for aging summer flounder. We included opercular bones because many studies, on a variety of species, have found them to be superior to other structures and to have very distinct and easy to read marks (for examples, see LeCren, 1947; Donald et al., 1992; Hostetter and Munroe, 1993). A second objective was to compare right and left otoliths for potential differences in age based on differences in the location of the focus. Calcified structures were evaluated in terms of preparation and reading times, confidence in presumed annual mark clarity, agreement between repeated age readings, structure growth with fish growth, age agreement between different structures of the same fish, and increases in the number of presumed annual marks with structure size and fish size. Finally, we discuss the formation of early, presumably false, marks on summer flounder otoliths and scales that resulted in difficulties in age interpretation

## Methods

### Sample collection

To minimize difficulties interpreting marks on the edge of the structures, collections of summer flounder were made far from the time of presumed annual mark formation, which occurs in May and June on the scales of Chesapeake Bay summer flounder (Desfosse, 1995). Summer flounder were collected from commercial fisheries in the Chesapeake Bay region from September through November of 1998 ($n$=165). Additional juvenile fish ($n$=11) were collected by the Virginia Institute of Marine Science juvenile bottom trawl survey in October of 1998 in the lower Chesapeake Bay and James River.

Fish were processed for total length (TL), total weight (TW), and sex, and the calcified structures were removed as follows. Both saggital otoliths were removed, wiped clean, and stored dry in tissue culture cell wells. Scales were removed from just above the lateral line anterior to the caudal peduncle (Shepherd, 1980; Dery, 1988) and stored in coin envelopes. Both opercular bones were removed according to the methods of LeCren (1947), stored in coin envelopes, and frozen.

The collection of summer flounder was stratified into six length-based categories of 100 mm each to include as many age groups as possible in the final study sample. A random sample of 15 fish was then chosen from the first five categories. The last category included the six largest fish, all of which were used in the comparison, for a total of 81 fish. All calcified structures in the final study sample were assigned random numbers before preparation and aging. Summer flounder in the final study sample ranged in size from 209 to 758 mm TL and from 80.8 to 7304.6 g TW and in age from 0 to 10 years (determined from sectioned otoliths, as reported in this study).

### Preparation of calcified structures for age determination

Whole otoliths were examined in water on a dark background with reflected light at 120 to 240× magnification. Thin opaque bands, which appeared white under reflected light, were presumed to represent annual marks (Fig. 1A). Two counting paths were used for mark enumeration. The primary counting path was from the focus to the anterior margin of the otolith. The secondary counting path, used to verify the primary counting path reading, was from the focus to the posterior margin of the otolith. With calipers to 0.05 mm, whole otolith total length (WOTL) was measured as the largest distance from the anterior to the posterior edge and whole otolith radial length (WORL) was measured from the center of the focus to the tip of the anterior edge. A paired sample $t$-test was used to test for right-left differences in WORL.

After all whole otolith readings were made, right and left otoliths were mounted sulcal groove down onto cardboard with crystal bond adhesive and sectioned transversely through the focus with a variable speed Beuhler Isomet saw. The resulting sections, about 0.5 mm thick, were mounted on clear glass slides and immersed in crystal bond. Sections were viewed with transmitted light and bright field at 240× magnification. Thin opaque bands, which appeared dark with transmitted light, were presumed to represent annual marks (Fig. 1B) and were counted along the ventral side of the sulcal groove. Sectioned otolith radial length (SORL) was measured to 0.001 mm along the ventral arm of the sulcal groove from the center of the focus to the otolith edge by using a compound video microscope with the Optimas image analysis system (Media Cybernetics, 1999). Broken otoliths were not measured if they were fractured along the focus. A paired sample $t$-test was used to test for right-left differences in SORL.

Opercular bones were prepared according to the methods of LeCren (1947). Briefly, they were soaked in cold tap water for several minutes to thaw and to partially loosen surrounding skin, then soaked for 1 minute in simmering water, after which the skin was easily removed with a toothbrush. The opercular bones were then rinsed with cold tap water and air-dried. Opercular bones were examined dry with transmitted light and in water with reflected light on a dark background. Presumed annual marks (Fig. 1C) were defined as sharp transitions from relatively narrow translucent zones to relatively wide opaque zones that were continuous from the anterior to the posterior margin of the bone (Bagenal and Tesch, 1978; Hostetter and Munroe, 1993). Translucent zones appeared white under




**Figure 1**

Marks on calcified structures taken from a 5-year-old (determined from sectioned otoliths) female summer flounder, TL = 687 mm, collected in mid-January. Arrows indicate individual marks counted (as described in the "Methods" section). (**A**) Whole otoliths, viewed with reflected light on a black background. Arrows indicate presumed annual marks along the primary counting path; dots indicate presumed annual marks along the secondary path. Whole otolith radial lengths were measured along the primary counting path. (**B**) Transverse otolith sections, viewed in transmitted light. Sectioned otolith radial lengths were measured along the counting path (indicated by arrows) from the center of the focus to the edge along the ventral arm of the sulcal groove. (**C**) Right opercular bone, viewed with reflected light on a black background. AA = articular apex. (**D**) Scale impressions, viewed in transmitted light. White arrows indicate marks that appear on only one of the scales. Asterisks indicate probable false marks. Both scales have a probable false mark prior to the first mark counted.

transmitted light and dark under reflected light, whereas opaque zones appeared dark under transmitted light and white under reflected light. The first presumed annual mark was defined as the first opaque zone after the first translucent zone, where the first translucent zone occupied the central focal area of the opercular bone. Both bones were examined, and the one with the clearest marks was used for aging. Opercular bone radial length (OpRL) was measured to 0.05 mm from the center of the articular apex to the anterior margin edge with calipers.

Scales were soaked in water until flexible and brushed gently with a soft bristle toothbrush. Then 5 or 6 clean, symmetrical, unregenerated scales were dried, taped to an acetate sheet, inserted between two new acetate sheets, and pressed in a Carver laboratory scale press for 2 minutes at 15,000 pounds of pressure and 60°C. Scale impressions were read with a Bell-Howell R753 microfiche reader at 20× and 32×. Presumed annual marks were identified with standard scale reading criteria as described in Smith et al. (1981), Dery (1988), and Almeida

et al. (1992). Briefly, readers enumerated marks (Fig. 1D) that exhibited "cutting over" in both lateral fields of the scale that was accompanied by a clear narrow zone in the anterior portion of the scale. Scale radial length (ScRL) was measured to 0.001 mm from the center of the focus to the anterior edge of the scale by using a compound video microscope with the Optimas image analysis system (Media Cybernetics, 1999).

### Evaluation of calcified structures

Each structure was examined for age by two readers—twice by reader 1 and once by reader 2. Structures were read in a randomly selected order with no knowledge of fish size or collection date. Ages were assigned on the basis of presumed annual mark counts. Different structures from the same fish were read independently, including right and left otoliths, and at least one week separated the first and second readings of the same structure.

Preliminary evaluations of structures included preparation times, reading times, confidence in the clarity of presumed annual marks, growth of the structures with size of the fish, and agreement in repeated age readings of the same structure (precision). Structures judged acceptable based on those criteria were then evaluated further for agreement in age readings between different structures from the same fish and to see if the number of presumed annual marks increased with structure size and fish size. Our preliminary evaluation indicated otoliths and scales to be superior to opercular bones; therefore opercular bones were not evaluated further.

Preparation time, a measure of the processing efficiency of a structure, was evaluated as the time taken to prepare structures for reading. Clarity of presumed annual marks on a structure was evaluated using both reading times and confidence scores. Reading time was measured as the time taken to read a given structure in an individual fish. Confidence scores, expressed on a scale of 1 (low) to 5 (high), were assigned by the reader to each reading based on the clarity of the marks. Differences in confidence scores between structures were tested at $\alpha = 0.05$ by using the normal approximation to the Mann-Whitney test for ordinal data (Zar, 1996).

The assumption that structure growth is directly related to fish growth was evaluated using regression analysis (Zar, 1996). Structure sizes (ScRL, OpRL, WOTL, WORL, SORL) were regressed on fish TL to determine if the relationships were significant and increasing. Sample sizes varied in these regressions, and in regressions of the number of presumed annual marks on structure size described below because some structures were broken in preparation and could not be measured.

Precision in age determinations for a given structure was evaluated using simple percent agreement in repeated readings within and between readers. Within-reader agreement compared the first and second readings by reader one, and between-reader agreement compared the first readings of each of the two readers. Reader comments on structure features were evaluated to determine the proximal causes of disagreements.

Scales that disagreed in the initial two readings by reader 1 were reread independently a third time by reader 1 to reach a consensus for use in between-structure comparisons. Likewise, right and left otoliths that disagreed in the initial two readings by reader 1 were read a third time to reach a consensus. Structures that showed no agreement in three readings (1 of 81 for scales, 1 of 81 for sectioned otoliths) were not included in between-structure comparisons.

Agreement in presumed annual mark counts between different structures of an individual fish was evaluated by using simple percent agreement between structures and simple linear regression procedures. For the regressions, ages determined by one structure were regressed on ages determined by another structure, and the slope of the regression line was tested to see if it differed significantly from one. A slope of one implies that $y = x$ and that the two structures give the same age. For each regression, we used as the $x$-variable the structure judged to be superior in the preliminary evaluations.

The assumption that the number of presumed annual marks on a structure is directly related to structure size and to fish size was evaluated using regression analysis (Zar, 1996). The number of presumed annual marks on a structure was regressed on structure size (ScRL, WOTL, WORL, SORL) and on fish TL to determine if the relationships were significant and increasing.

## Results

### Comparative appearance of calcified structures

All four calcified structures showed concentric marks that were interpreted as annual (Fig. 1). However, these structures differed greatly in the clarity of presumed annual marks.

Presumed annual marks on both whole and sectioned otoliths (Fig. 1, A and B) were typically clear, consistent, and easy to interpret, especially for sectioned otoliths. The right-left difference in the location of the focus had moderate effects on mark clarity for both whole and sectioned otoliths, as described below. Whole otolith marks were most easily read at younger ages, but age had little effect on sectioned otolith mark clarity. The few disagreements in otolith ages were primarily caused by an early, presumably false, mark that often occurred prior to the first presumed annual mark (Fig. 2). This early mark appeared as a thin opaque band close to, but distinct from, the focus and was found on both young (Fig. 2A) and older (Fig. 2B) fish. We tried not to count this early mark in our age readings, because it did not occur consistently in all fish. Finally, only one otolith of 81 pairs was poorly calcified and unable to be read whole, although its age was easily determined upon sectioning.

Presumed annual marks on opercular bones (Fig. 1C) were fairly clear in some fish, but they were more often poorly defined, inconsistent, and difficult to follow across the structure, making age interpretation difficult and highly subjective. Opercular bones commonly exhibited un-

clear transitions from translucent to opaque zones; the first one or two marks were particularly difficult to distinguish, even on young fish. Zone transitions were often easier to interpret towards the edge of the structure in older fish, although this too varied greatly from fish to fish. The example in Figure 1C is unusually clear and easy to read.

Presumed annual marks on scales (Fig. 1D) were clearer than those on opercular bones, but they still required much subjective interpretation. Figure 1D shows some of the common problems encountered with scales, including presumably false marks (asterisks) and marks that were present on only some scales from the same fish (white arrows). In addition, many fish had regenerated, asymmetrical, or otherwise damaged scales, making it difficult and time-consuming to choose acceptable scales to press. For example, about 20 scales were pressed in order to obtain two scales that were adequate to show in Figure 1D. Interpretation of age from scales of older fish was extremely difficult because marks at the scale edges were often obscured or crowded together, particularly in the narrow lateral fields. Finally, a major source of disagreement in age determination from scales resulted from an early, presumably false, mark that often occurred prior to the first presumed annual mark (Fig. 1D, asterisk). Because this early mark did not appear consistently in all fish or even on several scales from the same fish, we tried not to count it in our age readings.




**Figure 2**

Right whole otoliths showing an early, presumably false, mark. (**A**) is from a 299-mm-TL age-1 fish collected in September, and (**B**) is from a 442-mm age-4 fish collected in October. White arrows point to the early marks. Black arrows indicate primary counting path (anterior field), dots indicate secondary counting path (posterior field).

### Preparation times, reading times, and confidence in clarity of marks

Preparation times were short and reasonable for all structures, at less than 15 minutes per fish. Whole otoliths took by far the shortest time because no preparation was required before reading (Table 1). Sectioned right otoliths and opercular bones required 4 to 6 minutes to prepare, whereas scales and sectioned left otoliths took much longer to prepare, about 11 and 14 minutes, respectively. Left sectioned otoliths took much longer to prepare than right sectioned otoliths primarily because they broke much more frequently during sectioning.

Reading times were short and reasonable for all structures, at less than three minutes per fish. Sectioned right otoliths had by far the shortest reading time, at only 0.27 minutes per fish (Table 1). Whole otoliths and sectioned left otoliths had the next shortest reading time, at only about 0.4 to 0.6 minutes per fish. Scales (1.2 min) and opercular bones (2.4 min) both required much more reading time than otoliths, indicating that otoliths could be aged more easily.

Reader confidence scores varied greatly between structures. Sectioned otoliths had by far the highest confidence scores, with values of 4.9 and 4.8 for the right and left, respectively (Table 1). Whole otoliths had somewhat lower confidence scores, with values of 4.1 and 3.8 for the right and left, respectively. Confidence scores were much lower for scales (3.2) and especially for opercular bones (2.3), indicating that these structures were not as easily interpreted. All confidence scores were significantly different from one another ($Z$=2.10 to 4.18; $P$<0.0001 to 0.013; individual values not reported).

### Regression of structure size on fish size

All calcified structures grew in size as summer flounder body length grew, indicating that each structure could be useful for back-calculation studies. All regressions of structure size on total length were significant at $P < 0.001$, and all slopes were positive (Table 2). All regressions were strong and explained much of the variation in structure size, generally 90% or more, with coefficient of determination values (100 $r^2$) ranging from 72% to 98%. Values for 100 $r^2$ were less than 91% only for right and left sectioned otoliths, which were 72% and 85%, respectively.

### Agreement in age determinations for the same structure

Agreement (precision) between repeated age readings varied greatly between calcified structures. Precision by the same reader was highest by far (95% to 97%) for sectioned right and left otoliths and left whole otoliths (Table 3). Precision was somewhat lower in right whole otoliths (89%) than in

**Table 1**

Average preparation times (min), reading times (min) ± standard error (SE), and confidence scores (±SE) for summer flounder calcified structures.

| Structure | Preparation time | Reading time | Confidence score |
|---|---|---|---|
| Opercular bones | 4.63 | 2.43 ±0.20 | 2.31 ±0.16 |
| Scales | 10.50 | 1.20 ±0.13 | 3.21 ±0.15 |
| Sectioned otoliths | | | |
| Right | 5.86 | 0.27 ±0.04 | 4.91 ±0.04 |
| Left | 13.93 | 0.57 ±0.09 | 4.75 ±0.05 |
| Whole otoliths | | | |
| Right | 0.00 | 0.45 ±0.06 | 4.10 ±0.11 |
| Left | 0.00 | 0.41 ±0.04 | 3.84 ±0.10 |

**Table 2**

Regression statistics for relationships between structure size and summer flounder total length (TL). Structure abbreviations are defined in the "Methods" section of the text. $n$ = sample size. All regressions were significant at $P < 0.001$.

| Structure | Equation | $n$ | 100 $r^2$ |
|---|---|---|---|
| Opercular bones | $OpRL = -2.280 + 0.0772\ TL$ | 66 | 98 |
| Scales | $ScRL = -0.348 + 0.0126\ TL$ | 81 | 93 |
| Sectioned otoliths | | | |
| Right | $SORL = -0.015 + 0.0027\ TL$ | 66 | 85 |
| Left | $SORL = 0.015 + 0.0018\ TL$ | 47 | 72 |
| Whole otoliths | | | |
| Right | $WORL = 0.642 + 0.0089\ TL$ | 76 | 91 |
| Left | $WORL = 0.601 + 0.0111\ TL$ | 77 | 93 |
| Right | $WOTL = 1.280 + 0.0164\ TL$ | 76 | 94 |
| Left | $WOTL = 1.530 + 0.0156\ TL$ | 77 | 91 |

**Table 3**

Average percent agreement, within and between readers, for presumed annual mark counts on summer flounder calcified structures.

| Structure | Within reader | Between reader |
|---|---|---|
| Opercular bones | 46 | — |
| Scales | 80 | 58 |
| Sectioned otoliths | | |
| Right | 97 | 96 |
| Left | 95 | 88 |
| Whole otoliths | | |
| Right | 89 | 86 |
| Left | 97 | 87 |

left whole otoliths; however this could be attributed to the reader learning to use reflected lighting more effectively during the second reading, because 7 of the 9 consensus readings for right otoliths agreed with the second reading. Within-reader agreement was lower with the use of scales (80%), but precision varied with age. Agreement in repeated scale readings was actually high for ages 0 to 4 (92%, $n$=52), but it decreased to only 59% for fish over age 4 ($n$=29). Precision was lowest by far in opercular bones (46%), where there were no patterns in agreement by age. Because opercular bones showed the lowest precision and the poorest mark clarity, we did not include them in further evaluations.

Agreement in age determinations between readers also varied greatly among calcified structures. Precision between readers was highest by far (96%) for right sectioned otoliths (Table 3). Agreement was somewhat lower (86% to 88%) for left sectioned otoliths and whole otoliths. Agreement was lowest by far for scales (58%), reflecting the overall poor clarity of marks and the resulting subjectiveness in scale age readings compared with otolith age readings.

## Comparison of right and left otoliths

Differences in right and left radial lengths were observed for both whole and sectioned otoliths. The right radial length was significantly shorter than the left in whole otoliths (paired $t$=17.59, df=73, $P$<0.0001; Fig. 1A). However, for sectioned otoliths, the right radial length was significantly longer than the left (paired $t$ =–11.72, df=43, $P$<0.0001; Fig. 1B) because the right otolith is thicker at the focus, where the transverse cross section was taken.

Right and left whole otoliths generally gave the same age readings. Reader one had high age agreement between right and left whole otolith readings (96%), and the null hypothesis that the slope of the line equals one was not rejected ($P$=0.077, Fig. 3A).

Although right and left whole otoliths generally indicated the same age, they differed in mark clarity. When the posterior field (secondary counting path) was used to verify or determine the number of presumed annual marks, the right otolith was generally much easier to read than the left because of the greater distance between the focus and the posterior margin on the right otolith (Fig. 1A). This greater distance made the marks further apart and more easily distinguishable on the right than on the left otolith. The difference in mark clarity was greatest for older fish and was also reflected in significantly higher confidence scores for the right whole otolith than for the left (Table 1).

Right and left sectioned otoliths also generally gave the same age readings. Reader one had high age agreement between right and left sectioned otolith readings (94%), and the null hypothesis that the slope of the line equals one was not rejected ($P$=0.393, Fig. 3B).

Although right and left sectioned otoliths generally gave the same age, presumed annual marks were usually clearer and easier to interpret on the right otolith. Right sectioned otoliths had a much longer counting path and were therefore easier to age than left sectioned otoliths, where the marks were more crowded and less clearly defined (Fig. 1B). This difference was also reflected in higher confidence scores and lower reading times for the right sectioned otolith than for the left (Table 1).

## Comparison of different calcified structures from the same fish

Whole and sectioned otoliths generally gave the same age readings. The number of presumed annual marks on whole and sectioned otoliths showed high agreement (95%), with 100% agreement for fish under age 4 (Fig. 4A). In addition, the null hypothesis that the slope of the line equals one was not rejected ($P$=0.901).

Although whole and sectioned otoliths generally provided the same age, presumed annual marks were often clearer on sectioned otoliths than on whole ones, especially in older fish, where crowding of marks at the edge of whole otoliths became a problem. This observation is supported by the much higher confidence scores for sectioned otoliths (Table 1). As a specific example, the oldest fish in the comparison showed very clear marks and was aged 10 in every reading using both right and left sectioned otoliths (Fig. 5A), and all confidence scores were 5. Marks were less clear on the whole otolith (Fig. 5B), however, with between 8 and 10 marks counted in different readings, and an average confidence score of only 2.5. In general, the use of sectioned otoliths appeared to greatly increase mark clarity in fish over age 4 or 5.

Scales and sectioned otoliths often did not give the same age readings. Agreement in the number of presumed annual marks on scales and sectioned otoliths was undesirably low, at only 80% (Fig. 4B). In addition, the null hypothesis that the slope of the line equals one was rejected ($P$=0.047). Scales tended to overage compared with sectioned otoliths in fish age 4 and younger, but to underage in fish older than age 4. Agreement between scales and sectioned otoliths was fairly high for ages 0 to 4 (86%, $n$=56) but decreased to only 65% in fish over age 4 ($n$=23).

Scales and whole otoliths often did not give the same age readings. Agreement in the number of presumed annual marks on scales and whole otoliths was also undesirably low, at only 76% (Fig. 6). In addition, the null hypothesis that the slope of the line equals one was again rejected ($P$=0.039). As with sectioned otoliths, scales tended to overage compared with whole otoliths in fish age 4 and younger and to underage in fish older than age 4. Agreement between whole otoliths and scales was fairly high for ages 0 to 4 (85%, $n$=53) but decreased to only 56% in fish over age 4 ($n$=25).




**Figure 3**

Comparisons of presumed annual mark counts on the left otolith with mark counts on the right otolith for whole otoliths and sectioned otoliths in summer flounder. The 45° diagonal line represents 1:1 agreement. The number of fish is indicated at each symbol.

## Increase in number of marks with structure size and fish size

Mark counts on calcified structures increased as structure size and fish size increased, indicating that each structure tested could be useful in age determination. All regressions of mark counts on structure size were significant at $P$<0.001, and all slopes were positive (Table 4). Regressions were generally strong and explained much of the variation in mark counts because 100 $r^2$ values were high, generally from 80% to 86%. Values for 100 $r^2$ were lowest for left sectioned otolith radius and scale radius, at 67% and 73%, respectively. Likewise, all regressions of mark counts on fish size were significant at $P < 0.001$, and all slopes

were positive (Table 5). All regressions were again strong, with 100 $r^2$ values from 83% to 86%.

## Discussion

### Comparative evaluation of sectioned otoliths

Our findings indicate that sectioned otoliths are the best structure for aging summer flounder over the age range 0 to 10 years. Sectioned otoliths had the shortest reading times, the highest confidence scores, the highest within- and between-reader agreement, and they were consistently clearer and easier to read than whole otoliths, scales, and opercular bones. These findings are new for summer flounder because no published studies have used sectioned otoliths to age this species. These findings generally agree, however, with many studies on other species that have found sectioned otoliths to be the best aging structure (for examples, Beamish, 1979; Chilton and Beamish, 1982; Beamish and McFarlane, 1983; Lowerre-Barbieri et al., 1994).

Right sectioned otoliths were generally superior to left sectioned otoliths. Although we found high agreement in age between right and left sectioned otoliths, right otoliths were much easier to prepare, and they had a larger counting path, which made it easier to identify the marks, resulting in shorter reading times, higher confidence scores, and higher reader agreement.

Although we have found sectioned otoliths to be the best structure for determining the age of summer flounder, our studies have not proven their accuracy. To do so would require known-age methods or at least marginal increment methods. However, until validation is done, we feel there is sufficient evidence to recommend that sectioned otoliths replace the current practice of using scales for aging summer flounder.




**Figure 4**

Comparisons of presumed annual mark counts on whole otoliths (**A**) and scales (**B**) with mark counts on sectioned otoliths in summer flounder. The 45° diagonal line represents 1:1 agreement. The number of fish is indicated at each symbol.

### Comparative evaluation of whole otoliths

Our findings indicate that whole otoliths are the second best structure for aging summer flounder over the age range of 0 to 10 years. Whole otoliths had no preparation time and had the second shortest reading times, the second highest confidence scores, the second highest within- and between-reader agreement, and the highest agreement with sectioned otoliths. Whole otoliths were generally easy to read in fish less than age 4 or 5, and we feel they are adequate for these younger ages, especially in large-scale production aging where preparation time is important.

We found that the right whole otolith was often easier to read than the left when the secondary counting path was used. Therefore, although former studies have used the left whole otolith only (Poole, 1961; Eldridge, 1962; Smith and Daiber, 1977; Powell, 1982), we suggest that the right should be included in future work.

Our findings on preparation and reading times, confidence scores, within- and between-reader agreement and agreement with sectioned otoliths are generally new because the literature has not reported detailed evaluations of whole otoliths in summer flounder. Given our findings, we do not agree with the current preference for using scales rather than whole otoliths in summer flounder. Indeed, we disagree with the original reasons for rejecting otoliths, which included 1) poor calcification and poor contrast between opaque and translucent zones (Shepherd, 1980; Smith et al., 1981; Dery, 1988), 2) obscurement of the first mark as the fish ages (Powell, 1982), 3) deviation from the generalized pattern of opaque and translucent zone for-




**Figure 5**

Right sectioned (**A**) and whole (**B**) otolith from a female summer flounder, 10 years old (determined from sectioned otoliths) and 758 mm TL, collected in November. Arrows on the sectioned otolith indicate presumed annual marks. On the whole otolith, arrows indicate primary counting path (anterior field), dots indicate secondary counting path (posterior field). Ten marks are visible in the posterior field of the whole otolith, but only eight marks are visible in the anterior field.

**Table 4**

Regression statistics for relationships between the number of marks (*Marks*) and calcified structure size for summer flounder. Structure abbreviations are defined in the "Methods" section of the text. $n$ = sample size. All regressions were significant at $P < 0.001$.

| Structure | Equation | $n$ | 100 $r^2$ |
|---|---|---|---|
| Scales | $Marks = -2.64 + 1.080\ ScRL$ | 80 | 73 |
| Sectioned otoliths | | | |
| Right | $Marks = -3.39 + 5.424\ SORL$ | 65 | 80 |
| Left | $Marks = -3.36 + 6.996\ SORL$ | 46 | 67 |
| Whole otoliths | | | |
| Right | $Marks = -4.56 + 1.664\ RWOR$ | 75 | 85 |
| Left | $Marks = -4.47 + 1.367\ LWOR$ | 76 | 86 |
| Right | $Marks = -4.80 + 0.919\ RWOT$ | 75 | 86 |
| Left | $Marks = -4.80 + 0.934\ LWOT$ | 76 | 82 |

**Table 5**

Regression statistics for relationships between the number of marks (*Marks*) on calcified structures and summer flounder total length (*TL*). All regressions were significant at $P < 0.001$, and sample sizes were 80 fish.

| Structure | Equation | 100 $r^2$ |
|---|---|---|
| Scales | $Marks = -3.69 + 0.0151\ TL$ | 83 |
| Sectioned otoliths | $Marks = -3.86 + 0.0155\ TL$ | 85 |
| Whole otoliths | $Marks = -3.90 + 0.0157\ TL$ | 86 |

mation in temperate fishes (Smith et al., 1981), and 4) a narrow opaque zone as compared to the translucent zone (Smith et al., 1981). We address these issues in turn below.

We rarely observed poor calcification or poor contrast between opaque and translucent zones of whole otoliths. Rather, our procedures gave good contrast between opaque and translucent zones, so that we had high confidence in our age readings. In addition, we found only one otolith of 81 pairs to be poorly calcified. This otolith was easily aged once it was sectioned, and its pair was not poorly calcified and was aged with high confidence.

We saw little evidence that the first mark becomes obscured at older ages on whole otoliths, as indicated by our high agreement between whole and sectioned otoliths. The hypothesis that the first mark becomes obscured was

based on overlap in back-calculated sizes at the second and third marks on whole otoliths (Powell, 1982). However, size in any year class can vary greatly because summer flounder spawn over a protracted season (Smith, 1973; Morse, 1981; Able et al., 1990). Therefore, fish in adjacent year classes can be expected to overlap in size, and Powell's results do not necessarily mean that the first mark becomes obscured with age.

Smith et al. (1981) reported that summer flounder otoliths deviated from the general pattern of opaque and translucent zone formation seen in other temperate fishes and suggested that opaque zones formed in fall–winter, the reverse of the usual spring–summer formation in other temperate species. We saw no evidence of this reversal. Our fish were collected from October through December, so we should have observed opaque edges on the otolith if the timing of mark formation were reversed from other temperate fishes. Instead, we observed relatively wide translucent zones on the otolith edges. In addition, other studies have not found a reversal in the time of mark formation (Poole, 1961; Powell, 1982; Wenner et al., 1990), and Desfosse (1995) found that opaque zones appeared to form on whole otoliths at approximately the same time as scale marks (May through July). Finally, Smith et al. (1981) presented no data to support their hypothesis that opaque zones formed in the fall and winter. Indeed, their Figure 5 shows an opaque edge on a whole otolith from a summer flounder captured in June.




**Figure 6**

Comparison of presumed annual mark counts on scales and whole otoliths in summer flounder. The 45° diagonal line represents 1:1 agreement. The number of fish is indicated at each symbol.

In agreement with studies in other species (see references below), we found the translucent zone to be wider than the opaque zone on summer flounder otoliths. Smith et al. (1981) felt this was an anomalous occurrence and used it to reject whole otoliths. We disagree with their analysis, however, because many other fishes in our study area, including Atlantic croaker (Barbieri et al., 1994a), weakfish (Lowerre-Barbieri et al., 1994), and Spanish mackerel (Gaichas, 1997) have otoliths with a wide translucent zone and a narrow opaque zone. Such a pattern reflects the fact that opaque zones form over a short time period in these species: April–May in Atlantic croaker and weakfish (Barbieri et al., 1994a; Lowerre-Barbieri et al., 1994) and May–June in Spanish mackerel (Gaichas, 1997). In addition, although the sample size was limited ($n$=93), Desfosse (1995) found evidence, using marginal increments, that opaque zone formation on summer flounder otoliths occurs over a similarly short time period (May to July). Finally, regardless of whether opaque zones are narrower or wider than translucent zones, otoliths can be used for age determination if the mark can be proven annual.

## Comparative evaluation of scales

Our findings indicate that scales are inferior to, and much less desirable than, both sectioned and whole otoliths for aging summer flounder. Scales had significantly lower confidence scores and much higher reading times than sectioned and whole otoliths because marks on scales were often difficult to interpret using objective aging criteria. False marks were common, and different scales from the same fish often indicated different ages. As a result, both within- and between-reader percent agreement and agreement with whole and sectioned otolith age were undesirably low in scales, especially in fish over age 4. We feel that scales should not be used for aging summer flounder if otoliths, especially sectioned otoliths, are available.

The difficulties we found with summer flounder scales generally agree with reports in the literature. Dery (1988), Desfosse (1995), and Bolz et al. (2000), for examples, have reported similar problems interpreting scale marks. Like us, Desfosse (1995) found low within-reader scale agreement (only 46%) in fish over age 4. Desfosse (1995) reported high agreement between scales and whole otoliths (98%) for ages 0 to 5, much higher than the 85% agreement we found for ages 0 to 4. However, 90% of his fish ($n$=170) were ages 0 to 2 and only one was age 5, a likely explanation for his high percent agreement. Shepherd (1980) reported high agreement (91%) between scales and whole otoliths for moderately old fish (ages 4 to 6), but his sample size was only 21 fish, only one of which was age 6. Our study reported lower overall agreement between whole otoliths and scales (76%), but we examined fish over a much wider age range (ages 0 to 10) than previously reported.

## Comparative evaluation of opercular bones

Our comparative studies have found opercular bones to be inferior to both sectioned and whole otoliths in summer flounder, and even to scales. Opercular bones had the lowest confidence scores, the highest reading times, only 46% within reader agreement, and they often exhibited unclear transitions from translucent to opaque zones, particularly at early ages. For these reasons, we feel that opercular bones should not be used for aging summer flounder.

These findings are new for summer flounder, because no previous studies have used opercular bones to age this species. We were disappointed at the poor performance of opercular bones because they have been reported useful in many other species, including perch (LeCren, 1947), carp (McConnell, 1952), yellow perch (Bardach, 1955), northern pike (Frost and Kipling, 1957), tautog (Cooper, 1967; Hostetter and Munroe, 1993), and goldeye (Donald et al., 1992). Many of these studies show photographs of opercular bones with clear, easily recognized marks that have been interpreted as being formed annually. These studies, however, generally have not validated age determination in opercular bones; therefore it is unclear whether they give accurate ages in these other species.

### Formation of early marks on otoliths and scales

We sometimes observed an early, presumably false, mark prior to the first presumed annual mark on both otoliths and scales of summer flounder. Although we attempted not to count this early mark, it appeared to be the primary cause for disagreements between our readers in aging both otoliths and scales. This problem has not been reported in summer flounder otoliths, although there is evidence of this early false mark on scales (Dery, 1988; Bolz et al., 2000). Indeed, a primary problem cited by Bolz et al. (2000) for differences in interpretation of summer flounder scales was the choice of a first annual mark.

The early, presumably false, mark that sometimes occurred on summer flounder otoliths and scales appears similar to the first mark reported for Atlantic croaker otoliths (Barbieri et al., 1994a) and might be explained by similarities in certain life history traits of these two species. Both species have a protracted spawning season and spawn over a similar time frame in the Chesapeake Bay region: Atlantic croaker from mid-summer to late fall (Wallace, 1940; Haven, 1957; Barbieri et al., 1994b), and summer flounder from early fall to early winter (Smith, 1973; Morse, 1981; Able et al., 1990). Barbieri et al. (1994a) reported the formation of a first mark on Atlantic croaker otoliths in the first spring following hatching, at 5 to 10 months, with two patterns of early mark formation: 1) the first mark close to, but distinct from, the focus in early hatched fish, and 2) the first mark nearly continuous with the focus in late hatched fish. As with Atlantic croaker, we suggest that the first mark on summer flounder otoliths and scales, which we have referred to as an "early, presumably false, mark," might actually be laid down in the first spring following hatching, at 5 to 8 months, with the same two patterns of early mark formation.

Previous summer flounder aging studies interpret the first annual mark to be laid down on scales and otoliths in the second spring following hatching (Smith et al., 1981; Szedlmayer et al., 1992), at 17 to 20 months, one year after the first annual mark is laid down on Atlantic croaker otoliths. Despite this difference, fish from these two species that are hatched at the same time are currently placed in the same year class. It thus appears that the current age determination methods differ between these two species. For example, according to current conventions (Bolz et al., 2000), a summer flounder hatched in October 2000 would be called age 1 on 1 January 2002, at a biological age of 15 months. This age is several months *before* the first presumed annual mark is laid down on the structures in the second spring following hatching (2002), even though an "early" mark might have been laid down in the first spring following hatching (2001). Similarly, an Atlantic croaker hatched in October 2000 would be called age 1 on 1 January 2002 (Barbieri et al., 1994a), at a biological age of 15 months. However, this age is 8 months *after* the first annual mark is laid down on the otolith, which occurs in the first spring following hatching (2001). Therefore, the two species differ in the way the first annual mark is assigned.

To resolve the issue of early mark formation in summer flounder, we suggest that calcified structures of young-of-the-year fish be examined to determine when the early mark is formed, as Barbieri et al. (1994a) did for Atlantic croaker. Barbieri et al.'s (1994a) validated method automatically assigns an early first mark, formed at 5 to 10 months, to all Atlantic croaker otoliths, whether the mark is distinct or not. If the "early, presumably false, mark" in summer flounder is similar to the first annual mark in Atlantic croaker, an early first mark could likewise be assigned to summer flounder otoliths. If this were done, disagreements on the first mark on summer flounder structures would be fewer, and summer flounder and Atlantic croaker would be aged in exactly the same way. That is, both fish would already have a first annual mark on the structure when ages are advanced to 1 on the 1 January arbitrary birthdate.

## Acknowledgments

We would like to thank Chesapeake Bay commercial fishermen for their cooperation and James Owens (VIMS) for helping us to obtain samples from them. T. Ihde (VIMS) helped develop some of the aging methods and was the second reader in our study. J. Foster (VIMS) helped develop ideas for the early mark portion of the discussion. Financial support for this study was provided by a Wallop/Breaux Program Grant for Sport Fish Restoration from the U.S. Fish and Wildlife Service through the Virginia Marine Resources Commission, Project F-88-R-2.

## Literature cited

Able, K. W., R. E. Matheson, W. W. Morse, M. P. Fahay, and G. Shepherd.
1990. Patterns of summer flounder *Paralichthys dentatus* early life history in the Mid-Atlantic Bight and New Jersey estuaries. Fish. Bull. 88:1–12.

Almeida, F. P., R. E. Castaneda, R. Jesien, R. E. Greenfield, and J. M. Burnett.
1992. Proceedings of the NEFC/ASMFC summer flounder, *Paralichthys dentatus*, aging workshop, 11–13 June 1990, Northeast Fisheries Center, Woods Hole, Massachusetts. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NEC-89, 7 p.

Bagenal, T. B., and F. W. Tesch.
1978. Age and growth. *In* Methods for assessment of fish

production in fresh waters, 3rd ed. (T. B. Bagenal, ed.), p. 101–136. Blackwell Scientific Publications, Oxford.

Barbieri, L. R., M. E. Chittenden Jr., and C. M. Jones.
1994a. Age, growth, and mortality of Atlantic croaker, *Micropogonias undulatus*, in the Chesapeake Bay region, with a discussion of apparent geographic changes in population dynamics. Fish. Bull. 92:1–12.

Barbieri, L. R., M. E. Chittenden Jr., and S. K. Lowerre-Barbieri.
1994b. Maturity, spawning, and ovarian cycle of Atlantic croaker, *Micropogonias undulatus*, in the Chesapeake Bay and adjacent coastal waters. Fish. Bull. 92:671–685.

Bardach, J. E.
1955. The opercular bone of the yellow perch (*Perca flavescens*), as a tool for age and growth studies. Copeia 2:107–109.

Beamish, R. J.
1979. Differences in the age of Pacific hake (*Merluccius productus*) using whole otoliths and sections of otoliths. J. Fish. Res. Board Can. 36:141–151.

Beamish, R. J., and G. A. McFarlane.
1983. The forgotten requirement for age validation in fisheries biology. Trans. Am. Fish. Soc. 112:735–743.

Bigelow, H. B., and W. C. Schroeder.
1953. Fishes of the Gulf of Maine. U.S. Fish Wildl. Serv., Fish. Bull. 53:1–577.

Bolz, G. R., J. P. Monaghan Jr., K. L. Lang, R. W. Gregory, and J. M. Burnett.
2000. Proceedings of the summer flounder aging workshop, 1–2 February 1999, Woods Hole, Massachusetts. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NE-156, 15 p.

Chilton, D. E., and R. J. Beamish.
1982. Age determination methods for fishes studied by the Groundfish Program at the Pacific Biological Station. Can. Spec. Publ. Fish. Aquat. Sci. 60:1–102.

Cooper, R. A.
1967. Age and growth of the tautog, *Tautoga onitis* (Linnaeus), from Rhode Island. Trans. Am. Fish. Soc. 96:134–142.

Dery, L. M.
1988. Summer flounder, *Paralichthys dentatus*. *In* Age determination methods for Northwest Atlantic species (J. Penttila and L. M. Dery, eds.), p. 97–102. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 72.

Desfosse, J. C.
1995. Movements and ecology of summer flounder, *Paralichthys dentatus*, tagged in the southern Mid-Atlantic Bight. Ph.D. diss., College of William and Mary, Williamsburg, VA, 187 p.

Donald, D. B., J. A. Babaluk, J. F. Craig, and W. A. Musker.
1992. Evaluation of scales and operculum methods to determine age of adult goldeyes with special reference to a dominant year-class. Trans. Am. Fish. Soc. 121:792–796.

Eldridge, P. J.
1962. Observations on the winter trawl fishery for summer flounder, *Paralichthys dentatus*. M.S. thesis, College of William and Mary, Williamsburg, VA, 58 p.

Frost, W. E., and C. Kipling.
1957. The determination of the age and growth of pike (*Esox lucius* L.) from scales and opercular bones. J. Cons. 24:314–341.

Gaichas, S. K.
1997. Age and growth of Spanish mackerel, *Scomberomorus maculatus*, in the Chesapeake Bay region. M.S. thesis, College of William and Mary, VA, 111 p.

Ginsburg, I.
1952. Flounders of the genus *Paralichthys* and related genera in American waters. U.S. Fish Wildl. Serv., Fish. Bull. 52:267–351.

Gutherz, E. J.
1967. Field guide to the flatfishes of the Family Bothidae in the western North Atlantic. U.S. Fish Wildl. Serv. Circ. 263, 47 p.

Haven, D. S.
1957. Distribution, growth, and availability of juvenile croaker, *Micropogon undulatus*, in Virginia. Ecology 38:88–97.

Hildebrand, S. F., and W. C. Schroeder.
1928. Fishes of Chesapeake Bay. Bull. U.S. Bur. Fish 43:1–366.

Hostetter, E. B., and T. A. Munroe.
1993. Age, growth, and reproduction of tautog *Tautoga onitis* (Labridae:Perciformes) from coastal waters of Virginia. Fish. Bull. 91:45–64.

LeCren, E. D.
1947. The determination of the age and growth of the perch (*Perca fluviatilis*) from the opercular bone. J. Anim. Ecol. 16:188–204.

Leim, A. H. and W. B. Scott.
1966. Fishes of the Atlantic Coast of Canada. Fish. Res. Board Can., Bull. 155:1–485.

Lowerre-Barbieri, S. K., M. E. Chittenden Jr., and C. M. Jones.
1994. A comparison of a validated otolith method to age weakfish, *Cynoscion regalis*, with the traditional scale method. Fish. Bull. 92:555–568.

MAFMC (Mid-Atlantic Fishery Management Council).
1987. Fishery management plan for the summer flounder fishery, 159 p. MAFMC. Dover, DE.

McConnell, W. J.
1952. The opercular bone as an indicator of age and growth of the carp, *Cyprinus carpio* Linnaeus. Trans. Am. Fish. Soc. 81:138–149.

Media Cybernetics.
1999. Optimas, version 6.5. Media Cybernetics, Silver Spring, MD.

Morse, W. W.
1981. Reproduction of the summer flounder, *Paralichthys dentatus* (L.). J. Fish. Biol. 19:189–203.

Poole, J. C.
1961. Age and growth of the fluke in Great South Bay and their significance to the sport fishery. N.Y. Fish Game J. 8:1–18.
1962. The fluke population of Great South Bay in relation to the sport fishery. N.Y. Fish Game J. 9:93–117.

Powell, A. B.
1982. Annulus formation on otoliths and growth of young summer flounder from Pamlico Sound, North Carolina. Trans. Am. Fish. Soc. 111:688–693.

Shepherd, G.
1980. A comparative study of ageing methods for summer flounder (*Paralichthys dentatus*). Lab. Ref. Doc. 80-13, Northeast Fisheries Science Center, NMFS, Woods Hole, MA, 26 p.

Smith, R. W., and F. C. Daiber.
1977. Biology of the summer flounder, *Paralichthys dentatus*, in Delaware Bay. Fish. Bull. 75:823–830.

Smith, R. W., L. M. Dery, P. G. Scarlett, and A. Jearld Jr.
1981. Proceedings of the summer flounder (*Paralichthys dentatus*) age and growth workshop, 20–21 May 1980, NEFC, Woods Hole, Massachusetts. U.S. Dep Commer., NOAA Tech. Memo. NMFS-F/NEC-11, 30 p.

Smith, W. G.
1973. The distribution of summer flounder, *Paralichthys dentatus*, eggs and larvae on the continental shelf between Cape Cod and Cape Lookout. Fish. Bull. 71:527–535.

Szedlmayer, S. T., K. W. Able, and
R. A. Rountree.
1992. Growth and temperature induced mortality of young-of-the-year summer flounder (*Paralichthys dentatus*) in southern New Jersey. Copeia 1992:120–128.

Wallace, D. H.
1940. Sexual development of the croaker, *Micropogon undulatus*, and distribution of the early stages in Chesapeake Bay. Trans. Am. Fish. Soc. 70:475–482.

Wenner, C. A., W. A. Roumillat, J. E. Moran Jr., M. B. Maddox,
L. B. Daniel, and J. W. Smith.
1990. Investigations of the life history and population dynamics of marine recreational fishes in South Carolina: part 1. South Carolina Wildl. Mar. Resourc. Dep., Mar. Resourc. Res. Instit., 26 p.

Zar, J. H.
1996. Biostatistical analysis. Prentice Hall, Upper Saddle River, NJ, 662 p.

# Bridle efficiency of a survey trawl for flatfish

**David A. Somerton**
**Peter Munro**

Alaska Fisheries Science Center
7600 Sand Point Way NE
Seattle, Washington 98125
E-mail address (for D. A. Somerton): david.somerton@noaa.gov

**Abstract**—A bottom trawl catch of flatfish is composed of fish that were initially in the path of the trawl net and fish that were initially in the path of the bridles and were subsequently herded into the net path. Bridle efficiency (i.e. the proportion of fish in the bridle path that are herded into the net path) for seven species of flatfish was estimated by fitting a model of the herding process to data collected during a field experiment. The experiment consisted of repeatedly making trawl hauls with three different bridle lengths. The model was then fitted to the catch of each species, by length class, as a function of the widths of the net and door spreads at the three bridle lengths. Bridle efficiency was independent of body length for five species (yellowfin sole, *Limanda aspera*; flathead sole, *Hippoglossoides elassodon*; rock sole, *Lepidopsetta bilineata*; Dover sole, *Microstomus pacificus*; rex sole, *Glyptocephalus zachirus*) and ranged from 0.22 for rex sole to 0.40 for rock sole. Bridle efficiency declined with increasing body length for two species (English sole, *Parophrys vetulus*; Pacific sanddab, *Citharichthys sordidus*), ranging from about 0.35 to 0.10 over the lengths sampled.

Manuscript accepted 15 March 2001.
Fish. Bull. 99:641–652 (2001).

Swept-area estimates of biomass produced by bottom trawl surveys are typically used as relative indices of stock size rather than absolute measures because the sampling efficiency of the trawls[1] is rarely known. As a means to estimate trawl efficiency, Dickson (1993a) developed a model describing the trawling process and the associated fish behavior from an undisturbed state before arrival of a fishing vessel until passage of a trawl codend. For benthic species that remain closer to the sea bottom than the head-rope height, this model specifies that trawl efficiency is primarily determined by two processes: escapement of fish out of the net path by passing either under the footrope or through the mesh (net efficiency) and herding of fish into the net path by the doors, mudclouds, and bridles (bridle efficiency). In our study, we focused on fish herding, considering the problem of quantitatively estimating bridle efficiency or the proportion of fish encountering the bridles that are herded into the net path.

Underwater observations of trawls in operation have revealed that the process of herding differs between semipelagic species, such as Atlantic cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) that tend to swim near the bottom, and benthic species, such as flatfish that often remain in direct contact with the bottom (Main and Sangster, 1981b). Atlantic cod are apparently stimulated to herd by the sight of the trawl doors and the mudclouds that form in the wake of the doors, whereas flatfish are stimulated to herd by the close proximity or actual touch of the doors and lower bridles as they sweep along the bottom (Hemmings, 1969; High, 1969; Martyshevskii and Korotkov, 1969; Main and Sangster, 1981a, 1981b). Flatfish, once stimulated to move, tend to swim a relatively short distance in a direction perpendicular to the bridles where they slow down or settle on the bottom until the bridle again overtakes them. This pattern is usually repeated, producing what appears to an observer on the trawl headrope as a zigzag path toward the center of the trawl footrope (Main and Sangster, 1981b). While this is occurring, however, flatfish may either swim over the bridle or become exhausted and be overtaken by the bridle and thereby escape the herding process. Bridle efficiency is determined by the relative frequency of such events.

Perhaps because the mechanism of herding seems more obvious for flatfish than other types of fish, the earliest mathematical models of fish herding were based on the contact-swim-contact pattern of flatfish herding (Hemmings, 1969; Foster et al., 1981; Fuwa et al,. 1988; Fuwa, 1989). These models were focused on individual fish and typically considered both the physical aspects of the trawl, such as shape and speed, as well as the biological aspects of the fish, such as size and endurance. Although these models provided a more structured way of examining fish herding and allowed the quantitative evaluation of the relative importance of the various aspects of the process, they were of limited use in estimating bridle efficiency of specific trawls because some of the required parameters were unknown or difficult to estimate.

The approach to modeling fish herding changed markedly after Engås and Godø (1989a) conducted trawling ex-

[1] Throughout this paper, we will refer to the trawl as an entire fishing gear comprising the net, bridles, and doors.

periments to quantitatively examine the effects of variations in bridle length on catch for each area and fish size. Reasoning that the changes in catch that accompanied the changes in bridle length provided information on the efficiency of herding, Dickson (1993b) estimated the bridle efficiency coefficients for Atlantic cod and haddock by fitting a simple linear model of the trawl herding process to data from Engås Godø's (1989a) experiments. Ramm and Xiao (1995) conducted a similar trawl herding experiment and developed a new type of herding model appropriate for cases in which escapement at the center of the trawl is zero.

In our study, we extended Dickson's (1993a) herding model so that it is more specific to the peculiarities of flatfish herding and additionally extended Dickson's (1993b) approach to fitting the model to data by providing a more rigorous statistical foundation. The model was then applied to herding data for the 83-112 Eastern bottom trawl (Armistead and Nichol, 1993) used by the Alaska Fisheries Science Center (AFSC) to conduct its annual bottom trawl survey of the Eastern Bering Sea. These data were collected during two herding experiments, patterned after those of Engås and Godø (1989a), in which emphasis was placed on seven species of flatfish: yellowfin sole (*Limanda aspera*), flathead sole (*Hippoglossoides elassodon*), rock sole (*Lepidopsetta bilineata*), English sole (*Parophrys vetulus*), Dover sole (*Microstomus pacificus*), rex sole (*Glyptocephalus zachirus*), and Pacific sanddab (*Citharichthys sordidus*).

## Materials and methods

### Development of the herding model

Consider that the area of the bottom swept by a trawl consists of two components, the area between the wingtips of the net (net path), and the area between the wing tips and the doors (bridle path; Fig. 1). The catch ($N$) obtained in a trawl can be represented as some proportion, $k_n$, of the number of fish in the net path ($N_n$) plus some proportion, $P$, of the number of fish in the bridle path ($N_b$). If the probability of capture for a fish herded into the net path is identical to that of a fish initially in the net path, then $P$ can be considered equal to $k_n k_b$, where $k_b$ is the bridle efficiency[2] or the proportion of fish within the bridle path that is herded into the net path. Algebraically this is expressed as

$$N = k_n N_n + k_n k_b N_b. \quad (1)$$

The numbers of fish within the net and bridle paths are equal to the fish density ($D$) multiplied by the path areas, that is,

$$\begin{aligned} N_n &= DLW_n \\ N_b &= DL\,(W_d - W_n), \end{aligned} \quad (2)$$

where $L$ = tow length;
$W_n$ = the width of the net path; and
$W_d$ = the width of the door path (Fig. 1).

Combining all terms, the total catch then can be expressed as

$$N = k_n DLW_n + k_n k_b DL\,(W_d - W_n). \quad (3)$$

At this point the model is quite similar to the one proposed by Dickson (1993a) to describe the trawl catching process for Atlantic cod and haddock. We now modify the model to make it more specific to flatfish by considering that herding is restricted to the portion of the bridle path in which the lower bridle is in contact with the bottom. This can be expressed as

$$N = k_n DLW_n + k_n hDL\,(W_d - W_n - W_{off}), \quad (4)$$

where $W_{off}$ = the width of the path in which the bridle is not in contact with the bottom; and
$h$ = the herding coefficient or the proportion of fish in the bridle contact path that is herded into the net path.

Thus, $k_b$ is the average efficiency over the entire bridle path and $h$ is the average efficiency in the portion of the bridle path where herding actually occurs.

### Description of the herding experiments

The strategy that we used in our herding experiments was to repeatedly trawl in such a way that the width of the net path was held approximately constant, whereas the width of the door path was varied among three distances by changing the length of the bridles. The first of two herding experiments was conducted 25 July–2 August 1994 aboard the FV *Arcturus* in the eastern Bering Sea and focused on yellowfin sole, flathead sole, and rock sole. The second experiment was conducted 14–25 September 1994 aboard the RV *Alaska* off the coast of Washington State and focused on English sole, Dover sole, rex sole, and Pacific sanddab. For both experiments, we used a blocked sampling design to minimize the effects on catch of the spatial variation in fish density. In each geographic block, three nearby, but nonoverlapping, 30-min trawl hauls at a speed of 1.5 m/s were made with each of three bridle lengths chosen in random order. Bridles measured 27 m, 55 m (the standard length used on AFSC surveys), and 82 m in length and were constructed of 16-mm diameter steel cable. Tailchains connecting the doors to the bridles (Fig. 1) were always constructed of 13-mm diameter long link chain but differed in length between vessels (Table 1). Trawl doors were always a "V" style measuring 1.8 m × 2.7 m and weighing 910 kg. On all hauls, spreads of the doors were measured at the tops of the wings simultane-

[2] In Dickson (1993a) this same quantity is referred to as sweep efficiency ($k_s$). We use the term "bridle efficiency" because the 83-112 Eastern trawl does not have sweeps connecting the bridles to the doors as does the trawl considered by Dickson.




**Figure 1**

Definitions of the trawl measurements. Included are the tailchain length, bridle length, length of the bridle not in contact with the bottom ($L_{off}$), width of the door path ($W_d$), width of the net path ($W_n$), width of the bridle path not in contact with the bottom ($W_{off}$), and bridle angle ($\alpha$).

ously and continuously with an acoustic trawl mensuration system. Tow length was measured as the straight line distance between the GPS positions of first and last footrope contact with the bottom determined with a time-depth recorder attached to the headrope. Bottom temperature was recorded at two second intervals with a bathythermograph and averaged 3.0°C during the eastern Bering sea experiment and 7.0°C in the West Coast experiment. Bottom depth averaged 76 m during the eastern Bering sea experiment and 67 m in the West Coast experiment. The catch from each haul was first sorted to species; then the catch of each species was weighed in the aggregate and fish of each species were individually measured for total length in centimeters. Both experiments were conducted during daylight hours to best approximate the sampling protocol used on most AFSC bottom trawl surveys.

## Estimating $W_{off}$

The herding experiments provided data on $N$, $L$, $W_d$, and $W_n$, but obtaining data to estimate $W_{off}$ required a different type of experiment. We obtained the necessary data by directly observing the contact of the lower bridle with the bottom on an experimental cruise conducted 9–12 Septem-

**Table 1**

Trawl configuration parameters for each herding experiment. Included are the bridle and tail chain lengths that were used, the number of sampling blocks occupied, and the means and standard deviations (in parentheses) of the wing spread, door spread, and bridle angle. Also included are the mean values for the bridle off-bottom path width ($W_{off}$). WA = Washington State.

| Experiment | Bridle length (m) | Tailchain length (m) | Number of blocks | Wing spread (m) ($W_n$) | Door spread (m) ($W_d$) | Bridle angle (deg) ($\alpha$) | Bridle off-bottom path width (m) ($W_{off}$) |
|---|---|---|---|---|---|---|---|
| Bering Sea | 27.3 | 13.9 | 15 | 18.0 (0.6) | 44.6 (2.1) | 18.9 (1.2) | 23.3 |
| | 54.6 | 13.9 | 15 | 17.3 (0.6) | 58.7 (2.9) | 17.6 (1.0) | 21.8 |
| | 81.6 | 13.9 | 15 | 17.1 (0.5) | 71.6 (3.7) | 16.6 (1.0) | 20.5 |
| West Coast (WA) | 27.3 | 8.0 | 19 | 17.7 (0.3) | 41.9 (0.7) | 20.0 (0.5) | 24.1 |
| | 54.6 | 8.0 | 19 | 17.1 (0.6) | 57.5 (1.5) | 18.8 (0.7) | 23.1 |
| | 81.6 | 8.0 | 19 | 16.8 (0.3) | 70.0 (1.5) | 17.3 (0.5) | 21.3 |

ber 1995 aboard the RV *Alaska* off the coast of Washington State using the 83-112 Eastern trawl equipped with standard 55-m bridles. The experiment consisted of recording views of the lower bridle with a silicon-intensified target (SIT) video camera while the trawl was in operation. The camera was mounted in a positively buoyant case attached to the upper bridle with a 1-m tether line and aligned so that the lower bridle directly below the attachment point could be viewed. The tether was positioned at 5-m intervals between 23 m and 43 m behind the doors. One 15-min tow at each of the five bridle positions was taken at both depths of 20 m and 35 m.

The video tapes were subsequently analyzed to determine the degree of bottom contact at each bridle position. This was done by viewing each tow at ten randomly chosen 10-sec sampling intervals. Bottom contact was recognized by the presence of sediment that is mixed into the water at the point of contact. Because the bottom is irregular, when the bridle has weak contact, just the tops of the irregularities are touched, whereas when the bridle has strong contact, the entire bottom in the field of view is touched. The average percentage of the bridle in the field of view that was in contact with the bottom was scored according to the following four-level scale: 1) no contact, 2) ≤25% contact, 3) >25% and ≤75% contact, 4) >75% contact. The degree of bottom contact at each bridle position and depth was then estimated as the mean of the ten evaluations.

The length of the bridle not in contact with the bottom ($L_{off}$, Fig. 1) was defined as the distance between the attachment of the tailchain to the door and the point along the bridle where bottom contact was 50%. To estimate $L_{off}$, a logistic function was fitted to the bottom-contact and bridle-position data with generalized linear modeling (Venables and Ripley, 1994), then the fitted logistic equation was evaluated at a bottom contact score of 3.0 (i.e. bottom contact >25% and <75%). Variance of $L_{off}$ was estimated with bootstrapping (Efron and Tibshirani, 1993), where the bootstrap samples were obtained by randomly choosing, with replacement, from the haul mean scores at each bridle position.

In subsequent analysis, $L_{off}$ was treated as a constant for all three bridle lengths and for both vessels. However the variable actually used in the herding model, $W_{off}$, was calculated as $L_{off}$ multiplied by the sine of the bridle angle (Fig. 1) and therefore varied slightly between bridle lengths and experiments because the bridle angle varied.

In addition to the camera placements for measuring $L_{off}$, we also made placements at bridle positions both closer to the doors to examine for any evidence of herding in the area where the lower bridles were not in contact with the bottom and closer to the wing tips to determine if bottom contact was maintained continuously near the junction of the bridle and wing.

## Fitting the model to the herding data

The herding model (Eq. 4) was modified in several ways to clarify the way it was fitted to the experimental data and to better define its underlying statistical structure. First, because of the block design of the experiment, fish density ($D$) was considered to be a constant within each block but to vary between blocks. Furthermore, net efficiency ($k_n$) was also considered to be a constant for all tows within a block because depth and other bottom conditions are nearly constant but vary between blocks. Because $D$ and $k_n$ are both block-specific constants and are confounded in the model, they were combined into a new constant $k$. Second, herding can vary with fish length (Engås and Godø, 1989a; Dickson, 1993b), therefore Equation 4 was modified to allow length dependency. The modified equation is

$$N_{ijk} = k_i\left[LW_n\right]_{ij} + k_i h_k\left[L(W_d - W_n - W_{off})\right]_{ij} + \varepsilon \qquad (5)$$

where the subscript $i$ = block number;
$j$ = bridle length within a block;
$k$ = fish length class and
$\varepsilon$ = ~ $N(0, \sigma)$.

For each size class, fitting Equation 5 to the herding data required estimation of $N$+1 parameters, where $N$ is the number of blocks sampled (i.e. a unique value of $k$ for each block and a common value of $h$ for all blocks). Because the model is nonlinear, because of the product of $k$ and $h$, it was fitted to the catch and trawl measurement data by using nonlinear regression (Venables and Ripley, 1994). Fish length classes used in the calculations were chosen such that the number of length observations was approximately equal among classes. Ten length classes were usually chosen, but for species with narrow length ranges, as few as eight were used. After fitting the model, estimates of the variances of the parameters were obtained from the inverse of the information matrix (Seber and Wild, 1989).

Bridle efficiency ($k_b$) for each size class was calculated from the estimated value of the herding coefficient ($h$) by using the relationship

$$k_b = h\left(1 - \frac{W_{off}}{(W_d - W_n)}\right). \quad (6)$$

Although up to this point $h$ was treated as a constant for all bridle lengths, $k_b$ is not a constant because the proportion of the bridle path in contact with the bottom will vary with bridle length. Therefore $k_b$ was evaluated only for the standard bridle length. Variance of $k_b$ was approximated as

$$Var(k_b) = \left(1 - \frac{W_{off}}{(W_d - W_n)}\right)^2 Var(h). \quad (7)$$

Although the model that we used allows for length-dependent herding, herding may not be a length-dependent process in all species. We performed a simple test of linear length-dependency by regressing the estimated values of $k_b$ on the midpoints of the associated length intervals. For species in which the slope was not significantly different from zero, the herding model was refitted to the data with all length intervals combined. For species in which the slope differed significantly from zero, we calculated a functional relationship between $k_b$ and fish length. Among the possible relationships examined, the best fit was obtained with the exponential model

$$k_b = a + b \exp(cL), \quad (8)$$

where a, $b$, $c$ are parameters; and

$L$ = the midpoints of the length classes.

Model fitting was done by using nonlinear regression. Variance of the predicted value of $k_b$ was estimated with bootstrapping (Efron and Tibshirani, 1993), where the bootstrap samples were obtained by sampling the residuals from the regression, with replacement, and adding them to the predicted values of $k_b$. The functional relationship between the standard deviation of $k_b$ and fish length was approximated by using a parabolic model fitted by linear regression.

One implicit assumption of the herding model is that $h$ is constant for all three bridle lengths. We tested the validity of this assumption by comparing the goodness of fit of a modified version of the herding model to that of the original. In the modified model, the herding coefficient was represented as

$$h_l = h + dL,$$

where $h_l$ = the herding coefficient at bridle length $L$;

$h$ = the mean herding coefficient for all bridle lengths;

$d$ = a free parameter; and

$L$ = a coded value of bridle length equal to –1, 0, 1 for the short, medium, and long bridles (with this notation, negative values of $d$ indicate that $h_l$ declines with increasing bridle length).

Because this is a comparison of two nonlinear models with differing numbers of parameters, we used as a measure of fit the penalized sum of squares; that is, RSS/($n$–2$p$), where RSS is the residual sum of squares, $n$ is the number of data values, and $p$ is the number of parameters (Hilborn and Mangel, 1997, page 115). A linear change in $h_l$ is considered significant when the penalized sum of squares for the modified model is less than that for the original model.

## Results

### Herding experiments

Fifteen blocks were completed in the Bering Sea herding experiment; 19 blocks were completed in the West Coast herding experiment. Although the intention of the experiments was to maintain a constant trawl geometry at all bridle lengths, in both experiments wing spread decreased 0.9 m and bridle angle decreased an average of 2.5% from the shortest to the longest bridles (Table 1). The experimental change in bridle length had a pronounced effect on the area swept by the bridles. In the Bering Sea experiment, for example, total bridle length (i.e. bridle plus tailchain length) was increased by 143% from the shortest to the longest bridle, which, in turn, produced a 64% increase in door path width ($W_d$) and a 113% increase in total bridle path width ($W_d - W_n$). However, this increase in bridle length also produced a 930% increase in the bridle contact path width ($W_d - W_n - W_{off}$; Table 1) because only a small portion of the bridle remained in contact with the bottom at the shortest bridle length.

Catch per area swept by the net increased significantly with increasing bridle length for six of the seven species (rock sole, yellowfin sole, flathead sole, Dover sole, English sole, rex sole; weighted linear regression, $P$<0.010) indicating strong herding by the bridles (Fig. 2). For Pacific sanddab, however, the increase was not significant ($P$=0.096).

Fish length changed significantly ($P$<0.05) with increasing bridle length for only two species (English sole and Pacific sanddab; Table 2) and in both cases smaller fish were




**Figure 2**

Catch in numbers per hectare swept by the net (i.e. tow length multiplied by net width), by species, plotted against the bridle length in meters. A straight line, fitted to the data with weighted (inverse of the variance) linear regression, is also shown for each species. Significance of the slope of this line indicates that herding contributes to the catch.

caught at longer bridle lengths. Because four of the remaining five species also displayed declining, but nonsignificant, changes in fish length, a decrease in fish length with increasing bridle length seems to be a general flatfish pattern for this trawl. In all cases, however, the magnitude of the decrease was quite small (Fig. 3).

**Table 2**

Relationship of mean fish length to increasing bridle length. The total number of fish measured, the slope of the regression line fitted to mean length at each tow versus bridle length, and the probability that the slope equaled zero is shown for each species. Only English sole and Pacific sanddab are significant. Note that a negative slope indicates that smaller fish are caught with longer bridles.

| Species | Number of fish measured | Slope | $P$(slope=0) |
|---|---|---|---|
| Rock sole | 23,586 | –.075 | 0.23 |
| Yellowfin sole | 7231 | 0.008 | 0.83 |
| Flathead sole | 2927 | –0.311 | 0.31 |
| Rex sole | 22,774 | –0.052 | 0.56 |
| Dover sole | 12,507 | –0.036 | 0.75 |
| English sole | 21,968 | –0.277 | <0.01 |
| Pacific sanddab | 34,209 | –0.383 | <0.01 |

### Estimating $W_{off}$

The degree of contact of the lower bridles with the bottom varied with increasing distance from the doors. Contact never occurred in the first 28 m (Fig. 4). At greater distances, contact became intermittent and was focused primarily at the tops of sand ripples. Contact reached 50% at 36 m and nearly 100% at 40 m (Fig. 4). The estimated value for $L_{off}$ was 36.0 m (95% confidence intervals; ±1.4 m). This value was assumed to be constant for all three bridle lengths and both vessels.

### Fit of the model

Of the seven species examined for length-dependent herding, only Pacific sanddab and English sole had significant slopes in the regressions of $k_b$ on fish length (Table 3, Fig. 5). In both cases the slope was negative indicating that herding efficiency decreased with increasing fish length.

For the five species displaying no significant length-dependent herding, the estimated herding efficiency in the bridle contact path ($h$) varied from 0.50 for rex sole to 0.84 for rock sole, whereas the herding efficiency in the entire bridle path ($k_b$) varied from 0.22 to 0.40 (Table 4). For the two species showing significant length-dependent herding, English sole and Pacific sanddab, $k_b$ varied from about 0.35 at the shortest lengths to about 0.08–0.10 at the longest lengths (Fig. 5).

### Test of model assumptions

For six of the seven species, the fit of a modified herding model, with $h$ parameterized as a linear function of bridle length, produced a larger penalized sum of squares than the fit of the original model with a constant $h$. This finding indicates that the fit was not improved by the specification of a linear variation in $h$, and that the assumption of constancy of $h$ at all bridle lengths is valid. However, for Pacific sanddab the fit was significantly improved by the modification. Because the sign of the slope parameter of the linear change was negative, it indicates that for this species herding was less effective at longer bridle lengths.

**Table 3**

Regression of $k_b$ on the midpoints of fish-length classes. The number of length classes, the slope of the regression line, and the probability that the slope equals zero are shown for each species. Only English sole and Pacific sanddab have significant ($P$<0.05) slopes.

| Species | Number of size classes | Slope | $P$(slope=0) |
|---|---|---|---|
| Rock sole | 10 | –0.006 | 0.088 |
| Yellowfin sole | 10 | 0.001 | 0.767 |
| Flathead sole | 10 | –0.004 | 0.249 |
| Rex sole | 10 | –0.004 | 0.354 |
| Dover sole | 9 | –0.003 | 0.564 |
| English sole | 8 | –0.027 | 0.006 |
| Pacific sanddab | 10 | –0.016 | 0.008 |

## Discussion

### Bridle efficiency

Herding by the bridles of the 83-112 Eastern trawl substantially contributes to the catch of flatfish. For rock sole, as an example, approximately 49% of the catch comprised fish that were herded into the net path.[3] This contribution is ignored when swept-area estimates of abundance are based on wing spread rather than door spread. Although such an assumption could lead to an overestimate of abundance, herding is only one component of the trawl catching process and the gains from herding could be offset by the losses due to net escapement.

Bridle efficiency did not vary with fish length for five of the seven species examined (Table 3); consequently, for these species, herding by the bridles is not length selective. A similar lack of length selection was observed by Bridger (1969) in a study examining the herding of lemon sole (*Pseudopleuronectes americanus*). For English sole and Pacific sanddab, however, bridle efficiency clearly declined with increasing fish length and therefore herding is length selective.

Such length selectivity should lead to a change in mean fish length when the average duration of herding experienced by an individual fish is increased by an increase in bridle length. Without exception, the changes in mean fish length that we observed with increasing bridle length are

[3] From Equation 4, it can be shown that the herded proportion in the catch is equal to $hw_{on}/(w_u + hw_{on})$, where $w_{on} = w_d - w_u - w_{off}$.




**Figure 3**

Length-frequency distributions, expressed as proportions, by bridle length. Thin solid lines represent short bridles, medium solid lines represent standard bridles, and thick solid lines represent long bridles.

consistent with the observed changes in bridle efficiency with fish length. All five species with length-independent herding efficiency displayed no change in mean length, whereas both species with herding efficiency that declined with fish length displayed a decline in mean length with increasing bridle length (Tables 2 and 3).

**Table 4**

Estimates of bridle efficiency, $k_b$, the standard deviation (SD) of $k_b$ by species, and the $r^2$ for the model fitted without length dependency. For the two species where $k_b$ varies with length, parameters for predicting $k_b$ and SD($k_b$) as functions of fish length are provided. The two functions are $k_b = a + b \exp(cL)$ and SD($k_b$)$= a + bL + cL^2$, where $L$ = fish length in centimeters; and $h$ = the average efficiency in the portion of the bridle where herding occurs.

| | | | | | Parameters of $k_b = f$(fish length) | | | Parameters of SD($k_b$) = $f$(fish length) | | |
|---|---|---|---|---|---|---|---|---|---|---|
| Species | $r^2$ | $h$ | $k_b$ | SD($k_b$) | $a$ | $b$ | $c$ | $a$ | $b$ | $c$ |
| Rock sole | 0.91 | 0.840 | 0.400 | 0.056 | | | | | | |
| Yellowfin sole | 0.77 | 0.580 | 0.275 | 0.057 | | | | | | |
| Flathead sole | 0.70 | 0.510 | 0.242 | 0.065 | | | | | | |
| Rex sole | 0.71 | 0.502 | 0.216 | 0.061 | | | | | | |
| Dover sole | 0.80 | 0.636 | 0.274 | 0.064 | | | | | | |
| English sole | 0.81 | 0.384 | 0.165 | 0.041 | 0.033 | 42.667 | –0.221 | 0.3338 | –0.0232 | 0.00042 |
| Pacific sanddab | 0.83 | 0.161 | 0.070 | 0.030 | 0.046 | 36.783 | –0.354 | 0.0774 | –0.0059 | 0.00012 |

## Form of the herding model

The herding model that we propose is essentially the same as Dickson's (1993a) model except that it includes an additional parameter ($W_{off}$) needed to correct a potential shortcoming of the original. This shortcoming can be seen in the following example, which for simplicity considers a herding experiment with only two bridle lengths. Starting with Dickson's model (similar to Eq. 3 in our study), an estimator for $k_b$ can be derived from the quotient of the catches obtained with long bridles to the catches obtained with short bridles:

$$k_b = \frac{W_n - RW_n}{R(W_{d1} - W_n) - (W_{d2} - W_n)},$$

where $R = N_2 / N_1$;
$W_{d1}, W_{d2}$ = the door spreads with the shorter and longer bridles; and all other variables are the same as those in Equation 3.

Estimates of $k_b$, based on values of $W_{d1}$, $W_{d2}$, and $W_n$ from the Bering sea experiment, increase with catch ratio ($R$) and reach a value of 1.0 when $R$ equals the ratio of the door spreads ($W_{d2}/W_{d1}$). This increase indicates that the bridles are completely efficient ($k_b$=1.0) when the proportionate increase in the area subjected to herding equals the proportionate increase in catch. If $R$ exceeds the ratio of the door spreads, then $k_b$>1.0, which is not feasible. In our herding experiments, however, $R$ exceeded door spread ratios for all species except English sole and Pacific sanddab; therefore, in most cases the unmodified herding model failed to provide feasible estimates of $k_b$. By comparison, estimates of $k_b$ from the modified model are feasible over the entire observed range of catch ratios.

In our initial attempts to fit the herding model to experimental data, we treated $W_{off}$ as an additional parameter to be estimated in the fitting process. However, it soon became clear that $W_{off}$ and $h$ are confounded in the model and that convergent solutions required an independent estimate of $W_{off}$. We were able to obtain such an estimate and observe fish behavior near the bridle by using video methods, because in the standard configuration of the 83-112 Eastern trawl the bridles are not obscured by mudclouds over much of their lengths. However, in perhaps more typical cases where the bridles are obscured by the mudclouds, estimation of $W_{off}$ with video methods would be problematic. A further complication in such cases is that the mudclouds themselves might provide a herding stimulus and confuse the interpretation of $W_{off}$.




**Figure 4**

Relative bottom contact of the lower bridle plotted against the distance measured from the door for each camera position and depth. Also shown is the fit of a logistic equation. $L_{off}$ is defined as the distance (vertical dotted line) corresponding to a relative bottom score of 3.0 (horizontal dotted line).




**Figure 5**

Estimated bridle efficiency ($k_b$) plotted against fish length in centimeters. The fit of an exponential model (Eq. 8) to the data is also shown for English sole and Pacific sanddab, the two species with significant decreases in $k_b$ with increasing length.

## Model assumptions

The adequacy of the herding model and the appropriateness of our herding efficiency estimates rest on a variety of assumptions. First, we assumed herding efficiency was independent of bridle length. If, however, exhaustion was a primary determinant of bridle efficiency, then it is possible that individual endurance would be exceeded more

frequently at longer bridle lengths and would thereby diminish herding efficiency. This assumption was tested by comparing the fit of a herding model with a bridle-length dependent $h$ to that of an unmodified model. The modified model provided a better fit for only one species, Pacific sanddab. Because this species is the smallest of the seven (Fig. 3), it is likely that its swimming endurance might have been exceeded at the longest bridle length. However, for the remaining species the assumption is valid.

Second, we assumed that none of the fish in the area spanned by $W_{off}$ are herded into the net path. Although we recognize that the doors and the mudclouds immediately behind the doors must herd some fish toward the net path, such fish would be herded into the section of the bridle unobscured by the mudcloud and likely escape beneath the lower bridle. We attempted to confirm this possibility by examining video observations from this area of the bridle to see if flatfish escaped or showed any herding behavior, but the results were equivocal because few fish, and probably only moving fish, were seen.

Third, we assumed that in changing the length of the bridles we did not alter trawl geometry in any way other than by increasing the area exposed to herding. As in the studies of Engås and Godø (1989a) and Ramm and Xiao (1995), increased bridle length in our study produced a small, but significant, reduction in wing spread (Table 1). If this change produced a change in footrope contact and net efficiency, then it would lead to a biased estimate of herding efficiency (net efficiency is explicitly assumed to be independent of bridle length in Eq. 4). Other possible changes in trawl geometry are that 1) at the longest length the upper bridle may sag sufficiently to touch the bottom within $L_{off}$ and 2) at the shortest length the bridle tension may be sufficient to lift the wingtip off the bottom. With such changes, and perhaps even without, $L_{off}$ may not be constant at all bridle lengths. Although we examined the performance of the trawl along the entire distance from the wing tips to the doors at the standard bridle length, we had insufficient ship-time to examine the performance at all bridle lengths to verify that the bridles performed as intended. We recommend that further studies of herding efficiency include additional research to verify that standard trawl performance is maintained for all experimental configurations. One approach for doing this is to simultaneously estimate escapement under the footrope, perhaps by using an auxiliary net attached beneath the trawl net as in Engås and Godø (1989b), so that the assumed independence of net efficiency and bridle length could be tested.

Fourth, we have conducted these experiments in relatively small areas that do not encompass all of the bottom conditions found over the areas covered by the bottom trawl surveys; therefore the results are potentially biased and certainly less variable than if they were conducted over such a range of conditions.

Bridle efficiency is only one component needed to determine trawl efficiency (i.e. the proportion of fish within the door path that are caught). The remaining component, for flatfish at least, is net efficiency (i.e. the proportion of fish within the net path that are caught). If estimates of net efficiency are also available, perhaps obtained with the use of an auxiliary net to capture escaping fish (Engås and Godø, 1989b), they can be included with estimates of bridle efficiency in Dickson's (1993a) model of the trawl fishing process to produce estimates of trawl efficiency. Such estimates could then be used to convert relative indices of fish abundance from trawl survey to absolute estimates of abundance.

## Acknowledgments

We thank Asgeir Aglen, Martin Dorn, Olav Rune Godø, Jack Turnock, and Gary Stauffer for reviewing the manuscript and for making helpful suggestions.

## Literature cited

Armistead, C. E., and D. G. Nichol.
1993. 1990 bottom trawl survey of the eastern Bering Sea continental shelf. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-7, 190 p.

Bridger, J. P.
1969. The behaviour of demersal fish in the path of a trawl. FAO Fish. Rep. 62(3):695–715.

Dickson, W.
1993a. Estimation of the capture efficiency of trawl gear. I: Development of a theoretical model. Fish. Res. 16: 239–253.
1993b. Estimation of the capture efficiency of trawl gear. II: Testing a theoretical model. Fish. Res. 16:255–272.

Efron, B., and R. Tibshirani.
1993. An introduction to the bootstrap. Chapman and Hall, New York, NY, 436 p.

Engås, A., and O. R. Godø.
1989a. The effect of different sweep lengths on the length composition of bottom-sampling trawl catches. J. Cons. Int. Explor. Mer 45:263–268.
1989b. Escape of fish under the fishing line of a Norwegian sampling trawl and its influence on survey results. J. Cons. Int. Explor. Mer 45:269–276.

Foster, J. J., C. M. Campbell, and G. C. W. Sabin.
1981. The fish catching process relevant to trawls. Can. Spec. Publ. Fish. Aquat. Sci. 58:229–246.

Fuwa, S.
1989. Fish herding model by ground ropes considering reaction of fish. Nippon Suisan Gakkaishi 55:1767–1771.

Fuwa, S., O. Sato, K. Nashimoto, and N. Higo.
1988. Fish herding model by ground rope. Nippon Suisan Gakkaishi 54:1155–1159.

Hemmings, C. C.
1969. Observations on the behaviour of fish during capture by the Danish seine net and their relation to herding by trawl bridles. FAO Fish. Rep. 62:645–655.

High, W. L.
1969. Scuba diving, a valuable tool for investigating the behavior of fish within the influence of fishing gear. FAO Fish. Rep. 62(3):323–331.

Hilborn, R., and M. Mangel.
1997. The ecological detective: confronting models with data. Princeton Univ. Press, Princeton, NJ, 315 p.

Main, J., and G. I. Sangster.
1981a. A study of the sand clouds produced by trawl boards

and their possible effect on fish capture. Scott. Fish. Res. Rep. 20:1–20.

1981b. A study of the fish capture process in a bottom trawl by direct observations from a towed underwater vehicle. Scott. Fish. Res. Rep. 23:1–23.

Martyshevskii, V. N., and V. N. Korotkov.

1969. Fish behaviour in the area of the trawl as studied by bathyplane. FAO Fish. Rep. 62(3):781–791.

Ramm, D. C., and Y. Xiao.

1995. Herding in groundfish and effective pathwidth of trawls. Fish. Res. 24:243–259.

Seber, G. A. F., and C. J. Wild.

1989. Nonlinear regression. Wiley, New York, NY, 768 p.

Venables, W. N., and B. D. Ripley.

1994. Modern applied statistics with S-plus. Springer-Verlag, New York, NY, 462 p.

# Age and growth of red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico off Louisiana*

**Charles A. Wilson**
Coastal Fisheries Institute
and
Department of Oceanography and Coastal Sciences
School of the Coast and Environment
Louisiana State University
Baton Rouge, Louisiana 70803-7503
E-mail address: cwilson@lsu.edu

**David L. Nieland**
Coastal Fisheries Institute
School of the Coast and Environment
Louisiana State University
Baton Rouge, Louisiana 70803-7503

**Abstract**–The red snapper (*Lutjanus campechanus*) is currently under rigorous federal and state management in the Gulf of Mexico because of apparent overfishing. Management strategies implemented to promote recovery of the species are dependent upon knowledge of various demographic variables, including the ages of individuals, the distribution of these ages (cohort strength) within the population, and maximum longevity. Thus a dependable and precise aging method is of principal importance. Counts of annuli in otolith thin sections have been used to age many species of fish, including red snapper. However, the utility of this method for aging red snapper has been questioned by those who dispute both the apparent longevity (over 50 yr) of red snapper and the position of the first annulus within the red snapper otolith.

We counted annuli and assessed edge condition in sagittal otoliths of 3791 red snapper collected from the northern Gulf of Mexico off Louisiana during the periods from 1989 to 1992 and from 1995 to 1998. Opaque annuli were validated by marginal increment analysis to form once per year from December through June. Estimated ages ranged from 0.5 to 52.6 yr for individuals from 104 mm to 1039 mm total length and from 0.02 kg to 22.79 kg total weight. Among the 2546 specimens of known sex, both sexes evidenced rapid growth of individuals until about age 8–10 yr, after which growth slows considerably. Von Bertalanffy growth models for total length at age were significantly different for males and females, indicating differential growth between the sexes, with females typically obtaining larger sizes at older ages than do males.

Manuscript accepted 15 March 2001.
Fish. Bull. 99:653–664 (2001).

The red snapper, *Lutjanus campechanus* (Poey) (family: Lutjanidae), is resident on the continental shelves of the Gulf of Mexico (GOM) and northwest Atlantic Ocean from the Bay of Campeche, Mexico, to Massachusetts; however, it is found only occasionally north of Cape Hatteras, North Carolina (Rivas, 1966; Robins and Ray, 1986; Hoese and Moore, 1998). Although Rivas (1966) suggested that red snapper may also occur off Bermuda, the Bahamas, and northern Cuba, it has been reported neither from Bermuda by Smith-Vaniz et al. (1999) nor from the Bahamas by Böhlke and Chaplin (1993), nor from Cuba by Allen (1985). The species is replaced in the Caribbean Sea and southward by the Caribbean red snapper, *L. purpureus* (Rivas, 1966; Robins and Ray, 1986; Hoese and Moore, 1998). Although the appellation "red snapper" has been widely used to identify as many as 12 commercially marketed lutjanids (Camber, 1955; Carpenter, 1965), it is correctly applied only to *L. campechanus* (Robins et al., 1991). The binomina *L. aya, L. blackfordi,* and *Neomaenis aya*, all of which appear in the literature (e.g. Moseley, 1966) are synonyms of *L. campechanus* and refer to the red snapper *sensu* Robins et al. (1991).

The red snapper has been and remains a significant component of both the commercial and recreational fisheries in the GOM. However, documented commercial landings from United States territorial waters declined precipitously from historic highs of about 6389 metric tons (t) in 1965 to 1015 t in 1991; estimated recreational landings similarly waned from 4734 t in 1979 to 581 t in 1990 (Schirripa and Legault[1]). Since 1991 both fisheries have been constrained by size limits, creel or trip limits, and quotas as established by the Gulf of Mexico Fishery Management Council (GMFMC). The best efforts of the GMFMC and the commercial and recreational sectors notwithstanding, overfishing of red snapper in the GOM may persist (Schirripa and Legault[2]).

* Contribution LSU-CFI-00-01 of the Coastal Fisheries Institute, Louisiana State University, Baton Rouge, LA 70803-7503.

[1] Schirripa, M. J., and C. M. Legault. 1997. Status of the red snapper in U. S. waters of the Gulf of Mexico: Updated through 1996. Contribution rep. MIA-97/98-05 from Sustainable Fisheries Division, Miami Laboratory, Southeast Fisheries Science Center, National Marine Fisheries Service, 75 Virginia Beach Drive, Miami, FL 33149-1099. [Not available from NTIS.]

[2] Schirripa, M. J., and C. M. Legault. 1999. Status of the red snapper in U. S. waters of the Gulf of Mexico: Updated through 1998. Contribution rep. SFD-99/00-75 from Sustainable Fisheries Division, Miami Lab-

*continued*

Accurate information on the age structure of the red snapper populations in the GOM is essential for monitoring year-class strength, for conducting stock assessments, and for documenting population recovery. Previous efforts at estimating red snapper age have employed a variety of aging methods. Bradley and Bryan (1975) cited the long spawning season and a constant recruitment into the population as reasons for the difficulty in assigning red snapper ages from length-frequency data. Moseley (1966) used scale annuli to age red snapper to age 4 yr and advanced spawning as the causal factor in check formation. Among 240 red snapper taken off the west coast of Florida, Futch and Bruger (1976) estimated red snapper ages of 1 to 5 yr from 200 readable whole otoliths; however, they postulated ages up to 20 yr for larger individuals whose otoliths were unreadable. Comparisons of ages derived from whole otoliths, scales, and vertebrae by Bortone and Hollingsworth (1980) revealed all three hard parts to be of equal utility in aging red snapper at age 1 and 2 yr. Wade (1981) also used scales to age red snapper to 9 yr. Nelson and Manooch (1982) reported red snapper ages from 1 to 16 yr based on both scales and sectioned otoliths and demonstrated once yearly scale annulus formation in June and July from monthly mean marginal growth. A recent study of red snapper otoliths significantly extended the hypothesized longevity of red snapper in the GOM to 42 yr (Szedlmayer and Shipp, 1994). Render (1995) provided a preliminary validation of yearly opaque annulus formation in sagittal otoliths and reported ages from 0 to 53 yr for red snapper in Louisiana waters. Examinations of otolith sections from 537 red snapper captured in the northwestern Atlantic Ocean from Beaufort, North Carolina, south to the Florida Keys manifested a maximum longevity of 25 yr (Manooch and Potts, 1997). Among 907 red snapper from the GOM off Alabama, Patterson (1999) reported opaque annulus formation from January through June and maximum ages of 30 yr for females and 31 yr for males. Despite these efforts, the longevity of red snapper remains controversial. Small sample sizes, a paucity of older specimens, and the failure to present legitimate validations of ages from hard parts (Beamish and McFarlane, 1983) have variously hampered the above studies. It has further been speculated that larger and presumably older red snapper form numerous false annuli within otoliths (Rothschild et al.[3]). And both the timing of deposition and the position of the putative first annulus remain in question.

Otolith analyses have proven consistent in estimating ages of many fish species, including several from the temperate waters of the northern GOM (Johnson et al., 1983; Barger, 1985; Beckman et al., 1988; Beckman et al., 1990, 1991; Crabtree et al., 1992; Murphy and Taylor, 1994; Franks et al., 1998; Thompson et al., 1998). Herein we present our interpretations of the use of sagittal otoliths to estimate ages of red snapper from the GOM off Louisiana. Specifically we address the timing of formation and position of the first annulus, validation of the once yearly accretion of opaque annuli, longevity, and reader reproducibility. We further describe the growth of red snapper with von Bertalanffy growth models for both males and females.

## Methods and materials

Red snapper from recreational and commercial catches were sampled from 1989 to 1992 and from 1995 to 1998 by personnel of the Louisiana State University (LSU) Coastal Fisheries Institute and the Louisiana Department of Wildlife and Fisheries (LDWF). Although the vast majority of our sampling efforts were targeted at both wholesale facilities and charter boat docks located in Grand Isle and Port Fourchon, LA, the area of coverage in the northern GOM extended from off the Mississippi River Delta in the east to off Galveston, TX, in the west. Morphometric measurements (total length [TL] or fork length [FL] in mm, total weight [TW] or eviscerated body weight [BW], i.e. mass with liver, digestive tract, and reproductive organs removed, in g) were recorded, both sagittal otoliths were removed, and sex was determined, when possible, for each specimen. Eviscerated body weight was converted to TW, when necessary, with the equation $TW=1.101(BW)-26.32$ (linear regression, df=418; $P<0.001$; $r^2=0.996$) and TL was estimated from FL with the equation $TL=1.073\ (FL)+3.56$ (linear regression, df=1015; $P<0.001$; $r^2=0.999$).

All undamaged, intact otoliths were weighed to the nearer 0.1 g. The left sagittal otolith from each individual was embedded in an epoxy resin and subsequently sectioned with a low-speed saw equipped with a wafering blade as described in Beckman et al. (1988). In those instances where the left sagitta was damaged or unavailable, the right sagitta was substituted. Examinations of otolith sections were made with a compound microscope and transmitted light at 40× to 100× magnification. Counts of annuli (opaque zones) were accomplished by reading along the medial surface of the transverse section dorsal or ventral to the sulcus; annuli were often inconsistent in other regions of the otolith section. Annuli were counted by two readers without knowledge of date of capture or morphometric data. The appearance of the otolith margin was also coded as either opaque or translucent (Beckman et al., 1988). Sections were recounted a second time by both readers when initial counts disagreed. Rather than excluding the small number of individuals for which a consensus could not be reached after a second reading ($n=27$), the assigned annulus count for these was that of the more experienced reader (reader 1). Reader 2's annulus count and edge condition were used in those circumstances where reader 1's were missing ($n=2$). Annulus counting error between the two readers was evaluated after both the initial and second readings of the otolith sections. Reproducibility of the resultant age estimates was evaluated with the coefficient of variation, the index of precision (Chang, 1982), and average percent error (Beamish and Fournier, 1981).

[2] (continued) oratory, Southeast Fisheries Science Center, National Marine Fisheries Service, 75 Virginia Beach Drive, Miami, FL 33149-1099. [Not available from NTIS.]

[3] Rothschild, B. J., A. F. Sharov, and A. Y. Bobyrev. 1997. Red snapper stock assessment and management for the Gulf of Mexico. Report submitted to the National Marine Fisheries Service, Office of Science and Technology, 1315 East-West Highway, Silver Spring, MD 20910 USA. [Not available from NTIS.]

The periodicity of opaque annulus formation was determined by marginal increment analysis and by plotting the proportion of otoliths with opaque margins by month of capture (Beckman et al., 1988). To assess the possibility of false annulus formation among either younger or older red snapper, those individuals of age ≤5 yr and those >5 yr were also analyzed as above. Fork lengths at 100% maturity, 420 mm for males and 440 mm for females (Render, 1995), are achieved at about age 5. If one opaque and one translucent zone are shown to be formed each year, validation of annuli as being accreted once yearly is accomplished.

Ages of red snapper were estimated from opaque annulus counts and date of capture with the equation

$$Age(\text{days}) = -182 + (annulus\ count \times 365) + ((m-1)\times 30) + d,$$

where $m$ = the ordinal number (1–12) of the month of capture; and
$d$ = the ordinal number (1–31) of the day of the month of capture.

The 182 days that were subtracted from each age estimate are an accommodation for the uniform 1 July hatching date which was assigned for all specimens (Render, 1995; Collins et al., 1996). Age in yr was derived by dividing the age in days by 365. Thus a red snapper captured on 1 January which exhibits five opaque annuli (including an opaque margin [see below]) in its otoliths would have an estimated age of 1644 days or 4.5 yr. Our age estimation method also assumed that opaque annulus formation at the otolith margin uniformly commenced in January. The small number of individuals captured in September, October, November, and December that evidenced early formation of opaque annuli had their ages adjusted by subtracting 365 days from their age estimates. Conversely, a larger number of individuals captured in January, February, and March had otoliths with translucent margins—evidence of an assumed delay in opaque annulus formation. The age estimates of these were augmented by the addition of 365 days.

Total length-TW regressions were fitted with linear regression to the model $TL = a\ TW^b$ from $\log^{10}$-transformed data. Male and female regressions were compared with analysis of covariance (SAS, 1985). Only those red snapper for which sex could be determined were used to fit growth models. Von Bertalanffy growth models of TL at age were fitted with nonlinear regression by least squares (SAS, 1985) in the form

$$TL_t = L_\infty\left(1 - e^{-k(t-t_0)}\right),$$

where $TL_t$ = $TL$ at age $t$;
$L_\infty$ = the $TL$ asymptote;
$k$ = a growth coefficient;
$t$ = age in yr; and
$t_0$ = a hypothetical age when $TL$ is zero.

Growth models were generated for three groups of red snapper within which the age and TL of all individuals were extant: 1) all specimens, 2) all specimens of known sex, and 3) specimens of known sex for which growth models were fitted independently for each sex. Likelihood ratio tests (Cerrato, 1990) were used to test for differences between males and females, both in growth models and in growth parameter estimates. Significance level for statistical analyses was 0.05 unless indicated otherwise.

## Results

During eight years of variable collection effort, 3791 red snapper from recreational ($n$=274) and commercial ($n$=3517) catches were sampled for morphometric data and sagittal otoliths. Among the 1438 male and 1542 female specimens for which sex could be determined, females ranged from 242 to 1039 mm TL and from 0.16 to 22.79 kg TW; males were 245–946 mm TL and 0.19–13.70 kg TW. Composite ranges for all specimens of either known or unknown sex were 104–1039 mm TL and 0.02–22.79 kg TW; however, 67.6% of 3787 available TL were between 325 and 525 mm and 80.0% of 3718 available TW were less then 2.5 kg (Fig. 1). Neither the slopes (df=2,932; $F$=3.41; $P$<0.065) nor the intercepts (df=2,932; $F$=3.16; $P$<0.075) of the TL-TW regressions were found to differ significantly between males and females; thus data for the two sexes were combined and a single predictive equation was generated

$$TW = 1.17 \times 10^{-8}(TL)^{3.04}$$

Sagittae of red snapper are ovate, laterally compressed, and have an indented sulcus acousticus on the proximal surface (Fig. 2A). Although one can count purported annuli in relatively small whole otoliths of red snapper less than age 5 (Futch and Bruger, 1976), it is difficult to discern annuli in the larger otoliths of older individuals. Thin transverse sections of these older otoliths showed semidistinct translucent and opaque annuli that alternated from the core to the growing edge (Fig. 2, B and C). The presumptive first annulus posed the most consistent problem for the readers. This annulus appeared as a diffuse "smudge" of opaque material variously located from totally isolated and somewhat distant from the core (Fig. 2B) to contiguous to and continuous with the otolith core (Fig. 2C).

Annulus counts ranging from 0 to 53 and edge conditions were determined by at least one reader for all 3791 individuals sampled. Reader 2 considered all the otolith sections to be of sufficient quality to produce annulus counts; reader 1 provided annulus counts from all but two sections. After the initial counts, consensus between readers was achieved for 2804 individuals A second reading of the 987 sections for which annulus counts differed produced consensus for 3762 individuals. The degree of agreement in red snapper opaque annulus counts between the two readers in each of the two readings was assessed. Average percent error (APE), coefficient of variation (CV), index of precision (D), and percentages of absolute differences in counts are given in Table 1.




**Figure 1**

Frequency histograms for (**A**) total length in mm ($n$=3787) and (**B**) total weight in kg ($n$=3718) for red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico off Louisiana. Specimens collected from the recreational and commercial fisheries from 1989 to 1992 and from 1995 to 1998. The vertical arrow denotes the rod and reel world record red snapper caught on 23 June 1996.

Proportions of otoliths with opaque margins were plotted by month of capture for all individuals ($n$=3791), for those individuals presumed to be sexually immature (ages less than or equal to 5, $n$=2143), and for those from individuals of presumptive sexual maturity (ages greater than 5, $n$=948). Each of the three plots (Fig. 3) features a single broad peak and a single broad valley and conclusively demonstrates opaque annulus formation from December through June and translucent annulus formation from July through November. Thus, the assumption of one to one correspondence between opaque annulus counts and estimated red snapper age in years is validated. Furthermore, this correspondence is validated for immature and mature individuals of all ages.

Having demonstrated once yearly accretion of opaque annuli, we estimated ages from 0.5 to 52.6 yr from the annulus counts of the red snapper in our study. The vast majority of specimens examined were ages 2–5 and only 1.2% of the total number were greater than age 15 yr (Fig. 4). The few ages greater than 15 yr, which were not represented in our sample, were 24, 28, 31, 34, 39, 40, 42–46, and 49–50. The otolith section from the oldest specimen examined is shown in Figure 5.

Among 3787 individuals, the single von Bertalanffy growth model which best describes red snapper TL at age was

$$TL\ (\text{mm}) = 941\ [1 - e^{-0.18\,(t + 0.55)}], \qquad (r^2 = 0.72).$$




**Figure 2**

Medial view (**A**) and transverse sections (**B** and **C**) of red snapper (*Lutjanus campechanus*) left sagittal otoliths. Abbreviations for all are Do = dorsal, V = ventral, P = posterior, A = anterior, SA = sulcus acousticus, Di = distal, and Pr = proximal. Arrows with numbers indicate positions of opaque annuli and their enumeration.

Not surprisingly, the von Bertalanffy growth model generated for all individuals of known sex ($n$=2979) was quite similar to the above:

$$TL\ (\text{mm}) = 935\ [1 - e^{-0.18\,(t\,+\,0.54)}], \qquad (r^2=0.72).$$

However, a likelihood ratio test revealed that von Bertalanffy growth models for males ($n$=1438) and females ($n$=1541) were significantly different from one another ($\chi^2$=75.09; df=1,2979; $P$<0.0001). The resultant models for TL at age were

$$\textit{Female } TL\ (\text{mm}) = 977\ [1 - e^{-0.16\,(t\,+\,0.63)}], \qquad (r^2=0.71)$$
$$\textit{Male } TL\ (\text{mm}) = 904\ [1 - e^{-0.19\,(t\,+\,0.48)}], \qquad (r^2=0.73)$$




**Figure 3**

Marginal increment analysis of red snapper (*Lutjanus campechanus*) otoliths for specimens of all ages ($n$=3791), for ages >5 ($n$=2143), and for ages >5 ($n$=948).

Predicted TL at age for males and females (Figs. 6 and 7) generated with the above equations illustrated rapid and roughly equivalent growth to an age of approximately 8–10 yr after which the growth curves diverge. Differential growth between males and females was further demonstrated with likelihood ratio tests which indicated significant differences in $L_\infty$ ($\chi^2$=13.05; df=1,2979; $P$<0.0003) and $k$ ($\chi^2$=7.16; df=1,2979; $P$<0.0075). No significant difference was detected in $t_0$ ($\chi^2$=1.34; df=1,2979; $P$<0.247). Owing to the large variability in observed TL at age (Fig. 6), TL was a poor estimator of red snapper age within the range of TL encountered in our sampling efforts.

**Table 1**

Average percent error (APE), coefficient of variation (CV), and index of precision (D) differences in red snapper (*Lutjanus campechanus*) otolith annulus counts for two readers on first and second readings. $n$ = 3791.

| | 1st reading | 2nd reading |
|---|---|---|
| APE | 3.736 | 0.0934 |
| CV | 0.045 | 0.0011 |
| D | 0.032 | 0.0008 |
| 0 | 73.96% | 99.29% |
| ±1 | 23.27% | 0.61% |
| ±2 | 2.24% | 0.08% |
| ≥ ±3 | 0.53% | 0.02% |

## Discussion

Otolith annuli as indicators of age in years have been validated for many freshwater and marine fish species, including the Australian lutjanids *L. adetii* and *L. quinquelineatus* (Newman et al., 1996) and *L. argentimaculatus*, *L. bohar*, *L.carponotatus*, *L. erythropterus*, *L. gibbus*, *L. johnii*, *L. malabaricus*, *L. monostigma*, *L. rivulatus*, *L. sebae*, and *L. vitta* (Cappo et al., 2000). Previous studies of red snapper have used scales (Moseley, 1966; Wade, 1981), otoliths (Futch and Bruger, 1976; Szedlmayer and Shipp, 1994; Render, 1995; Manooch and Potts, 1997; Patterson, 1999), scales and otoliths (Nelson and Manooch, 1982) and scales, otoliths, and vertebrae (Bortone and Hollingsworth, 1980) to estimate ages. Among these, early attempts to validate age estimation from circuli of scales and annuli of otoliths have suffered from two shortcomings: 1) a small sample size and 2) a paucity of individuals over age 10 yr. Nevertheless, they have produced a general consensus




**Figure 4**

Age-frequency histogram ($n$=3791) for red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico. Specimens were collected from the recreational and commercial fisheries of the Gulf of Mexico off Louisiana from 1989 to 1992 and from 1995 to 1998.

that transparent annuli (Nelson and Manooch, 1982) are formed during the spawning season, May to September in the GOM (Collins et al., 1996). Our validation of opaque annulus formation in otoliths of red snapper during the winter and spring seasons is in substantial agreement with previous efforts. Given that yearly formation of opaque annuli has been validated for substantial numbers of red snapper from the Atlantic waters off North Carolina south to Florida (Manooch and Potts, 1997) and the GOM waters off Alabama (Patterson, 1999) and Louisiana (Render, 1995; our study), and the validation among Australian congeneric species cited above, the one-to-one correspondence between annuli and age in years should be indisputable.

Certainly, the reproducibility statistics indicate that the annuli of red snapper otoliths are more difficult to count than those in otoliths of some other species. Comparisons of between-reader age estimates in several species of the family Sciaenidae have yielded almost 100% agreement (Beckman et al., 1988; Beckman et al., 1990; Barbieri et al., 1993; Lowerre-Barbieri et al., 1995). Sciaenid otoliths are comparatively massive and annuli are especially well defined. Conversely, red snapper otoliths are relatively thin and fragile and the annuli become less well defined with increasing age. But, even given the above, a first reading followed by a second reading produced consensus in age estimates for 99.29% of those red snapper considered in our study. Patterson (1999) reported 93.8% between-reader consensus of red snapper annulus counts after two readings. Quite unlike the situation in sciaenids, training and experience are critical to achieving high between-reader consensus on red snapper annulus counts.

The variable position and the diffuse appearance of the first annulus formed during the first winter following hatching (age approximately 6 months) are presumed to be functions of both the protracted red snapper spawning season and the rapid growth rate of juvenile red snapper. Those individuals that are spawned early in the season will experience proportionally more growth (and presumably more translucent zone accretion adjacent to the otolith core) than will a late spawned individual before opaque annulus accretion begins during the following winter; thus the first opaque annulus will be more distant from the otolith core in the former instance than under the latter circumstance. Also with the first opaque annulus accreting at a rate theoretically corresponding to the rapid growth rate experienced during the juvenile stage, the resulting first annulus is broader and more diffuse in appearance than annuli produced during times of reduced growth rates in later life. A more complete understanding of first annulus formation in red snapper could improve insights into both recruitment patterns and growth rates of individuals within a spawning season.

The distribution of ages among our sample population (Fig. 4) is certainly not reflective of the age distribution of red snapper in the GOM off Louisiana. Age-0 and age-1 specimens have been largely unavailable to our sampling efforts owing to minimum size limits applied to the recreational and commercial fisheries during the 1990s. Also




**Figure 5**

Cross section of sagittal otolith from the oldest red snapper, *Lutjanus campechanus*, from among 3791 specimens sampled from the recreational and commercial fisheries of the Gulf of Mexico off Louisiana from 1989 to 1992 and from 1995 to 1998. Abbreviations are Do = dorsal, V = ventral, SA = sulcus acousticus, Di = distal, and Pr = proximal. Arrows with numbers indicate positions of opaque annuli and their enumeration.

the dominance of ages 2–5 may reflect the practices of the fishermen who target red snapper and a migratory aspect of the species' life history. Age-0, and to a lesser extent age-1, red snapper are known to inhabit shallow water areas devoid of complex habitats or vertical relief where some are vulnerable to capture in trawls. This behavior is illustrated in fishery-independent trawl data from the GOM, specifically the Fall Groundfish Survey and the Summer SEAMAP Survey, in which the great majority of red snapper captured are age 0 and 1 (Schirripa and Legault[2]). It has been hypothesized that the disappearance of red snapper from the trawl data at age 1 represents their migration to structures such as oil and gas platforms that presumably provide refuge from large predators (Render, 1995). It is during this residence at the numerous oil and gas platforms off Louisiana that red snapper become vulnerable to fishing gear. Because the platforms are easily located and potentially harbor large populations of red snapper and other fish species (Stanley and Wilson, 1996, 1998), they are the preferred destinations for both commercial and recreational fishermen. The very low numbers of individuals of age >6 in our sample population likely result from both removal from the population through fishing and natural mortalities and emigration away from the oil and gas platforms to alternative habitats where they are less susceptible to capture.

It is difficult to compare the maximum observed red snapper longevity reported in our study with those reported in earlier studies (Moseley, 1966; Futch and Bruger, 1976; Wade, 1981; Nelson and Manooch, 1982; Render, 1995; Szedlmayer and Shipp, 1994; Patterson, 1999) because of the assortment of aging techniques (scales, whole or sectioned otoliths, length frequencies) and the variety of sources (commercial, recreational, or both) used. All show a predominance of relatively young individuals (<10 yr). However, recent advances and refinements in otolith preparation technology have allowed red snapper to be aged reliably up to the following ages: 42 yr (Szedlmayer and Shipp, 1994), 53 yr (Render, 1995), 31 yr (Patterson, 1999), and 52 yr (our study). Despite the sparsity of old red snapper among these research efforts, there can be little doubt that red snapper at least have the potential to achieve ages of 40–50 yr and more.

The red snapper growth models that we present are similar to those of earlier studies (Nelson and Manooch, 1982; Szedlmayer and Shipp, 1994; Manooch and Potts, 1997; Patterson, 1999) which did not produce separate models for the two sexes and variously applied weighted




**Figure 6**

Observed total length (mm) at age and relationship of age to total length predicted from von Bertalanffy growth models for male ($n$=1438) and female ($n$=1542) red snapper, *Lutjanus campechanus*, from the northern Gulf of Mexico. Closed circles and narrow line represent males; open squares and thick line represent females.

or unweighted analyses (Fig. 7). All models predict rapid, and very much similar, growth during the first 8–10 years of life and slower growth thereafter. Asymptotic lengths among the above varied from 936 to 1025 mm. Our von Bertalanffy growth models also predicted a greater asymptotic TL and slightly faster growth for female red snapper. Among marine teleosts from the GOM, a similar pattern of growth has been shown for red drum, *Sciaenops ocellatus* (Beckman et al., 1988), sheepshead, *Archosargus probatocephalus* (Beckman et al., 1991), and cobia, *Rachycentron canadum* (Franks et al., 1998). However, this phenomenon in red snapper may be the result of a preponderance of data from the commercial fishery included in our analyses. Owing to minimum size limits enforced in both the commercial and recreational fisheries, we had access to few red snapper less than age 2 and few less than 250 mm TL. Conversely, the preference of commercial red snapper fishermen and wholesalers for smaller, plate-size individuals afforded us little opportunity to sample larger, and presumably older, red snapper; these are the individuals that can influence estimation of $k$ and which ultimately drive estimation of $L_\infty$. The addition of another 20–30 old specimens of both genders could have profound effects on the estimations of $L_\infty$ for both sexes. Furthermore, growth studies of lutjanids in Australian waters (Davis and West, 1992; McPherson and Squire, 1992; Newman et al., 1996) report faster growth and larger size at age among males. Proportionally greater expenditures of energy in the production of gametes by females is advanced to explain this observation (Newman et al., 1996). Thus, although the growth rates and asymptotic lengths for male and female red snapper are shown to differ statistically in our study, questions of the biological veracity and the biological significance of these differences remain unresolved.

In addition to the differences in red snapper estimated growth rates between the sexes, there is an obvious high degree of diversity in individual growth rates. Owing to the large variability in age at a given TL (Fig. 6), this variable is a poor estimator of red snapper age. Our data indicate that red snappers of 400 mm, 600 mm, and 800 mm TL could be ages 2–7 yr, 3–9 yr, and 5–35+ yr, respectively. As a more concrete example, consider the International Game Fishing Association world-record red snapper caught by rod and reel, the otoliths of which were given to us for age analysis. This individual was caught off the coast of Louisiana by Doc Kennedy of Grand Isle, LA, on 23 June 1996; it was 22.79 kg (50 lb, 4 oz) TW, 1039 mm (40.9 in) TL, and 965 mm (38 in) FL. Given the immense size of this specimen, one would reasonably expect it to be ancient by red snapper standards. However, our analysis




**Figure 7**

Comparative von Bertalanffy growth models for red snapper, *Lutjanus campechanus*, from the western north Atlantic Ocean. S&S 94 = Szedlmayer and Shipp (1994), P 99 = Patterson (1999), N&M 82 = Nelson and Manooch (1982), and M&P 97 = Manooch and Potts (1997). Males and Females are from our study.

revealed it to be only 19.98 yr. Conversely, the two oldest red snapper we encountered, age 52.63 and 51.73 yr, were a comparatively small 851 mm TL and 862 mm TL, respectively, and 7.886 kg TW and 9.188 kg TW, respectively. A similar pattern was noted by Patterson (1999) among the red snapper that he sampled from the GOM off Alabama.

Personnel at the LSU Coastal Fisheries Institute continue to investigate the nuances of deriving red snapper ages from sagittal otoliths. Although our marginal increment analysis demonstrates that a single opaque increment is formed each year, our sample size among older individuals, albeit larger than any previous investigation, is probably inadequate for absolute validation of this phenomenon. Thus, some have and will continue to question once yearly annulus accretion among red snapper older than 20 yr. A solution for this problem may lie in radiometric aging techniques with protocols that analyze various radionuclides in the otoliths. Also, core-to-first-annulus measurements made on otolith sections from age-0 and age-1 individuals would contribute to a better understanding of when and how the first annulus is accreted.

## Acknowledgments

We appreciate the able assistance of Ed Moss (LDWF), Bruce Thompson, Scott Baker, and many others in our red snapper sampling efforts. To Louise Stanley (reader 1) and Andrew Fischer (reader 2), we offer our gratitude not only for assistance in sampling, but also for the many long hours at the microscope counting annuli. Jane and Lonnie Black of USA Fish, Golden Meadow and Cameron, LA, graciously allowed us access to their facilities. We also thank Joey Trosclair and the employees of Trosclair Canning Company, Cameron, LA, for their help and the many commercial and recreational fishermen who allowed us to sample their catches. Michael Schirripa, National Marine Fisheries Service, is gratefully recognized for his many tangible contributions to our red snapper research. The comments of Richard Shaw, Brigitte Nieland, and three anonymous reviewers contributed greatly to the manuscript. Funding for this research was provided by the U. S. Department of Commerce Marine Fisheries Initiative (MARFIN) Program (grant numbers NA90AA-H-MF762, NA57FF0287, and NA77FF0544). Finally, we take great pride in dedicating this paper to the memory of Jeffery H. Render, our friend and colleague, for his pioneering efforts in the study of red snapper life history in the northern GOM.

## Literature cited

Allen, G. R.
1985. FAO species catalogue. Vol. 6: Snappers of the world: an annotated and illustrated catalogue of lutjanid species known to date, 208 p. FAO, Rome.

Barbieri, L. R., M. E. Chittenden Jr, and C. M. Jones.
1993. Age, growth, and mortality of Atlantic croaker, *Micropogonias undulatus*, in the Chesapeake Bay region, with a discussion of apparent geographic changes in population dynamics. Fish. Bull. 92:1–12.

Barger, L. E.
1985. Age and growth of Atlantic croakers in the northern Gulf of Mexico, based on otolith sections. Trans. Am. Fish. Soc. 111:847–850.

Bradley, E., and C. E. Bryan.
1975. Life history and fishery of the red snapper (*Lutjanus campechanus*) in the northwestern Gulf of Mexico. Proc. Gulf Caribb. Fish. Inst. 27:77–106

Beamish, R. J., and D. A. Fournier.
1981. A method for comparing the precision of a set of age determinations. Can. J. Fish. Aquat. Sci. 38:982–983.

Beamish, R. J., and G. A. McFarlane.
1983. The forgotten requirement for age validation in fisheries biology. Trans. Am. Fish. Soc. 112:735–743.

Beckman, D. W., C. A. Wilson, and A. L. Stanley.
1988. Age and growth of red drum, *Sciaenops ocellatus*, from offshore waters of the northern Gulf of Mexico. Fish. Bull. 87:17–28.

Beckman, D. W., A. L. Stanley, J. H. Render, and C. A. Wilson.
1990. Age and growth of black drum in Louisiana waters of the Gulf of Mexico. Trans. Am. Fish. Soc. 119:537–544.
1991. Age and growth-rate estimation of sheepshead *Archosargus probatocephalus* in Louisiana waters using otoliths. Fish. Bull. 89:1–8.

Böhlke, J. E., and C. C. G. Chaplin.
1993. Fishes of the Bahamas and adjacent tropical waters. Univ. Texas Press, Austin, TX, 771 p.

Bortone, S. A., and C. L. Hollingsworth.
1980. Aging red snapper, *Lutjanus campechanus*, with otoliths, scales, and vertebrae. Northeast Gulf Sci. 4:60–63.

Camber, C. I.
1955. A survey of the red snapper fishery of the Gulf of Mexico, with special reference to the Campeche Banks. Fla. Board Conserv. Tech. Ser. No. 12:1–64.

Cappo, M., P. Eden, S. J. Newman, and S. Robertson.
2000. A new approach to validation of periodicity and timing of opaque zone formation in the otoliths of eleven species of *Lutjanus* from the central Great Barrier Reef. Fish. Bull. 98:474–488.

Carpenter, J. S.
1965. A review of the Gulf of Mexico red snapper fishery. U.S. Dep. Interior, Fish and Wildl. Serv., Bur. Comm. Fish. Circ. 208:1–35.

Cerrato, R. M.
1990. Interpretable statistical tests for growth comparisons using parameters in the von Bertalanffy equation. Can. J. Fish. Aquat. Sci. 47:1416–1426.

Chang, W. B.
1982. A statistical method for evaluating the reproducibility of age determination. Can. J. Fish. Aquat. Sci. 39: 1208–1210.

Collins, L. A., A. G. Johnson, and C. P. Keim.
1996. Spawning and annual fecundity of the red snapper (*Lutjanus campechanus*) from the northeastern Gulf of Mexico. *In* Biology, fisheries and culture of tropical groupers and snappers (F. Arreguin-Sanchez, J. L. Munro, M. C. Balgos, and D. Pauly, eds.), p. 174–188. ICLARM Conf. Proc. 48.

Crabtree, R. E., E. C. Cyr, R. E. Bishop, L. M. Falkenstein, and J. M. Dean.
1992. Age and growth of tarpon, *Megalops atlanticus*, larvae in the eastern Gulf of Mexico, with notes on relative abundance and probable spawning areas. Env. Biol. Fishes 35: 361–370.

Davis, T. L. O., and G. J. West.
1992. Growth and mortality of *Lutjanus vittus* (Quoy and Gaimard) from the North West Shelf of Australia. Fish. Bull. 90:395–404.

Franks, J. S., J. R. Warren, and M. V. Buchanan.
1998. Age and growth of cobia, *Rachycentron canadum*, from the northeastern Gulf of Mexico. Fish. Bull. 97:459–471.

Futch, R. B., and G. E. Bruger.
1976. Age, growth, and reproduction of red snapper in Florida waters. *In* Proceedings: colloquium on snapper-grouper fishery resources of the western central Atlantic Ocean (H. R. Bullis Jr. and A. C. Jones, eds.), p. 165–183. Florida Sea Grant College Program Report 17.

Hoese, H. D., and R. H. Moore.
1998. Fishes of the Gulf of Mexico. Texas A&M Univ. Press, College Station, TX, 422 p.

Johnson, A. G., W. A. Fable Jr., M. L. Williams, and L. E. Barger.
1983. Age, growth, and mortality of king mackerel, *Scomberomorus cavalla*, from the southeastern United States. Fish. Bull. 81:97–106.

Lowerre-Barbieri, S. K., M. E. Chittenden Jr., and L. R. Barbieri.
1995. Age and growth of weakfish, *Cynoscion regalis*, in the Chesapeake Bay region with a discussion of historical changes in maximum size. Fish. Bull. 93:643–656.

Manooch, C. S., III, and J. C. Potts.
1997. Age and growth of red snapper, *Lutjanus campechanus*, Lutjanidae, collected along the southeastern United States from North Carolina through the east coast of Florida. J. Elisha Mitchell Sci. Soc. 113:111–122.

McPherson, G. R., and L. Squire.
1992. Age and growth of three dominant *Lutjanus* species of the Great Barrier Reef inter-reef fishery. Asian Fish. Sci. 5:25–36.

Moseley, F. N.
1966. Biology of red snapper, *Lutjanus aya* Bloch, of the northwestern Gulf of Mexico. Publ. Inst. Mar. Sci., Univ. Texas 11:90–101.

Murphy, M. D., and R. G. Taylor.
1994. Age, growth, and mortality of spotted seatrout in Florida waters. Trans. Am. Fish. Soc. 123:482–497.

Nelson, R. S., and C. S. Manooch III.
1982. Growth and mortality of red snappers in the west-central Atlantic Ocean and northern Gulf of Mexico. Trans. Am. Fish. Soc. 111:465–475.

Newman, S. J., D. McB. Williams, and G. R. Russ.
1996. Age validation, growth and mortality rates of the tropical snappers (Pisces : Lutjanidae) *Lutjanus adetii* (Castelnau, 1873) and *L. quinquelineatus* (Bloch, 1790) from the central Great Barrier Reef, Australia. Mar. Freshwater Res. 47:575–584.

Patterson, W. F., III.
1999. Aspects of the population ecology of red snapper *Lutjanus campechanus* in an artificial reef area off Alabama. Ph.D. diss., Univ. South Alabama, Mobile, AL, 164 p.

Render, J. H.
1995. The life history (age, growth, and reproduction) of red snapper (*Lutjanus campechanus*) and its affinity for oil and gas platforms. Ph.D. diss., Louisiana State Univ., Baton Rouge, LA, x + 76 p.

Rivas, L. R.
1966. Snappers of the western Atlantic. Commer. Fish. Rev. 32(1):41–44 .

Robins, C. R., and G. C. Ray.
1986. A field guide to Atlantic Coast fishes of North America. Houghton Mifflin Company, Boston, MA, 354 p. + 64 pl.
Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott.
1991. Common and scientific names of fishes from the United States and Canada, 5th edition. Am. Fish. Soc. Spec. Publ. 20, Bethesda, MD, 183 p.
SAS (Statistical Analysis System).
1985. SAS user's guide: statistics, version 5 ed. SAS Institute, Cary, NC, 956 p.
Smith-Vaniz, W. F., B. B. Collette, and B. E. Luckhurst.
1999. Fishes of Bermuda: history, zoogeography, annotated checklist, and identification keys. Allen Press Inc., Lawrence, KS, 424 p.
Stanley, D. R., and C. A. Wilson.
1996. Abundance of fishes associated with a petroleum platform as measured with dual-beam hydroacoustics. ICES J. Mar. Sci. 53:743–475.
1998. Spatial variation in fish density at three petroleum platforms as measured with dual-beam hydroacoustics. Gulf Mexico Sci. 16:73–82.
Szedlmayer, S. T., and R. L. Shipp.
1994. Movement and growth of red snapper, *Lutjanus campechanus*, from an artificial reef area in the northeastern Gulf of Mexico. Bull. Mar. Sci. 55(2-3):887–896.
Thompson, B. A., M. Beasley, and C. A. Wilson.
1998. Age distribution and growth of greater amberjack, *Seriola dumerili*, from the north-central Gulf of Mexico. Fish. Bull. 97:362–371.
Wade, C. W.
1981. Age and growth of spotted seatrout and red snapper in Alabama. Proc. Ann. Conf. S.E. Assoc. Fish & Wildl. Agencies 35:345–354.

# Stomach content analysis of cobia, *Rachycentron canadum*, from lower Chesapeake Bay*

**Michael D. Arendt**
School of Marine Science
College of William and Mary
Virginia Institute of Marine Science
Gloucester Point, Virginia 23062
Present address: Marine Resources Research Institute
South Carolina Department of Natural Resources Division
217 Fort Johnson Road
Charleston, South Carolina 29422-2559
E-mail address: arendtm@mrd.dnr.state.sc.us

**John E. Olney**
Department of Fisheries Science
School of Marine Science
College of William and Mary
Virginia Institute of Marine science
Gloucester Point, Virginia 23062

**Jon A. Lucy**
Sea Grant Marine Advisory Program
Virginia Institute of Marine Science
Glooucester Point, Virginia 23062

Cobia (*Rachycentron canadum*) is a migratory pelagic species that is found in tropical and subtropical seas of the world, except in the central and eastern Pacific Ocean. The United States ranks third in total commercial production of cobia, however, recreational landings generally exceed commercial landings by an order of magnitude (Shaffer and Nakamura, 1989). In the western Atlantic Ocean, cobia migrate to Chesapeake Bay in spring and summer to spawn, and the productive waters of the Bay are believed to constitute important foraging grounds (Joseph et al., 1964; Richards, 1967). Cobia are known to move to areas of high food abundance, particularly crustacean abundance (Darracott, 1977).

Recent feeding studies in the northern Gulf of Mexico and off North Carolina have reported geographic differences in cobia diet and have indicated that the relative importance of fishes versus crustaceans is variable and that cephalopods constitute the least significant prey items. In the northern Gulf of Mexico, Franks et al. (1996) reported that fish (primarily anchovies, *Anchoa* sp.) dominated (% index of relative importance [IRI], Pinkas et al., 1971) the diet of juvenile cobia (236–440 mm FL). Meyer and Franks (1996) reported crustaceans (primarily portunid crabs) occurred in 79.1% of stomachs and represented 77.6% of total prey items consumed by cobia (373–1,530 mm FL) in the northern Gulf of Mexico. Fish (primarily hardhead catfish, *Arius felis*, and American eel, *Anguilla rostrata*) increased in importance with increasing cobia size. Fish were found in 58.5% of all cobia stomachs (20.3% of total prey) but occurred in 84.4% of stomachs of cobia 1150–1530 mm FL (Meyer and Franks, 1996). In contrast, Smith (1995) observed decreased importance of teleosts in the diet of cobia (39–142 cm FL) in North Carolina. Elasmobranch fishes and portunid crabs dominated the diet of cobia >9 kg (Smith, 1995).

Tag-recapture data collected between 1995 and 1999 document localized movement of cobia within lower Chesapeake Bay during summer, as well as the return of individual cobia to specific locations or general regions of the lower Bay in subsequent summers.[1] Although Chesapeake Bay is an important destination for migrating cobia, feeding habits of cobia in the Bay have never been thoroughly examined. Our study documents cobia feeding habits in Chesapeake Bay and compares findings with similar cobia studies from North Carolina and the northern Gulf of Mexico.

## Methods

Cobia were sampled opportunistically at marinas and fishing tournaments in lower Chesapeake Bay between June and July 1997. Intact stomachs were removed by cutting above the cardiac sphincter (esophagus) and below the pyloric sphincter (large intestine). Stomachs were labeled, bagged, transported on ice to the VA Institute of Marine Science, and examined in relatively fresh condition. An incision was made along the longitudinal axis and the contents of stomachs were emptied onto a 500-µm mesh sieve for rinsing and sorting. Contents were blotted dry on paper towels before counts, displaced volumes (1-L graduated cylinder), and identifications to the lowest possible taxon were made. When possible, carapace widths (mm) of crabs were measured with calipers. An index of relative importance (IRI) for all prey items combined was calculated with the formula *(% Number + % Volume)* × *(% Frequency)*, as described by Pinkas et al. (1971) and subsequently used by Smith (1995), Meyer and Franks

* Contribution 2390 of the Virginia Institute of Marine Science, Gloucester Point, VA 23062.

[1] Annual reports. Virginia Game Fish Tagging Program, Marine Resources Commission, 968 Oriole Dr. South, Suite 102, Virginia Beach, VA 23451.

Manuscript accepted 13 April 2001.
Fish. Bull. 99:665–670 (2001).




**Figure 1**

Volume (mL) of portunid crab consumed in lower Chesapeake Bay as a function of cobia size.

(1996), and Franks et al. (1996). Frequency (*%F*) was the percent of all stomachs that contained food.

## Results and discussion

Stomach contents from 114 adult cobia (37–141 cm FL) were examined. Seventy-eight stomachs (68%) contained at least one identifiable, nonbait prey item. One species of bivalve, one species of hydroid, six species of crustacean, one elasmobranch, and 16 species of teleost were observed (Table 1). Mean volume of prey per stomach was 150.6 mL (range: 2–680 mL). Mean number of different prey per stomach was 1.9 species (range, 1–5 species). Blue crab (*Callinectes sapidus*) and lady crab (*Ovalipes ocellatus*) dominated the diet of cobia.

Index of relative importance (IRI) values for blue crab (5257) and lady crab (3665) were two orders of magnitude higher than IRI values for other prey items. Larger cobia consumed greater volumes of crab (Fig. 1). Additionally, larger cobia consumed larger (sublegal, <13 cm) blue crabs, although similar-size lady crabs were consumed by all sizes of cobia (Table 2). Atlantic croaker (*Micropogonias undulatus*), hogchoker (*Trinectus maculatus*), and fish and crab remains constituted the top food items volumetrically after portunid crabs. Although relatively high volumes were observed, IRI values for Atlantic croaker and hogchoker were low owing to infrequent occurrence of these items. High volume of crab remains was consistent with high volumes, counts, and frequency of occurrence of portunid crabs in cobia stomachs. Fish remains likely resulted from finfish bait (predominantly menhaden, *Brevoortia tyrannus*, and Atlantic croaker, *Micropogonias undulatus*). Crab and fish remains not identifiable to family and finfish bait were excluded from IRI calculations.

Consumption of blue crab was greatest along the western shore of the Bay and least at the mouth of the Bay. Conversely, consumption of lady crab was greatest at the mouth of the Bay and least along the eastern shore of the Bay (Fig. 2A). A greater percentage of lady crabs were consumed by male cobia than by female cobia, whereas a greater percentage of blue crabs were consumed by female cobia (Fig. 2B). IRIs for blue crab and lady crab were similar in June, but dramatically different in July. In July, IRI for blue crab was twice as high as that for lady crab. Prey availability and habitat utilization were likely responsible for determining location, sex, and within-season differences in composition of portunid crabs consumed by cobia. June samples were predominantly male cobia collected from the Bay mouth and eastern shore of the Bay, whereas July samples were predominantly female cobia collected from the western Bay shore.

Cobia feeding habits in lower Chesapeake Bay were more similar to feeding habits reported for cobia from North Carolina (Smith, 1995) than to feeding habits of cobia from the northern Gulf of Mexico (Franks et al., 1996; Meyer and Franks, 1996). Portunid crabs dominated the diet of cobia in Chesapeake Bay and North Carolina (Smith, 1995). Elasmobranch fishes were consumed exclusively by large cobia in Chesapeake Bay and North Carolina (Smith, 1995). In North Carolina (Smith, 1995), cobia fed on stingrays (*Dasyatis* sp.) and smooth dogfish (*Mustelus canis*). Cownose ray (*Rhinoptera bonasus*), previously unreported as a food item of cobia, was the only elasmobranch consumed by cobia in Chesapeake Bay. Cownose ray was observed from eight cobia stomachs; however, on-

**Table 1**

Index of relative importance (IRI) of prey items consumed by cobia ($n$=78) in lower Chesapeake Bay, June–July 1997. IRI was calculated as *(% Number + % Volume) × (% Frequency)*, as described in Smith (1995) and Meyer and Franks (1996). Frequency (*%F*) is the percent of stomachs that contained food. UnID = unidentified.

| | Frequency (stomachs) | *%F* | Number (counts) | *%N* | Volume (mL) | *%V* | IRI |
|---|---|---|---|---|---|---|---|
| Phylum Cnidaria | | | | | | | |
| Class Hydrozoa | | | | | | | |
| Family Sertulariidae | | | | | | | |
| *Sertularia* sp. | 1 | 1.28 | 1 | 0.14 | 8 | 0.07 | 0 |
| Phylum Mollusca | | | | | | | |
| Class Bivalva | | | | | | | |
| Family Mytilidea | | | | | | | |
| *Mytilus edulis* | 2 | 2.56 | 47 | 6.60 | 17 | 0.14 | 17 |
| Phylum Arthropoda | | | | | | | |
| Class Crustacea | | | | | | | |
| Family Squillidae | | | | | | | |
| *Squilla empusa* | 2 | 2.56 | 2 | 0.28 | 11 | 0.09 | 1 |
| Family Pagruidae | 3 | 3.85 | 3 | 0.42 | 5 | 0.04 | 2 |
| Family Portunidae | | | | | | | |
| *Callinectes sapidus* | 46 | 58.97 | 226 | 31.74 | 6736 | 57.31 | 5252 |
| *Ovalipes ocellatus* | 43 | 55.13 | 321 | 45.08 | 2505 | 21.31 | 3660 |
| Family Canceridae | | | | | | | |
| *Cancer* sp. | 2 | 2.56 | 2 | 0.28 | 26 | 0.22 | 1 |
| *Cancer irroratus* | 1 | 1.28 | 1 | 0.14 | 5 | 0.04 | 0 |
| Family Xanthidae | | | | | | | |
| *Neopanope* sp. | 1 | 1.28 | 3 | 0.42 | 5 | 0.04 | 1 |
| Phylum Chordata | | | | | | | |
| Class Chonrichthyes | | | | | | | |
| Family Myliobatidae | | | | | | | |
| *Rhinoptera bonasus* | 7 | 8.97 | 8 | 1.12 | 81 | 0.69 | 16 |
| Class Osteichthyes | | | | | | | |
| Family Clupeidae | | | | | | | |
| *Opisthonema oglinum* | 1 | 1.28 | 3 | 0.42 | 160 | 1.36 | 2 |
| *Brevoortia tyrannus* | 1 | 1.28 | 1 | 0.14 | 180 | 1.53 | 2 |
| UnID Clupeidae | 2 | 2.56 | 2 | 0.28 | 275 | 2.34 | 7 |
| Family Batrachoididae | | | | | | | |
| *Opsanus tau* | 1 | 1.28 | 1 | 0.14 | 185 | 1.57 | 2 |
| Family Ophidiidae | | | | | | | |
| *Ophidion marginatum* | 1 | 1.28 | 1 | 0.14 | 42 | 0.36 | 1 |
| *Ophidion* sp. | 3 | 3.85 | 4 | 0.56 | 47 | 0.40 | 4 |
| Family Sygnathidae | | | | | | | |
| *Syngnathus floridae* | 1 | 1.28 | 1 | 0.14 | 2 | 0.02 | 0 |
| *Syngnathus* sp. | 2 | 2.56 | 2 | 0.28 | 9 | 0.08 | 1 |
| *Hippocampus erectus* | 1 | 1.28 | 1 | 0.14 | 3 | 0.03 | 0 |
| *Hippocampus* sp. | 5 | 6.41 | 49 | 6.88 | 52 | 0.44 | 47 |
| Family Pomatomidae | | | | | | | |
| *Pomatomus saltatrix* | 1 | 1.28 | 1 | 0.14 | 180 | 1.53 | 2 |
| Family Sciaenidae | | | | | | | |
| *Micropogonias undulatus* | 3 | 3.85 | 3 | 0.42 | 663 | 5.64 | 23 |
| Family Uranoscopidae | | | | | | | |

*continued*

**Table 1 (continued)**

| | Frequency (stomachs) | %F | Number (counts) | %N | Volume (mL) | %V | IRI |
|---|---|---|---|---|---|---|---|
| Family Uranoscopidae (continued) | | | | | | | |
| *Astroscopus* sp. | 1 | 1.28 | 1 | 0.14 | 40 | 0.34 | 1 |
| UnID Pleuronectiformes | 3 | 3.85 | 4 | 0.56 | 5 | 0.04 | 2 |
| Family Bothidae | | | | | | | |
| *Paralichthyes dentatus* | 2 | 2.56 | 2 | 0.28 | 25 | 0.21 | 1 |
| *Scopthalamus aquosus* | 1 | 1.28 | 1 | 0.14 | 15 | 0.13 | 0 |
| UnID Bothidae | 1 | 1.28 | 1 | 0.14 | 2 | 0.02 | 0 |
| Family Soleidae | | | | | | | |
| *Trinectes maculatus* | 9 | 11.54 | 14 | 1.97 | 440 | 3.74 | 66 |
| UnID Soleidae | 3 | 3.85 | 3 | 0.42 | 15 | 0.13 | 2 |
| Family Diodontidae | | | | | | | |
| *Chilomycterus schoepfi* | 3 | 3.85 | 3 | 0.42 | 15 | 0.13 | 2 |
| Totals | | | 712 | | | | 11754 |

**Table 2**

Consumption of portunid crabs as a function of five cobia size classes in lower Chesapeake Bay.

| | Cobia size class (cm FL) | | | | |
|---|---|---|---|---|---|
| | <100 | 100–109.9 | 110–119.9 | 120–129.9 | >130 |
| *Callinectes sapidus* | | | | | |
| *n* | 9 | 16 | 12 | 23 | 42 |
| Mean carapace width (mm) | 55.2 | 72.3 | 75.0 | 116.7 | 94.7 |
| Range of carapace width (mm) | 33–86 | 42–120 | 45–120 | 42–120 | 35–120 |
| *Ovalipes ocellatus* | | | | | |
| *n* | 42 | 37 | 38 | 32 | 27 |
| Mean carapace width (mm) | 39.1 | 41.2 | 37.7 | 45.3 | 37.1 |
| Range of carapace width (mm) | 26–50 | 30–56 | 20–51 | 20–51 | 27–55 |

ly the jaw plates (22–49 mm width) remained; thus, the IRI for cownose ray is likely underestimated. Flatfishes and syngnathids represented the teleosts most frequently consumed by cobia in Chesapeake Bay and North Carolina. Hogchoker (*Trinectes maculatus*) was more frequently consumed by cobia in Chesapeake Bay (our study) than in North Carolina (Smith, 1995). Blackcheek tonguefish (*Symphurus plaigusa*) was regularly consumed by cobia in North Carolina (Smith, 1995) but was absent from cobia stomachs in Chesapeake Bay. In Chesapeake Bay, *Syngnathus* sp. (pipefish) and *Hippocampus* sp. (seahorse) were only important in the diet of smaller cobia, whereas in North Carolina, these fishes were important in the diet of all cobia (Smith, 1995).

Cobia in our study predominantly consumed benthic and epibenthic prey items, most notably portunid crabs. Feeding studies on other large predatory fishes in Chesapeake Bay during the summer months, such as bluefish, *Pomatomous saltatrix*, and weakfish, *Cynoscion regalis*, reveal that these species consume prey items associated predominantly with pelagic food webs (Hartman and Brandt, 1995a, 1995b). Striped bass, *Morone saxatalis*, a large predator present in Chesapeake Bay year-round, is also reported to feed predominantly on pelagic fishes (Hartman and Brandt, 1995a, 1995b; Walters, 1999). During the summer, portunid crabs were consumed by bluefish, weakfish, and striped bass in Chesapeake Bay; however, portunid crab consumption was typically less than 5% of total prey items present in the stomachs of these fishes (Hartman and Brandt, 1995a, 1995b; Walters, 1999). Red drum, *Sciaenops ocellatus*, a large predator that uses Chesapeake Bay between spring and fall is also reported to selectively consume portunid crabs in estuarine environments in the Gulf of Mexico (Scharf and Schlict, 2000); however, feeding habits of red drum in Chesapeake Bay have not been documented. Although




**Figure 2**

Consumption of blue crab (*Callinectes sapidus*) and lady crab (*Ovalipes ocellatus*) by cobia in lower Chesapeake Bay by location (**A**), sex (**B**), and month (June and July, **C**).

other large predators in Chesapeake Bay occasionally consume portunid crabs, cobia is the only large predator in Chesapeake Bay for which selective consumption of portunid crabs is documented.

## Acknowledgments

Stomachs were collected in conjunction with a study on cobia reproduction in Chesapeake Bay. We thank Susan Crute, Jason Romine, and Holly Simpkins for technical assistance. Donnie Wallace and Harry Johnson Jr. provided access to their facilities at Wallace's Bait and Tackle, Fox Hill, VA. We thank the recreational and commercial fishermen who allowed their catches to be sampled. This project was supported by grants (RF96-8, RF97-9) from the Virginia Marine Resources Commission upon recommendation by the Recreational Fishing Advisory Board.

## Literature cited

Darracott, A.
1977. Availability, morphometrics, feeding, and breeding activity of a multi-species, demersal fish stock of the western Indian Ocean. J. Fish. Biol. 10(1):1–16.

Franks, J. S., N. K. Garber, and J. R. Warren.
1996. Stomach contents of juvenile cobia, *Rachycentron canadum*, from the northern Gulf of Mexico. Fish Bull. 94(2):374–380.

Hartman, K. J., and S. B. Brandt.
1995a. Trophic resource partitioning, diets, and growth of sympatric estuarine predators. Trans. Am. Fish. Soc. 124:520–537.
1995b. Predatory demand and impact of striped bass, bluefish, and weakfish in the Chesapeake Bay: applications of bioenergetics models. Can. J. Fish. Aquat. Sci. 52:1667–1687.

Joseph, E. B., J. J. Norcross, and W. H. Massmann.
1964. Spawning of cobia, *Rachycentron canadum*, in the Chesapeake Bay area, with observations of juvenile specimens. Ches. Sci. 5:67–71.

Meyer, G. H., and J. S. Franks.
1996. Food of cobia, *Rachycentron canadum*, from the north-central Gulf of Mexico. Gulf Res. Rep. 9(3):161–167.

Pinkas, L., M. S. Oliphant, and I. L. K. Iverson.
1971. Food habits of albacore, bluefin tuna, and bonito in California waters. Calif. Dep. Fish Game Fish Bull. 152, 105 p.

Richards, C. E.
1967. Age, growth, and fecundity of the cobia, *Rachycentron canadum*, from Chesapeake Bay and adjacent mid-Atlantic waters. Trans. Am. Fish. Soc. 96(3):343–350.

Scharf, F. S., and K. K. Schlicht.
2000. Feeding habits of red drum (*Sciaenops ocellatus*) in Galveston Bay, Texas: Seasonal diet variation and predator-prey size relationships. Estuaries 23(1):128–139.

Shaffer, R. V., and E. L. Nakamura.
1989. Synopsis of biological data on the cobia, *Rachycentron canadum* (Pisces: Rachycentridae). U.S. Dep. Commer., NOAA Tech Rep NMFS 82 [FAO Fisheries Synopsis 153], 21 p.

Smith, J. W.
1995. Life history of cobia, *Rachycentron canadum* (Osteichthyes: Rachycentridae), in North Carolina waters. Brimleyana 23:1–23.

Walters, J. F., III.
1999. Diet composition and feeding habits of large striped bass (*Morone saxatilis*) in Chesapeake Bay. M.S. thesis, School of Marine Science, College of William and Mary, Gloucester Point, VA, 124 p.

# Preliminary genetic population structure of southern flounder, *Paralichthys lethostigma*, along the Atlantic Coast and Gulf of Mexico

**Ivonne R. Blandon**
**Rocky Ward**
Perry R. Bass Marine Fisheries Research Station
Coastal Fisheries Division
Texas Parks and Wildlife Department
Palacios, Texas 77465
E-mail address (for I. R. Blandon): ivonne.blandon@tpwd.state.tx.us

**Tim L. King**
U.S. Geological Survey-Biological Resource Division
Leetown Science Center
Aquatic Ecology Laboratory
1700 Leetown Road
Kearneysville, West Virginia 25430

**William J. Karel**
Perry R. Bass Marine Fisheries Research Station
Coastal Fisheries Division
Texas Parks and Wildlife Department
Palacios, Texas 77465

**James P. Monaghan Jr.**
North Carolina Division of Marine Fisheries
3441 Arendell Street
Morehead City, North Carolina 28557

Southern flounder, *Paralichthys lethostigma*, inhabit coastal waters from Albemarle Sound, North Carolina to the Baja Laguna Madre del Sur in northern Mexico, but they are apparently absent from southern Florida (Ginsburg, 1952). This species inhabits coastal bays, sounds, and lagoons from spring to fall and migrates offshore to spawn in late fall and winter (Stokes, 1977). Valuable sport and commercial fisheries for southern flounder exist in both the northern Gulf of Mexico (Warren et al.[1]; Robinson et al.[2]) and the western North Atlantic (Monaghan[3]).

Declines in southern flounder absolute abundance in some regions (e.g. Texas during the 1980s; Fuls and McEachron[4]) have prompted some management agencies to institute restrictions on recreational and commercial fisheries including reductions in bag limits and minimum size. Should these measures fail to recover this fishery, other measures may be considered by managers, including further restrictions on harvest, or artificial propagation and stocking, or both. Implementation of such enhancement programs requires that genetic surveys be conducted to determine genetic variability and stock structure of managed fish populations (King et al., 1995). Failure to understand underlying genetic structure prior to implementing stocking programs places the genetic resources of target species at risk (Allendorf et al., 1987) and may result in the reduction or loss of among-population variability and changes in within-population genetic characteristics. Genetic analyses of population structure may also provide insight into management options that do not require stocking (Nelson and Soulé, 1987).

The objective of our study was to characterize population structure of southern flounder in coastal regions of the northern Gulf of Mexico and northwestern Atlantic Ocean and to test the null hypothesis of no genetic differentiation within the region surveyed. If this hypothesis was rejected, a number of processes would operate to structure southern flounder population(s). Genetic differentiation in some nearshore organisms in the northern Gulf of Mexico (e.g. *Sciaenops ocellatus*; Gold and Richardson, 1999) has been explained as isolation by distance (Wright, 1943). This model describes a population structured by isolation caused by limited individual migration potential in relation to the size of the species' distribution (Kimura and Weis, 1964). The hypothesis of isolation by distance is supported when geographic distance and genetic distance are positively correlated. Alternatively, differentiation may arise as an adaptive response to localized environmental conditions (King and Zimmerman, 1993) or from the operation of physical barriers, such

[1] Warren, T. A., L. M. Green, and K. W. Spiller. 1994. Trends in finfish landings of sportboat anglers in Texas marine waters May 1974–May 1992, 259 p. Manage. Data Ser. 109, Texas Parks and Wildlife (TPW), Coastal Fish. Div., Austin, TX 78744.

[2] Robinson, L., P. Campbell, and L. Butler. 1995. Trends in Texas commercial fishery landings, 1972–1994, 133 p. Manage. Data Ser. 117, Texas Parks and Wildlife (TPW), Coastal Fish. Div., Austin, TX 78744.

[3] Monaghan, J. P., Jr. 1996. Migration of paralichthid flounders tagged in North Carolina. Study 2 *in* Life history aspects of selected marine recreational fishes in North Carolina, 44 p. Completion Rep. Grant F-43, Segments 1–5, North Carolina Division of Marine Fisheries, 3441 Arendell Street, Morehead City, NC 28557.

[4] Fuls, B., and L. W. McEachron. 1997. Trends in relative abundance and size of selected finfishes and shellfishes along the Texas coast: November 1975–December 1995, 108 p. Manage. Data Ser. 137, Texas Parks and Wildlife (TPW), Coastal Fish. Div., Austin, TX 78744.

Manuscript accepted 9 February 2001.
Fish. Bull. 99:671–678 (2001).




**Figure 1**

Collection sites for southern flounder in North American waters. LLM = Lower Laguna Madre; MAT = Matagorda Bay; GAL = Galveston Bay; SAB = Sabine Lake; MS = Mississippi; AL = Alabama; STA = St. Augustine, Florida; NC = North Carolina.

as current patterns (King et al., 1994). Support for competing models comes from examination of specific patterns observed in structured populations. Such patterns, if observed, may have important management implications.

## Materials and methods

Southern flounder were collected during the summers of 1996 and 1997 by rod and reel, flounder gigs, or Texas Parks and Wildlife (TPW) gill nets in four Texas bays (Sabine Lake, Galveston Bay, Matagorda Bay, and lower Laguna Madre), by pound nets in Core Sound, North Carolina, from gill nets in estuarine waters near Biloxi, Mississippi, and from commercial fish houses in Dauphin Island, Alabama, and St. Augustine, Florida (Fig. 1). Southern flounder from commercial fish houses were reported by the house operator to be caught locally. A majority of individuals collected were adult and were not reliably assignable to year classes. Samples were screened by using isoelectric focusing (IEF) of sarcoplasmic proteins (methods of Ward et al., 1995) to insure that individuals belonging to other *Paralichthys* species were not included in our analyses (necessary because of accidental inclusion of congenerics both in samples obtained from commercial sources and in samples of juvenile flounder obtained during routine resource sampling in Texas).

Skeletal muscle, liver, heart, and kidney tissues were excised from fresh or frozen fish. Sample preparation and electrophoretic techniques and conditions followed those of King and Pate (1992). Gel and electrode buffers used were tris-borate-EDTA, pH 8.0 (Selander et al., 1971), tris-citrate, pH 8.0. (Selander et al., 1971), lithium hydroxide, pH 8.0 (Selander et al., 1971), borate buffer, pH 9.0 (modified from Sackler, 1966), and Poulik's discontinuous system, pH 8.7 (Selander et al., 1971). Histochemical methods were primarily taken from Manchenko (1994). Genetic nomenclature followed the recommendations of Shaklee et al. (1990).

BIOSYS-1 (Swofford and Selander, 1981) was used to generate an allele frequency table, to estimate the proportion of loci heterozygous ($H$) in the average individual, the proportion of polymorphic loci in individuals from each bay population, and genetic divergence using the $F$-statistics of Wright (1978). Exact tests calculated by GENEPOP (v. 3.1; Raymond and Rousset, 1995) were used to test for conformance of genotypic frequencies at each locus within a sample to Hardy-Weinberg expectations, genotypic linkage disequilibrium, and allelic and genotypic heterogeneity. Pairwise differences between samples were tested by using the genic differentiation randomization test in GENEPOP. Results were combined over loci with Fisher's method (Sokal and Rohlf, 1994). Differences between each pair of populations were summarized by using the chord distance of Cavalli-Sforza and Edwards (1967). An unrooted phylogenetic tree was fitted to the chord distance matrix by using the neighbor-joining (NJ) algorithm. TreeView (Page, 1996) was used to visualize

the tree. The strength of support for each node in the tree was tested by bootstrapping over loci with NJBPOP (Cornuet et al., 1999). To further quantify spatial heterogeneity, the fixation index ($F_{ST}$) was calculated for each locus to provide measures of interpopulation differentiation and estimates of reduction in heterozygosity of a subpopulation due to population subdivision. A $\chi^2$ test was used to test the null hypothesis, $F_{ST}$=0 (Workman and Niswander, 1970). All $\chi^2$ probability values from tests for conformance to Hardy-Weinberg expectations, heterogeneity, linkage disequilibrium, and $F_{ST}$ were adjusted for multiple simultaneous tablewide tests by using sequential Bonferroni adjustments to minimize type-I statistical inference errors (Rice, 1989).

Partitioning of variance components among geographic regions and within samples was accomplished by using a hierarchical analysis of molecular variance (AMOVA, Cockerham, 1969, 1973) with the package ARLEQUIN (version 2, Schneider et al., 1999). Sample sites were nested into regional groupings for separate analyses (Atlantic versus Gulf of Mexico, and Atlantic combined with eastern Gulf sites versus western Gulf). A phenogram was generated from the chord distance matrix with the neighbor-joining (N-J) algorithm. The N-J phenogram, with bootstrap estimates (as percentage of 10,000 replications) obtained by resampling loci within samples, was generated with NJBPOP (Cornuet, et al., 1999). The significance of the relationship between genetic (i.e. chord) and geographic (bay to bay shoreline distance) distance matrices was determined by sampling the randomization distribution generated from 1000 replications with the MXCOMP (matrix comparison) routine in NTSYS-PC 2.0 by Rohlf (1997) to allow a Mantel test (Mantel, 1967). An assignment test (WHICHRUN 4.1, Banks and Eichert, 2000) tested the ability to discriminate population of origin based on an individual's multilocus genotypic profile.

Clinal trends in heterozygosity and allele frequency were examined by using nonparametric correlation analyses (SAS Institute, 1989). Significance of Spearman's correlation coefficients was determined as the probability a correlation differed from zero. Probabilities less than 0.05 were considered statistically significant (Snedecor and Cochran, 1980).

## Results

Misidentifications detected by IEF resulted in a reduced sample size in some samples especially from Alabama and Florida. Examination of 46 enzymes and structural protein systems in southern flounder produced scorable phenotypes for 68 putative gene loci. Two dimeric esterase loci (*ESTD-1** and *ESTD-2**, 3.1.1.. [IUBMBNC, 1992]) a tripeptide aminopeptidase locus (*PEPB-2**, 3.4.13..), glycerol-3-phosphate dehydrogenase (*G3PDH**, 1.1.1.8), two glucose-6-phosphate isomerase loci (*GPI-A** and *GPI-B**, 5.3.1.9), phosphoglucomutase (*PGM**, 5.4.2.2) and two glucose-6-phosphate dehydrogenase loci (*G6PDH-1** and *G6PDH-2**, 1.1.1.49) were resolved, scored as variable, and included in analyses. The remaining 59 loci were monomorphic or could not be scored consistently and were omitted. All polymorphic loci had the same common allele across all localities (Table 1). *ESTD-1**, *ESTD-2**, *GPI-B**, and *G6PD-1** each expressed an allele unique to a single locality. The percentage of polymorphic loci ($P_{0.99}$) averaged 7.5% and ranged from 4.41% to 10.29% (Table 1). Mean individual heterozygosity ranged from 0.03 (SE=0.02) in North Carolina and Florida to 0.12 (SE=0.12) in Matagorda Bay. Statistically significant clinal relationships were found in the Gulf of Mexico between the frequency of the common allele of the *G6PDH-2** locus and degrees west longitude ($r_s$=–0.829, $P$<0.05) and degrees north latitude ($r_s$=0.829, $P$<0.05). Further differentiation of populations in the western Gulf of Mexico was observed at the *G3PDH** locus. The *G3PDH*1* allele occurred at a frequency of 12.0% in the Laguna Madre, declined to a frequency of 1.4% in Galveston Bay, and was absent in all collections east of Galveston Bay. Four loci had alleles confined to a single locality in this same region of Gulf of Mexico (Table 1), including *G6PDH-1*2* which was limited to the Laguna Madre. Samples were in Hardy-Weinberg equilibrium at the nine allozyme loci surveyed for each of the 72 possible comparisons of variable loci for each locality, except *ESTD-2** in Galveston Bay. No statistically significant genotypic linkage disequilibrium was observed between any loci for any population.

Tests for homogeneity of allelic frequencies at variable loci across all localities were statistically significant at six of the nine loci: *ESTD-2** ($P$<0.01), *G3PDH** ($P$<0.01), *GPI-A** ($P$=0.01), *GPI-B** ($P$<0.01), *PGM** ($P$<0.01), and *G6PDH*-2* ($P$<0.01). Tests for homogeneity of genotypic frequencies across all localities were significant at four loci: *G3PDH** ($P$<0.01), *GPI-B** ($P$<0.01), *PGM** ($P$<0.01), and *G6PDH-2** ($P$<0.01). $F_{ST}$ averaged 0.088 and varied from 0.181 for *G6PDH-2** to 0.001 for *G6PDH-1**. All estimates of $F_{ST}$ were found to be statistically different from 0, except for loci *G6PDH-1**, *PEPB-2**, and *GPI-A**.

Regional differentiation among southern flounder was demonstrated (Fig. 2) by using chord distance and N-J clustering. The greatest discontinuity was between all samples from Galveston Bay eastward and a cluster composed of Matagorda Bay and the Laguna Madre. No apparent differentiation exists between Atlantic Coast populations and populations collected from the eastern Gulf of Mexico. This pattern is supported by the outcome of the assignment test (Table 2). Percentage correct assignment ranged from 0% for Florida to 78% for Matagorda Bay. However, when assignments between groups identified in the cluster analysis were examined, the overall correct percentage increased to 81%. The assignment test supports the interpretation of the cluster analysis, suggesting within-region differentiation was minimal and not geographically consistent, but between-region differences were considerable. A different pattern was discerned by using AMOVA (Table 3). The majority of the variance (>99%) was distributed within samples and the among-sites-within-coasts component was nonsignificant. Comparison of Atlantic versus Gulf of Mexico sample sites was statistically significant ($P$=0.04) and comparison of western Gulf of Mexico sample sites with those from the east-

**Table 1**

Allele frequencies, sample sizes ($n$), percentage of polymorphic loci per population, and mean heterozygosity per locus. LM = Laguna Madre, Texas; MAT = Matagorda Bay, Texas; GAL = Galveston Bay, Texas; SAB = Sabine Lake, Texas; MIS = off Biloxi, Mississippi; ALA = Mobile Bay, Alabama; STA = off St. Augustine, Florida; NC = Core Sound, North Carolina.

| Locus | Allele | LM | MAT | GAL | SAB | MIS | ALA | STA | NC |
|---|---|---|---|---|---|---|---|---|---|
| *ESTD-1** [1,3] | *1 | 0.972 | 0.918 | 0.889 | 0.917 | 0.885 | 1.000 | 1.000 | 0.959 |
| | *2 | 0.028 | 0.082 | 0.111 | 0.083 | 0.115 | 0.000 | 0.000 | 0.027 |
| | *3 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.014 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 37 |
| *ESTD-2** [1,3] | *1 | 0.009 | 0.000 | 0.014 | 0.000 | 0.000 | 0.000 | 0.000 | 0.054 |
| | *2 | 0.981 | 1.000 | 0.917 | 1.000 | 1.000 | 1.000 | 1.000 | 0.946 |
| | *3 | 0.009 | 0.000 | 0.069 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 37 |
| *PEPB-2** [1,3] | *1 | 0.074 | 0.016 | 0.014 | 0.000 | 0.038 | 0.024 | 0.025 | 0.000 |
| | *2 | 0.926 | 0.984 | 0.986 | 0.958 | 0.962 | 0.976 | 0.975 | 1.000 |
| | *3 | 0.000 | 0.000 | 0.000 | 0.042 | 0.000 | 0.000 | 0.000 | 0.000 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 37 |
| *G3PDH** [2,3] | *1 | 0.120 | 0.098 | 0.014 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | *2 | 0.778 | 0.779 | 0.972 | 0.848 | 0.808 | 0.833 | 0.875 | 0.946 |
| | *3 | 0.102 | 0.123 | 0.014 | 0.152 | 0.192 | 0.167 | 0.125 | 0.054 |
| | $n$ | 54 | 61 | 36 | 23 | 13 | 21 | 20 | 37 |
| *GPI-A** [1,3] | *1 | 1.000 | 0.992 | 1.000 | 1.000 | 0.962 | 0.929 | 0.975 | 1.000 |
| | *2 | 0.000 | 0.008 | 0.000 | 0.000 | 0.038 | 0.071 | 0.025 | 0.000 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 36 |
| *GPI-B** [1,4] | *1 | 0.000 | 0.000 | 0.000 | 0.125 | 0.000 | 0.000 | 0.000 | 0.000 |
| | *2 | 1.000 | 1.000 | 0.986 | 0.875 | 1.000 | 1.000 | 1.000 | 1.000 |
| | *3 | 0.000 | 0.000 | 0.014 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 37 |
| *PGM** [1,4] | *1 | 0.009 | 0.008 | 0.086 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | *2 | 0.991 | 0.992 | 0.829 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |
| | *3 | 0.000 | 0.000 | 0.086 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | $n$ | 54 | 61 | 35 | 24 | 13 | 21 | 20 | 37 |
| *G6PDH-1** [1,3] | *1 | 0.991 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |
| | *2 | 0.009 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| | $n$ | 54 | 61 | 36 | 24 | 13 | 21 | 20 | 37 |
| *G6PDH-2** [1,3] | *1 | 0.000 | 0.000 | 0.114 | 0.000 | 0.000 | 0.095 | 0.025 | 0.000 |
| | *2 | 0.639 | 0.648 | 0.871 | 0.896 | 0.962 | 0.881 | 0.975 | 1.000 |
| | *3 | 0.361 | 0.352 | 0.014 | 0.104 | 0.038 | 0.024 | 0.000 | 0.000 |
| | $n$ | 36 | 61 | 35 | 24 | 13 | 21 | 20 | 37 |
| % polymorphic loci | | 10.29 | 8.82 | 10.29 | 7.35 | 7.35 | 5.88 | 5.88 | 4.41 |
| Heterozygosity | | 0.10 | 0.12 | 0.09 | 0.08 | 0.08 | 0.09 | 0.03 | 0.03 |
| Direct count (SE) | | (0.05) | (0.06) | (0.04) | (0.03) | (0.03) | (0.04) | (0.02) | (0.02) |

[1] Buffer = TVB (Tris-borate-EDTA, pH 8.0).
[2] Buffer = borate 9.
[3] Tissue = muscle.
[4] Tissue = liver.

ern Gulf and the Atlantic coast approached significance ($P$=0.07). The results of chi-square tests for population differentiation (Table 4) allowed a pairwise examination of differences among the eight sampling sites. Eastern sample sites were not significantly different ($P$>0.01) from their nearest geographic neighbors. Western sites, from Sabine Lake to the lower Laguna Madre, were statistically different from at least one neighbor, suggesting more pronounced genetic differentiation in the western Gulf of Mexico. Among Gulf of Mexico samples, no sta-

**Table 2**

Assignment test outcomes. Values indicate frequency of assignment of individuals from a collection locality (rows) to the locality to which it is most similar (columns). Numbers on the diagonal indicate correct assignments. The "East" category combines collections from North Carolina to Sabine Lake, Texas, and the "West" category combines Texas sites from Galveston Bay to Laguna Madre. NC = Core Sound, North Carolina; STA = off St. Augustine, Florida; ALA = Mobile Bay, Alabama; MIS = off Biloxi, Mississippi; SAB = Sabine Lake, Texas; GAL = Galveston Bay, Texas; MAT = Matagorda Bay, Texas; LM = Laguna Madre, Texas.

| | Assigned to | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| From | NC | STA | ALA | MIS | SAB | GAL | MAT | LM |
| NC | 29 | 0 | 4 | 0 | 2 | 0 | 0 | 0 |
| STA | 13 | 0 | 5 | 1 | 0 | 1 | 0 | 0 |
| ALA | 8 | 0 | 6 | 3 | 0 | 0 | 0 | 0 |
| MIS | 4 | 1 | 0 | 4 | 3 | 1 | 0 | 0 |
| SAB | 15 | 1 | 4 | 2 | 10 | 5 | 0 | 0 |
| GAL | 9 | 0 | 2 | 1 | 2 | 22 | 1 | 0 |
| MAT | 8 | 0 | 6 | 6 | 1 | 0 | 14 | 25 |
| LM | 7 | 0 | 4 | 0 | 0 | 1 | 3 | 19 |
| East | | | 158 | | | | | 1 |
| West | | | 33 | | | | | 61 |

tistically significant correlations were observed between chord distances and geographic distances ($r$=0.416, $t$=1.549, $P$=0.06).

## Discussion

Genetic differentiation was not extensive over most of the range of *P. lethostigma* examined in this study. Samples collected from Core Sound in North Carolina to Sabine Lake on the upper Texas coast were genetically similar. However, a discontinuity in allele frequencies was identified on the Texas coast between Galveston and Matagorda Bays. In addition, statistically significant clines in allele frequencies at the *G6PDH-2** locus and in average individual heterozygosity were observed across the Gulf of Mexico. These observations do not suggest the occurrence of independent stocks of southern flounder in the Gulf of Mexico but do support the hypothesis that genetic structuring is present. Southern flounder samples collected off St. Augustine, FL, and off North Carolina cluster with samples in the Gulf of Mexico from Galveston Bay, Texas, eastward, despite a modern-era distributional gap that encompasses the southern reaches of Florida from the Loxahatchee River on the Atlantic Coast to the Caloosatchee Estuary on the Gulf Coast (NOAA[5]). This apparent gap may not represent an effective barrier to gene flow or may be of such recent origin that differentiation has been minimal. It is also possible that differences existed that were undetected by techniques used in our study.

Significant correlation between genetic and geographic distance was not found, lending no support to application of the isolation by distance model (Wright, 1943) to pop-




**Figure 2**

Unrooted neighbor-joining tree depicting the underlying structure found in the pairwise Cavalli-Sforza and Edwards chord distance matrix for eight collections of southern flounder. Numbers indicate bootstrap support in 1000 replications and white ellipses indicate nodes with bootstrap support below 50%. LLM = Lower Laguna Madre; MAT = Matagorda Bay; GAL = Galveston Bay; SAB = Sabine Lake; MS = Mississippi; AL = Alabama; STA = St. Augustine, Florida; NC = North Carolina.

[5] NOAA. 1985. Gulf of Mexico coastal and ocean zones strategic assessment:data atlas, 188 p. Strat. Assess. Branch, Ocean Assess. Div., Off. Oceanography Mar. Assess., Nat. Ocean Serv. and the Southeast Fish. Center, NMFS, NOAA, U.S. Dep. Commer., 75 Virginia Beach Dr, Miami, FL 33149.

ulation structuring in southern flounder. The estimated $F_{ST}$ value of 0.088 for southern flounder was greater than that found for *Sciaenops ocellatus* ($F_{ST}$=0.022, Gold et al., 1994), *Cynoscion nebulosus* ($F_{ST}$=0.009, King and Pate, 1992), or *Pogonius chromis* ($F_{ST}$=0.013, Karel, unpubl. data) in this region. Physical or biotic factors may have caused greater isolation for southern flounder than for other nearshore species. Cluster analysis suggested that the region of greatest differentiation occurs along the middle Texas coast. Similar genetic structure in *Crassostrea virginica* may be explained by seasonal current patterns in Corpus Christi Bay, Texas (King et al., 1994). A comparable mechanism may operate in southern flounder; offshore currents may have resulted in reduced dispersion of eggs and larvae between the upper and middle Texas coasts. Currents off the Texas coast are seasonally variable and complex (Cochrane and Kelly, 1986) and may aid in reducing egg and larval dispersion between regions of the western Gulf of Mexico.

Many marine species spawn in the open ocean, have eggs and larvae with an extensive planktonic stage, or are highly mobile as adults. It is not surprising that such organisms are often panmictic, or exhibit subdivision on only broad levels (e.g. *Sciaenops ocellatus*; Gold et al., 1994). When genetic differentiation is found in marine organisms (e.g. *Cynoscion nebulosus*; King and Pate, 1992), extensive regions of clinal change in allele frequency may be seen and may be an adaptive feature (King and Zimmerman, 1993). A critical question for fishery management is how much differentiation is necessary to indicate biologically significant population structuring. Gold et al. (1994), for instance, found that red drum were subdivided (albeit weakly) between Atlantic and Gulf of Mexico subpopulations despite relatively high levels of gene flow between the populations and failure of cluster analyses to consistently segregate localities into proper geographic regions. However, significant differences in allele frequencies for two loci were found and the researchers were able to demonstrate, through hierarchical gene-diversity analysis, that 20% of the variation in gene diversity was related to within-region differences.

**Table 3**

Hierarchical analyses of molecular variation (AMOVA) among mtDNA composite haplotypes of southern flounder (*Paralichthys lethostigma*) from the U.S. Atlantic coast and Gulf of Mexico.

| Source of variation | Variance | % variation | $P^1$ |
|---|---|---|---|
| All sites | | | |
| Among coasts | 0.00026 | 0.06 | 0.04 |
| Among sites within coasts | –0.00020 | –0.04 | 0.92 |
| Within sites | 0.44985 | 99.99 | 0.74 |
| Eastern versus western cluster | | | |
| Among clusters | 0.00042 | 0.09 | 0.07 |
| Among sites within regions | –0.00034 | –0.08 | 0.10 |
| Within sites | 0.44985 | 99.98 | 0.76 |

[1] Probability of finding a more extreme variance component by chance alone (1000 permutations).

**Table 4**

Tests ($\chi^2$) of pairwise population differentiation for southern flounder (*Paralichthys lethostigma*) samples obtained from sites on the U.S. Gulf and Atlantic coasts based on variability at eight allozyme loci. The $\chi^2$ value tests the hypothesis of no difference in allelic distribution across collection localities. The associated probability ($P$) is given in parentheses. Inf. = infinite.

| | North Carolina | Florida | Alabama | Mississippi | Sabine | Galveston | Matagorda |
|---|---|---|---|---|---|---|---|
| Florida | 12.875 (0.38) | | | | | | |
| Alabama | 22.791 (<0.01) | 4.478 (0.81) | | | | | |
| Mississippi | 28.820 (<0.01) | 13.350 (0.21) | 7.723 (0.66) | | | | |
| Sabine | 26.087 (0.01) | 17.208 (0.14) | 18.942 (0.09) | 21.959 (0.04) | | | |
| Galveston | 59.146 (<0.01) | 40.903 (<0.01) | 42.314 (<0.01) | 37.970 (<0.01) | 44.407 (<0.01) | | |
| Matagorda | Inf. (<0.01) | Inf. (<0.01) | Inf. (<0.01) | Inf. (<0.01) | Inf. Inf. (<0.01) | (<0.01) | |
| Laguna Madre | Inf. (<0.01) | Inf. (<0.01) | Inf. (<0.01) | 42.178 (<0.01) | Inf. (<0.01) | Inf. (<0.01) | 26.956 (0.07) |

In our study, allele frequency discontinuities and clinal variation in genetic characters were identified among southern flounder inhabiting the western portion of the species' range. Short-term goals (e.g. supplementing exploited populations) that fail to account for this structuring could result in management programs that undermine the long-term resource management objective: maintaining the evolutionary potential of this species. The results of this survey suggest that southern flounder populations should be considered potentially distinct pending further resolution of population differentiation and clinal variation. Stocking efforts, especially those involving interbay transfers, should be undertaken only after careful consideration of all pertinent information, and be contingent upon careful studies of genetic variation at the appropriate local level. Observed discontinuities and clinal variations in allele frequencies may indicate adaptation to localized conditions and should be incorporated into comprehensive management strategies developed for southern flounder.

## Acknowledgments

Financial support for this research was received from the Sport Fish Restoration Program of the U. S. Fish and Wildlife Service. Field personnel of the TPW Coastal Fisheries Division provided invaluable assistance in sample collection. Additional samples were provided by Greg Stunz of Texas A&M University, and Bradley Randall and his colleagues in the Fishery Department of the Gulf Coast Research Laboratory of Ocean Springs, Mississippi. We thank Larry McEachron and Bob Colura for administrative support and assistance with sample collection, and Roberta Vickers, David Westbrook, and Eric Young for their laboratory assistance, as well as Andrew Shaw for the preparation of Figure 1.

## Literature cited

Allendorf, F. W., N. Ryman, and F. M. Utter.
1987. Genetics and fishery management: past, present, and future. *In* Population genetics and fishery management (N. Ryman and F. Utter, eds.), p. 1–19. Univ. Washington Press, Seattle, WA.

Banks, M. A., and W. Eichert.
2000. WHICHRUN (version 3.2) a computer program for population assignment of individuals based on multilocus genotype data. J. Hered. 91:87–89.

Cavalli-Sforza, L. L., and A. W. F. Edwards.
1967. Phylogenetic analysis: models and estimation procedures. Am. J. Hum. Gen. 19:233–257.

Cochrane, J. D., and F. J. Kelly.
1986. Low-frequency circulation on the Texas-Louisiana continental shelf. J. Geophys. Res. 91:10645–10659.

Cockerham, C. C.
1969. Variance of gene frequencies. Evolution 23:72–83.
1973. Analysis of gene frequencies. Genetics 74:679–700.

Cornuet, J.-M., S. Piry, G. Luikart, A. Estoup, and M. Solignac.
1999. New methods employing multilocus genotypes to select or exclude populations as origins of individuals. Genetics 153:1989–2000.

Ginsburg, I.
1952. Flounders of the genus *Paralichthys* and related genera in American waters. Fish. Bull. 52:267–351.

Gold, J. R., T. L. King, L. R. Richardson, D. H. Bohlmeyer, and G. C. Matlock.
1994. Genetic studies in marine fishes. VII: allozyme differentiation within and between red drum (*Sciaenops ocellatus*) from the Gulf of Mexico and Atlantic Ocean. J. Fish Biol. 116:517–185.

Gold, J. R., and L. R. Richardson.
1999. Population structure of two species targeted for marine stock enhancement in the Gulf of Mexico. Bull. Natl. Res. Inst. Aquacult. (suppl. 1):79–87.

IUBMBNC (International Union of Biochemistry and Molecular Biology, Nomenclature Committee).
1992. Enzyme nomenclature. Academic Press, Orlando, Florida, 862 p.

Kimura, M., and G. H. Weis.
1964. The stepping stone model of population structure and the decrease of genetic correlation with distance. Genetics 49:561–576.

King, T. L., and H. O. Pate.
1992. Population structure of spotted seatrout inhabiting the Texas Gulf coast: an allozymic perspective. Trans. Am. Fish. Soc. 121:746–756.

King, T. L., R. Ward, I. R. Blandon, R. L. Colura, and J. R. Gold.
1995. Using genetics in the design of red drum and spotted seatrout stocking programs in Texas: a review. *In* Uses and effects of cultured fishes in aquatic ecosystems (H. L. Schramm Jr. and R. G. Piper, eds.), p. 499–502. Am. Fish. Soc. Sym. 15, Am. Fish. Soc., Bethesda, MD.

King, T. L., R. Ward, and E. G. Zimmerman.
1994. Population structure of eastern oysters (*Crassostrea virginica*) inhabiting the Laguna Madre, Texas and adjacent bay systems. Can. J. Fish. Aquat. Sci. 51:215–222.

King, T. L., and E. G. Zimmerman.
1993. Clinal variation at aspartate aminotransferase-2 in spotted seatrout, *Cynoscion nebulosus* (Cuvier), inhabiting the north-western Gulf of Mexico. Anim. Gen. 24:59–61.

Manchenko, G. P.
1994. Handbook of detection of enzymes on electrophoretic gels. CRC Press, Boca Raton, FL, 341 p.

Mantel, N. A.
1967. The detection of disease clustering and a generalized regression approach. Cancer Res. 27:209–220.

Nelson, K., and M. Soulé.
1987. Genetical conservation of exploited fishes. *In* Population genetics and fishery management (N. Ryman and F. Utter, eds.), p. 345–368. Univ. Washington Press, Seattle, WA.

Page, D. D. M.
1996. TREEVIEW: an application to display phylogenetic trees on personal computers. Comput. Appl. Biosci. 12: 357–358.

Raymond, M., and F. Rousset.
1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenism. J. Heredity 86:248–249.

Rice, W. R.
1989. Analyzing tables of statistical tests. Evolution 43: 223–225.

Rohlf, F. J.
1997. NTSYS-pc: numerical taxonomy and multivariate analysis system, version 2.0. Exeter Software, Setauket, NY, 32 p.

Sackler, M. L.
1966. Xanthine oxidase from liver and duodenum of the rat:

histochemical localization and electrophoretic heterogeneity. J. Histochem. Cytochem. 14:326–333.

SAS Institute Inc.
1989. SAS/STAT user's guide, version 6, 4th ed., vol. 1. SAS Institute, Cary, NC, 890 p.

Schneider, S., D. Roessli, and L. Excoffier.
1999. ARLEQUIN ver. 2.0: a software for population genetic data analysis. Genetics and Biometry Laboratory, Univ. Geneva, Geneva, Switzerland.

Selander, R. K., M. H. Smith., S. Y. Yang, W. E. Johnson, and J. B. Gentry.
1971. Biochemical polymorphism and systematics in the genus *Peromyscus*. Variation in the old field mouse (*Peromyscus polionotus*). Studies in Genetics IV, Univ. Texas Publ. 7103:49–90.

Shaklee, J. B., F. W. Allendord, D. C. Morizot, and G. S. Whitt.
1990. Gene nomenclature for protein-coding loci in fish. Trans. Am. Fish. Soc. 119:2–15.

Snedecor, G. W., and W. G. Cochran.
1980. Statistical methods. Iowa State Univ. Press, Ames, IA, 524 p.

Sokal, R. R., and F. J. Rohlf.
1994. Biometry: the principles and practice of statistics in biological research, 3rd ed. W. H. Freeman and Co., New York, NY, 880 p.

Stokes, G. M.
1977. Life history studies of southern flounder (*Paralichthys lethostigma*) and Gulf flounder (*P. albigutta*) in the Aransas Bay areas of Texas. Texas Parks and Wildlife Dep. Tech. Ser. 25:1–37.

Swofford, D, L., and R. B. Selander.
1981. BIOSYS-1: a Fortran program for the comprehensive analysis of electrophoretic data in population genetics and systematics. J. Heredity 72:282–283.

Ward, R., I. R. Blandon, and B. W. Bumguardner.
1995. Hybridization among members of the genus *Morone* (Pisces: Percichthyidae) in Galveston Bay, Texas. Texas J. Sci. 47:155–158.

Workman, P. L., and J. D. Niswander.
1970. Population studies on southwestern Indian tribes II. Local genetic differentiation in the Papago. Am. J. Hum. Gen. 22:24–49.

Wright, S.
1943. Isolation by distance. Genetics 28:114–138.
1978. Evolution and the genetics of populations, vol. 4. Variability within and among natural populations. Univ. Chicago Press, Chicago, IL, 580 p.

# Age determined from the daily deposition of concentric rings on common octopus (*Octopus vulgaris*) beaks

**José L. Hernández-López**
**José J. Castro-Hernández**
Departamento de Biología
Universidad de Las Palmas de Gran Canaria
Apdo. 550, Las Palmas de Gran Canaria
Canary Islands, Spain
E-mail address (for J. J. Castro-Hernández, contact author): josejuan.castro@biologia.ulpgc.es

**Vicente Hernández-García**
Sea Fisheries Institute, Research Station
72-600 Swinoujscie, Poland

The common octopus (*Octopus vulgaris* Cuvier, 1797) is an Atlantic and Mediterranean species (Guerra, 1992; Mangold, 1998). It is one of the most important target species of the Northwest African fisheries (Hernández and Bas, 1993; Foucher et al., 1998). The common octopus catch reported for this area in 1994 was 137,844 t, representing 47.17% of the total world octopus catch. In 1996 it was 156,300 t, representing 50.03% of the total world octopus catch (FAO, 1998).

Octopus age and growth have been determined by laboratory rearing studies (Itami et al., 1963; Nixon, 1969; Mangold and Boletzky, 1973; Smale and Buchan, 1981; Villanueva, 1995) and by field studies (Guerra, 1979; Hatanaka 1979; Pereiro and Bravo de Laguna, 1979). Growth rates can be calculated for animals maintained in the laboratory, but comparison with growth under natural conditions is questionable (Mangold, 1983). In field studies, growth and age can be correlated when there is clear evidence that a single year class from a stable population is under consideration, but where the spawning season is very long as in the common octopus (Mangold, 1983; Guerra, 1992), identifying year classes is difficult (Guerra, 1979, Hatanaka, 1979).

Cephalopod age has been determined by several methods: Guerra (1979), Pereiro and Bravo de Laguna (1979), and others have reported growth and age correlations by following a single year class from a stable population. Concentric rings on statoliths (Young, 1960), the internal shell, and eye lenses (Gonçalves, 1993) of *Octopus vulgaris* have been reported. Raya and Hernández-González (1998) observed marks on the internal rostral area of beaks from common octopus, possibly related to daily growth.

None of these methods has been validated for known-age *Octopus vulgaris*. Furthermore, all require fairly complex methods for preparing structures prior to observation under the microscope (polishing, embedding in resin, and sectioning with a diamond, etc.) which hinders their application to field studies. This paper provides an easy new method of determining *Octopus vulgaris* age based on the upper beak microstructure, validated for the paralarval period.

## Material and methods

The study was carried out on 275 common octopus (164 males and 111 females) collected between January 1998 and May 1999, from catches of the small-scale fishery off the island of Gran Canaria (The Canary Islands, central-east Atlantic).

An additional sample of 27 *Octopus vulgaris* paralarvae was obtained from spawning females that deposited and incubated egg bunches in plastic burrows inside a 12,000-L tank. The embryonic development took between 25–30 days at a temperature range of 19–22°C. Once hatched, the paralarvae were transferred to transparent 12-L containers with open seawater flow, in July 1997 and June and July 1999, and reared in the laboratory at 19–22°C water temperature and natural photoperiod. The dates of hatching and death of each paralarva were recorded. The bottom was siphoned daily to remove dead individuals. During rearing, paralarvae were fed with recently hatched crab zoeae (see Hernández-García et al., 2000).

Ventral mantle length (VML) was measured in both benthic octopus and paralarvae to the nearest 0.1 mm. We used VML as the body measurement because we consider it to be more accurate than dorsal mantle length. Total body weight (TW) of benthic octopus was recorded to the nearest 0.01 g (to the nearest 0.0001 g in paralarvae). With the exception of the paralarvae, all the specimens were sexed.

The beak of each animal, including paralarvae, was removed and stored in 70% ethyl alcohol. Lower and upper beaks were sagittally sectioned with scissors to obtain two symmetrical half beaks (Fig. 1). The half beaks were cleaned with water and the mucus covering the inner part of the lateral walls was removed by rubbing it softly with the fingers (obviously, this operation was not necessary in the case of the paralarvae beaks).

By using a stereoscopic microscope, the concentric rings in the lateral wall of each beak were counted from the rostral tip area to the opposite end of the lateral wall. Because of the lack of pigmentation in paralarvae beaks, the concentric rings in their lateral wall were more easily counted with a microscope. Rings of each beak were

Manuscript accepted 10 April 2001.
Fish. Bull. 99:679–684 (2001).




**Figure 1**

Diagram of upper beak of *Octopus vulgaris* and lateral wall showing growth bands, rostral tip area, the distal posterior beak end, and counting line.

counted at least three times by the same person, and those with less than two identical counts were rejected from analysis.

The number of rings in beaks of paralarvae was compared with the number of days each one lived.

## Results

Octopus obtained from the small-scale fishery ranged in size from 4.8 to 165 mm VML, and weighed between 0.38 and 3926 g. Females ranged from 60 to 165 mm VML and weighed from 215 to 3926 g. Males ranged from 58 to 160 mm VML and from 200 to 3167 g in weight. We were unable to sex two individuals (4.8 mm VML, 0.38 g and 8.1 mm VML, 0.60 g). Paralarvae ranged from 1.0 to 2.7 mm VML and from 0.001 to 0.005 g in weight.

The internal lateral walls of upper beaks from 302 individuals (27 paralarvae and 275 individuals in benthic stages of *Octopus vulgaris*) revealed a pattern of concentric bands deposited from the rostral tip of the beak to the opposite margin of the lateral wall, parallel to the beak edges. Both halves of the upper beaks showed similar spatial and density patterns of microstructures. Lower beaks showed no regularity in the pattern of bands along the lateral walls and were discarded. On the upper beaks the distance from the rostral tip to the distal end of the lateral wall showed a positive correlation with the VML (Pearson's correlation: $r$=0.825; $P$<0.001) (Fig. 2).

Ring counts were more difficult near the rostral tip where rings were frequently discontinuous. Counts were easier to make near the edges of the lateral wall which was less highly pigmented.

Paralarvae survived in tanks from 3 to 26 days. For 48.1% of paralarvae, the concentric ring count in the lateral wall of the upper beak equaled the number of days that they lived. Otherwise, in 22.2% and 29.6% of paralarvae the number of rings counted were one more or one less, respectively, than the number of days of age. These data (Fig. 3) indicate that daily deposition of a growth increment in the lateral wall of the upper beak begins on day one after hatching. The weight-age and VML-age relationships of paralarvae (Fig. 4, A and B) were similar to those found by Villanueva (1995), with differences attributable to rearing conditions. Given the correlation between increment counts and age of paralarvae, we applied the upper beak ring count method for age determination of 272 common octopus, ranging from 4.8 to 165 mm VML. Results should be taken with caution for the benthic stages of octopus pending the validation of growth of adults and the frequency of rings deposition.

Increments counted on beaks from octopus collected in the wild ranged from 53 to 398 corresponding to individuals of 0.38 and 3926 g body weight, respectively (4.8 and 165 mm VML). Males and females had no difference in the number of rings counted in the lateral walls of upper beaks (ANOVA, $F$=0.0006, $P$=0.98). The age of males ranged between 3.2 and 12.3 months (95–369 rings), and females ranged between 3.1 and 13.3 months (93–398 rings), see Figure 5 (A and B) for weight-age and MVL-age relationships for benthic octopus.

## Discussion

In *Octopus vulgaris* and other shallow-water cephalopods, regular patterns of activity and evidence of endogenous rhythms induced by the light–dark cycles have been reported in both field and laboratory animals (Cobb et al., 1995). These endogenous rhythms may be reflected in a




**Figure 2**

Relationship of beak lateral wall length to ventral mantle length (VML) of benthic *Octopus vulgaris*.




**Figure 3**

Lateral wall ring counts versus real age in 27 paralarvae from 3 to 26 days old.

chitinous structure such as the beaks (Raya and Hernández-González, 1998) or in calcium deposits in statoliths. Statoliths are the hard structures most commonly used for cephalopod age estimation (Lipinski, 1986, 1993; Arkhipkin, 1993), although the presence of concentric rings in the internal shell, beaks, and eye lenses have also been used (Clarke, 1965; Gonçalves, 1993; Raya and Hernández-González, 1998). When beaks are used, erosion of the rostral area during the life of the animal may bias age determination toward underestimation (some of the first rings may be eroded and therefore not counted). We found evidence of incomplete increments on the edge of the lateral wall, near the rostral tip area; therefore, ages we provide for benthic adults are to be considered minimum estimates.

If rings on the lateral walls of the upper beaks are laid down daily and can be accurately counted even in the oldest specimens (as indicated by the pattern in paralarvae), then our results are consistent with a lifespan of 12–13 months in the Canary Island waters. Raya and Hernández-González (1998) gave a lifespan of 10–12 months for octopus caught off the coast of northwest Africa (21–26°N)




**Figure 4**

Plot of the number of rings counted on the lateral walls of the upper beak against (**A**) body weight and (**B**) ventral mantle length (VML) of paralarvae of *Octopus vulgaris*.

although they reported some heavier but younger specimens than we found. This difference could be due to discrepancies in the aging methods or, as in the case of Mangold (1983), areas off the coast may have different growth patterns and lifespans. Thus, Smale and Buchan (1981) proposed a lifespan of 9–12 months in females and 12–15 in males *Octopus vulgaris* from the South African coast.

Several authors have noted that size (and probably weight) may not reliably indicate age in field-caught cephalopods (Mangold and Boletzky, 1973; Hixon, 1980) because it may vary greatly depending upon factors such as food and temperature (Van Heukelem, 1979; Mangold, 1983). Cephalopods reveal great morphological variability with latitude attributed to environmental influences on development (Hernández-García and Castro, 1998), and probably on lifespan. The length and weight ranges of octopus caught off the Canary Islands are within the ranges reported for this species off East Africa, are the limits of range (upper and lower) off South Africa (Smale and Buchan, 1981) and in the western Mediterranean Sea (Mangold, 1983), although the range recorded in our study (Canary Islands) is closer to that reported for the Mediterranean Sea.

The smallest octopi that we examined from fishery catches were 4.8 and 8.1 mm VML (0.38 and 0.60 g TW, respectively), well outside the minimum commercial length (90–100 mm VML). Their estimated ages were 51 and 91 days old. In the English Channel, the planktonic phase for common octopus has been estimated at 3 months (Rees, 1950; Rees and Lumby, 1954); and average weight of 0.2 g at settling may be normal regardless of temperature (Mangold, 1983). *Octopus* typically spend the first 5–12 weeks of life as an active predator on plankton (Mangold, 1983); they change gradually from a planktonic to benthic life style (Boletzky, 1977) in some way dependent upon temperature (Mangold 1983). We did not observe marked differences in ring pattern spacing indicative of the transition between planktonic and benthic life styles. However, the distance between rings does change during the benthic phase of life—a feature that seems related to water temperature—the rings being larger than average during winter and smaller during summer.




**Figure 5**

Plot of the number of rings counted on the lateral walls of the upper beak against (**A**) body weight and (**B**) ventral mantle length (VML) of benthic *Octopus vulgaris*.

## Acknowledgments

We thank Ana Y. Martín-Gutiérrez for her assistance in data collection. This study was funded by the Secretariat for Fishing of the Autonomous Government of the Canary Islands (Spain).

## Literature cited

Arkhipkin, A.
1993. Age, growth, stock structure and migratory rate of pre-spawning short-finned squid *Illex argentinus* based on statolith ageing investigations. Fish. Res. 16(4):313–338.

Boletzky, S. v.
1977. Post-hatching behaviour and mode of life in cephalopods. Symp. Zool. Soc., Lond. 38:557–567.

Clarke, M. R.
1965. "Growth rings" in the beaks of the squid *Moroteuthis ingens* (Oegopsida: Onychoteuthidae). Malacologia 3(2):287–307.

Cobb, C. S., S. K. Pope, and R. Williamson.
1995. Circadian rhythms to light-dark cycles in the lesser octopus, *Eledone cirrhosa*. Mar. Freshwater Behav. Physiol. 26: 47–57.

FAO (Food and Agriculture Organization of the United Nations).
1998. Fishery statistics. Capture production 1996. FAO yearbook, vol. 82. FAO, Rome.

Foucher, E., M. Thiam, and M. Barry.
1998. A GIS for the management of fisheries in West Africa: preliminary application to the octopus stock in Senegal. *In* Cephalopod biodiversity, ecology and evolution (A. I. L. Payne, M. R. Lipinski, M. R. Clarke, and M. A. C. Roeleveld, eds.), p. 337–346. S. Afr. J. Mar. Sci. 20.

Gonçalves, J. M. A.
1993. *Octopus vulgaris* Cuvier, 1797 (polvo-comun): sinopse da biologia e exploração. Ms.C. thesis, Universidade dos Açores, Horta, Açores 470 p.

Guerra, A.
1979. Fitting a von Bertalanffy expression to *Octopus vulgaris* growth. Inv. Pesq. 43:319–327.
1992. Mollusca cephalopoda. Fauna Iberica, vol. 1. Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas, Madrid, 327 p.

Hatanaka, H.
1979. Studies on the fisheries biology of common octopus off northwest coast of Africa. Bull. Far Seas Fish. Res. Lab. 17: 13–124.

Hernández, V., and C. Bas.
1993. Análisis de la evolución de las tallas de los cefalópodos explotados en la costa del Sáhara (división 34.1.3 de CECAF) entre los períodos 1967–70 y 1989–90. Bol. Inst. Esp. Oceanogr. 9(1):215–225.

Hernández-García, V., and J. J. Castro.
1998. Morphological variability in *Illex coindetii* (Cephalopoda: Ommastrephidae) along the North-west coast of Africa. J. Mar. Biol. Assoc. U.K. 78:1259–1268.

Hernández-García, V., A. Y. Martín, and J. J. Castro-Hernández.
2000. Evidences of external digestion of crustaceans in *Octopus vulgaris* (Cuvier, 1797) paralarvae. J. Mar. Biol. Assoc. U.K. 80:559–560

Hixon, R. F.
1980. Growth, reproductive biology, distribution and abundance of three species of loliginid squid (Myopsida, Cephalopoda) in the northwest Gulf of Mexico. Ph.D. diss., Univ. Miami, Coral Gables, FL, 233 p.

Itami, K., Y. Izawa, S. Maeda, and K. Nakai.
1963. Notes on the laboratory culture of the Octopus larvae. Bull. Jap. Soc. Sci. Fish. 29:514–520.

Lipinski, M.
1986. Methods for the validation of squid age from statoliths. J. Mar. Biol. Assoc. U.K. 66(2):505–526.
1993. The deposition of statoliths: a working hypothesis. *In* Recent advances in cephalopod fisheries biology (T. Okutani, R. K. O'Dor, and T. Kubodera, eds.), p. 241–262. Tokai University Press, Tokyo.

Mangold, K.
1983. *Octopus vulgaris*. *In* Cephalopod life cycles, vol. I, species accounts (P. R. Boyle, ed.), p. 335–364. Academic Press, London.
1998. The Octopodinae from the eastern Atlantic Ocean and Mediterranean Sea. *In* Systematic and biogeography of cephalopods, vol. II (N. A. Voss, M. Vecchione, R. B. Toll, and M. I. Sweeney, eds.), p. 521–528. Smithsonian Contributions to Zoology 586.

Mangold, K., and S. Boletzky.
1973. New data on the reproductive biology and growth of *Octopus vulgaris*. Mar. Biol. 19:7–12.

Nixon, M.
1969. The lifespan of *Octopus vulgaris* Lamarck. Proc. malacol. Soc. Lond. 38:529–540.

Pereiro, J. A., and J. Bravo de Laguna.
1979. Dinámica de la población y evaluación de los recursos del pulpo del Atlántico Centro-oriental. Bol. Inst. Esp. Oceanogr. 5:69–105.

Raya, C. P., and C. L. Hernández-González.
1998. Growth lines within the beak microstructure of the *Octopus vulgaris* Cuvier, 1797. *In* Cephalopod biodiversity, ecology and evolution (A. I. L. Payne, M. R. Lipinski, M. R. Clarke, and M. A. C. Roeleveld, eds.), p. 135–142. S. Afr. J. Mar. Sci. 20.

Rees, W. J.
1950. The distribution of *Octopus vulgaris* Lamarck in British waters. J. Mar. Biol. Assoc. U.K. 29:361–378.

Rees, W. J., and J. R. Lumby.
1954. The abundance of Octopus in the English Channel. J. Mar. Biol. Assoc. 33:515–536.

Smale, M. J., and P. R. Buchan.
1981. Biology of *Octopus vulgaris* off the east coast of South Africa. Mar. Biol. 65(1):1–12.

Van Heukelen, W. F.
1979. Environmental control of reproduction and life span in octopus: an hypothesis. *In* Reproductive ecology of marine invertebrates (S. E. Stancyk, ed.), p. 123–133. The Belle W. Baruch Library in Marine Science 9. Univ. South Carolina Press, Columbia, SC.

Villanueva, R.
1995. Experimental rearing and growth of planktonic *Octopus vulgaris* from hatching to settlement. Can. J. Fish. Aquat. Sci. 52:2639–2650.

Young, J. Z.
1960. The statocyst of *Octopus vulgaris*. Proc. R. Soc., Ser. B 152:3–29.

# Reproduction of female spiny dogfish, *Squalus acanthias*, in the Oslofjord

**Thomas S. Jones**
Hans Oeverlandsvei 10
1363 Hoevik, Norway
E-mail address: tomtrussel@hotmail.com

**Karl I. Ugland**
Department of Biology
University of Oslo, Pb. 1064
0316 Oslo, Norway

The spiny dogfish (*Squalus acanthias*) is a relatively small shark with a characteristic spine in front of each dorsal fin. Its dorsal side is grayish and has sporadic white spots. Although it may reach a length of 160 cm, most individuals in the North Sea are in the range of 80–100 cm (Ford, 1921). It is distributed worldwide, absent only from tropical and polar regions (Compagno, 1984).

The spiny dogfish has been harvested for more than 100 years mostly for its oil-rich liver (Ketchen, 1986). At first, the oil was used for lamp fuel and as a lubricant in machines. The oil was later (during W.W.II) used as a source of vitamin A. Today the dogfish is valued as food in many countries (Gordon, 1986).

The reproduction cycle of the spiny dogfish begins with mature females bearing several large (over 40-mm) yellow eggs. As the eggs pass through the shell gland they are fertilized and become enclosed in a protective capsule (candle). The candle passes down the reproductive tract and comes to rest in the uterus. The embryos live off the large yolk sac attached under the gill region (Gilbert, 1981). As the embryos grow, they slowly absorb the yolk sack. Embryos that have completely absorbed the yolk sac may still remain in the uterus for some time before being born (Ford, 1921). According to Jones and Geen (1977), the embryos also bear an internal yolk sac which nourishes them for up to 2 months after birth. The dogfish reproductive cycle takes almost 2 years, one of the longest gestation periods of any living vertebrate (up to 24 months) (Ketchen, 1972; Nammack et al., 1985).

This shark, like other sharks, is very susceptible to overfishing, not only because of its long gestation period, but also because of slow growth, late maturity, and because it bears a small number of offspring (up to 15) (Nammack et al., 1985; Fahy, 1989). Extensive fishing since the early 1960s has led to a marked decrease in the North Sea stock. Fishing has also affected the population in the Oslofjord where the annual catch declined from 704 tons in 1979 to less than 300 tons per year during the 1990s (Official Statistics of Norway, 1996). We investigated dogfish reproduction in the Oslofjord by comparing the reproduction parameters of dogfish caught in 1987 and 1997. The focus of our study was to evaluate the reproductive parameters of spiny dogfish in the Oslofjord and to look for possible changes in these parameters.

## Materials and methods

### Sampling

Dogfish were sampled monthly off the Hvaler Islands throughout 1987 and 1997 in gill nets and by longline at depths ranging from 50 to 460 m (Fig. 1). The gill nets were composed of monofilament line (0.60 mm) with a mesh size of 285 mm. This large mesh size accounts for catches consisting mainly of larger dogfish (over 70 cm). The longline was composed of a 5-mm line connected to a 7/0 dogfish hook. The nets were usually checked every 24 hours while the longline was taken up after a few hours. The samples consisted of 132 females in 1987 and 101 females in 1997. Total length and weight were measured to the nearest 0.5 cm and 5 g, respectively, as described by Saunders and McFarlane (1993).

The same fisherman, fishing grounds and fishing gear were used in both sampling years, so that sampling bias was avoided. The fishing gear accounted for catches of dogfish that were mainly over 70 cm in length. There was a relatively small amount of unmarketable-size fish caught and the discard rates were approximately the same in both years.

### Age determination

The first and second dorsal spines were removed from 217 dogfish. The remaining 16 dogfish had spines that were either missing, or broken to such an extent that age determination was not possible for these individuals. The spines were air-dried at least 1 week before being cooked for approximately 3 minutes each in tap water. The flesh around the base of the spine was then removed with a scalpel and tweezers. The cleaned spines were subsequently dipped in ethyl alcohol and then dried with a soft cloth. Next, the spines were viewed under an Olympus SZH 10 zoom stereo microscope and aged according to the method described by Ketchen (1975). For large individuals with worn spines, age was modified by Ketchen's (1975) correction curve for age

$$Y = 0.5097X^{2.55},$$

where $X$ = the diameter of the spine base in millimeters; and
$Y$ = the additional age of the spine due to it being worn.

Ketchen's correction curve is based on fish caught in the Strait of Georgia, British Colombia. It is quite likely that there are differences in juvenile growth in the respective dogfish populations. The result may therefore be bi-

Manuscript accepted 15 March 2001.
Fish. Bull. 99:685–690 (2001)




**Figure 1**

Map of the Oslofjord. The gray shaded area represents the capture location of dogfish in the outer Oslofjord.

ased and in future work a correction curve will have to be made for the stock in the Oslofjord.

### Embryo development

Both ovaries of mature females were rinsed to expose eggs. The rinsing process consisted of removing and then carefully opening the ovaries with a scalpel. The eggs were counted and registered as belonging to the right or left ovary. In pregnant females the embryos were counted and sexed. The embryos were then measured to the nearest 0.1 cm on millimeter paper. Three stages of embryo development were identified according to Gauld[1]: stage 1 (candled embryos)—eggs, apparently fertilized, are present in a protective capsule in the uteri; stage 2 (free-living embryos)—the candle has ruptured and embryos bearing an external yolk sac are free in the uteri; and stage 3 (full term embryos)—fully developed embryos are present in the uteri. The yolk sac is fully absorbed and the umbilical slit is more or less closed.

## Results

### Biological parameters

The fish ranged in length from 54 to 110 cm in 1987 (mean 87 cm) and from 68 to 108 cm (mean 88 cm) in 1997. Weights ranged from 0.4 to 5.7 kg (mean 2.6 kg) in 1987 and from 1.1 to 5.5 kg (mean 2.9 kg) in 1997. Ages ranged from 9 to 35 years (mean 25 years) in 1987 and from 10 to 38 years (mean 23 years) in 1997.

On average, the fish in 1987 were 1.8 cm longer than fish in 1997 at the same age ($P$=0.02). However, there was a large variation in the length, and age could only account for 38% of this variability. Length accounted for 76% of the variability of the weight in 1987 and 93% in 1997. Fish between 80 and 110 cm were about 500 g heavier in 1997 ($P$<0.01). This difference was reflected in the condition factor, which was 3.8 in 1987 compared with 4.2 in 1997. This indicates that female dogfish caught in 1987 were longer and lighter than in 1997.

### Growth

Growth appeared to be best represented by a linear growth equation for dogfish between the ages of 11 and 38 years, with an average annual increment of 0.7 cm per year (Fig. 2). There was no sign of reduction in growth with increasing age; that is, no asymptote was detected in these data.

### Age and length at maturity

Maturity was defined as females bearing large ovarian eggs with a diameter of over 2 cm, candled young or free-living embryos. In both 1987 and 1997, most of the females matured between 12 and 26 years of age. PROBIT analysis indicated that dogfish reached 50% maturity at an age of 17.6 years in 1987 and 17.0 years in 1997 (Fig. 3). The

[1] Gauld, J. 1979. Reproduction and fecundity of the Scottish-Norwegian stock of spurdogs, *Squalus acanthias* (L.). ICES, Council Meeting (CM) 1979/H:54, 13 p. Directory of fisheries, biblioteket, Pb. 185, 5001 Bergen.




**Figure 2**

Growth of spiny dogfish in 1987 and 1997. ▲ = 1997 sample ($n$=79); ● = 1987 sample ($n$=92).




**Figure 3**

Percentage of mature fish in the different age groups 10–12, 13–15, 16–18,19–21, 22–24, 25–27 and 28–30 years in 1987 and 1997.

1997 sample also had a larger proportion of mature fish less than 15 years of age.

No individual was mature under 76 cm, but all individuals over 88 cm were mature, indicating that maturity occurs between 76 and 88 cm. The length at 50% maturity was 81 cm in both 1987 and 1997. There may be a difference in the growth rate between the two years; fish from 1997 reached maturity earlier than fish from 1987. However, because of the small sample size and difficulty with age determination, caution should be used to interpret this result.

## Fecundity

In 1987 the number of ovarian eggs with a diameter over 2 cm (indicating mature fish) varied between 5 and 14 (mean=8.2, SD=2.2). In 1997 the range was 3–17 eggs (mean=8.9, SD=3). The relationship between egg number and adult length in the 1987 and 1997 samples was not significantly different. On average, the egg production increased by one egg per 4-cm adult length.

The number of free-living embryos in 1987 varied from 2 to 14 per female (mean=6.6, SD=2.7). In 1997, free-living embryos numbered between 3 and 15 (mean=7.5, SD=2.6). The increment of free-living embryos per adult length (Fig. 4) in the 1987 and 1997 sample was not significantly different. However, the level of the regression line was significantly higher in 1987 ($P$=0.04). On average, the females in 1987 carried 1.2 more free-living embryos. It should be emphasized that owing to the large variability, length could only explain 45% to 56% of the difference in the number of free-living embryos.

## Reproduction cycle

Eggs on which the blastoderm was visible, were recognized as recently fertilized. This was part of the candle

stage (from the newly fertilized eggs up to embryos 10 cm long) that lasted from October to November the following year. Fertilization occurred from the beginning of October to the beginning of February. Free-living embryos were observed from October to September of the following year, followed by parturation of fully mature embryos from September to December. The duration of the pregnancy for spiny dogfish in the Oslofjord ranged from 18 to 24 months. This period of time is calculated from the time of fertilization to the time of parturation. These estimates of the reproduction phases are summarized in Figure 5.

### Embryonic growth

The embryos ranged from 4.4 to 24.9 cm total length. The growth of the combined 1986 and 1996 year classes of dogfish embryos was followed throughout their second year of development. Growth was approximately 1 cm per month for both year classes in the size range from 8 to 24 cm, which corresponds approximately to the embryos in the second year of development. However, growth was not uniform; it could be divided into two growth phases (Fig. 6). The first phase lasted from October to May with a slow growth of 0.6 cm/month. The second phase lasted from May until December and was characterized by a more rapid growth of 1.2 cm/month.




**Figure 4**

Number of free-living embryos in relationship to the length of the adult fish in 1987 and 1997. ▲ = 1987 sample; ● = 1997 sample.




**Figure 5**

Schematic time representation of the reproductive cycle of the female spiny dogfish by month ($n$=194).

## Discussion

### Growth

The lack of an asymptote is probably due to an absence of older, larger fish. This is most likely due to over-harvesting of dogfish in the Oslofjord. According to local fishermen, heavy catches of large individuals (exceeding 110 centimeters) were taken in the late 70s and early 80s.[2]

### Age and length at maturity

Age at 50% maturity was found to be 17.6 years in 1987 and 17.0 years in 1997. For spiny dogfish captured between the East Coast of Scotland and Norway, Aasen (1961) estimated age at 50% maturity to be in the range 12 to 14 years. In an area northwest of Scotland, Holden and Meadows (1962) reported age at 50% maturity of 9 years. However, Aasen (1961) and Holden and Meadows (1962) did not take into account worn spines, which combined with the difficulty of identifying all of the annual rings, means that their estimates are most likely biased downwards.

Length at 50% maturity in the Oslofjord should not deviate greatly from sizes observed elsewhere in the northeast Atlantic. Length at 50% maturity in the Oslofjord was found to be 81 cm in both 1987 and 1997. This is in general agreement with other observations in the northeast Atlantic where length at 50% maturity varied from 74 to 83 cm (Hickling, 1930; Holden and Meadows, 1964; Fahy, 1989; Gauld[1]).

### Fecundity

There was no significant difference between the number of eggs observed in 1987 and 1997. However, the number of eggs in the ovaries was higher than the number of embryos in the uterus (average: 8.1 eggs and 6.6 free-living embryos in 1987 and 8.9 eggs and 7.5 free-living embryos in 1997). This is probably due to the absorption of eggs in the ovaries following fertilization that was observed by Hanchet (1988), Holden and Meadows (1964), and Nammack et al. (1985).

[2] Johansen, P. A. 1997. Personal commun. Dogfish fisherman from the Oslofjord. Gudeberg Allè 1, 1600 Fredrikstad, Norway.




**Figure 6**

Length of the embryos during their second year of development for the combined 1987 and 1997 sample.

Fecundity was within the bounds found elsewhere in the northeast Atlantic (Aasen, 1961; Gauld[1]). Free-living embryos showed the same increase in number per length in 1987 and 1997, but the level was significantly lower in 1997 (on average one less embryo per unit of length). In addition the fish were shorter and heavier in 1997. However, these results must be taken with some caution because of the large variability in the data.

A small sample size and annual variability in both embryo production and food availability may have contributed to the lower fecundity in 1997. With regard to the sample sizes (31 and 38 fishes), it is emphasized that the covariation between the number of free-living embryos versus length was rather low. Although the differences between 1987 and 1997 were statistically significant, the overlap was so large that caution must be used in interpretation of the data.

## Reproduction cycle and embryonic growth

In the Oslofjord, fertilization occurs from October to February and parturation from October to December. Research in other areas has shown that the pregnancy lasts 22–23 months (Ford, 1921; Gauld[1]). Thus, the duration of pregnancy in the Oslofjord (18 to 24 months) seems to have a greater variation than these observations. This variation may have been caused by the sample from the Oslofjord, which included dogfish that were fertilized early in the fertilization period, as well as dogfish that were fertilized towards the end of the fertilization period. This, in turn, indicates the possible minimal and maximal duration of pregnancy.

Behavioral factors may influence pregnancy duration, as well. Some dogfish may inhabit cooler water for longer periods of time than other dogfish, which may increase the duration of their pregnancy.

The differences in the embryo growth rate during their second year of development may be related to the water temperature. Hickling (1930) found that females migrate from deep to shallow water as pregnancy proceeds. This migration pattern exposes the embryos to different temperature levels that may influence embryonic growth, as found in Newfoundland by Templeman (1944). Embryos outside of Newfoundland had an average growth rate of 1.1 cm/month and a 24-month pregnancy period (Templeman, 1944). Outside of Woods Hole, Hisaw and Albert (1947) found an average growth rate of 1.3 cm/month and a pregnancy period of 20–22 months. Nammack et al. (1985) stated that the lower sea temperature outside of Newfoundland could be a possible reason for this difference.

In phase 1, growth is slow and corresponds to a period of low water temperature (winter and spring). As females begin to migrate towards shallower water, they encounter warmer water (summer and fall). The increased growth rate in phase 2 is therefore most likely related to an increase in water temperature.

## Acknowledgments

We would like to thank P. A. Johansen for making the collection of the sharks possible, as well as giving us valuable suggestions. We thank scientists and personnel of the biological station in Drøbak and H. E. Karlsen for the use of equipment associated with age determination. We are also particularly grateful to T. R. Wiley and H. Zidowitz for their comments on the manuscript.

## Literature cited

Aasen, O.
1961. Pigghå-undersøkelsene. Fisken og havet 1:1–9.
Compagno, L. J. V.
1984. FAO species catalogue. Vol. 4: Sharks of the world: an annoted and illustrated catalogue of shark species known to date. Part 1. Hexiformes to Lamniformes. FAO Fish. Synop. 125:1–249.
Fahy, E.
1989. The spurdog *(Squalus acanthias)* fishery in South West Ireland. Ir. Fish. Invest. ser. B. no. 32, 22 p.
Ford, E.
1921. A contribution to our knowledge of the life-histories of the dogfishes landed at Plymouth. J. Mar. Biol. Assoc. U.K. 12:468–505.
Gilbert, P. W.
1981. Patterns of shark reproduction. Oceanus 24:30–39.
Gordon, D. G.
1986. The spiny dogfish. Oceans 19:56–59.
Hanchet, S.
1988. Reproductive biology of *Squalus acanthias* from the east coast, South Island, New Zealand. N.Z.. J. Mar. Freshwater Res. 22:537–549.
Hickling, C. F.
1930. A contribution towards the life history of the spurdog. J. Mar. Biol. Assoc. U.K. 16:529–576.
Hisaw, F. L., and A. Albert.
1947. Observations on the reproduction of spiny dogfish, *Squalus acanthias.* Biol. Bull. 92:187–199.
Holden, M. J., and P. S. Meadows.
1962. The structure of the spine of the spur dogfish (*Squalus acanthias*) and its use for age determination. J. Mar. Biol. Assoc. U.K. 42:179–197.

1964. The fecundity of the spurdog (*Squalus acanthias*). J. Cons. Perm. Int. Explor. Mer 28:418–424.

Jones, B. C., and G. H. Geen.
1977. Reproduction and embryonic development of spiny dogfish (*Squalus acanthias*) in the Strait of Georgia, British Colombia. J. Fish. Res. Board Can. 34:1286–1292.

Ketchen, K. S.
1972. Size at maturity, fecundity, and embryonic growth of the spiny dogfish (*Squalu sacanthias*) in British Colombia Waters. J. Fish. Res. Board Can. 29:1717–1723.
1975. Age and growth of dogfish, *Squalus acanthias*, in British Colombia Waters. J. Fish. Res. Board Can. 32: 43–59.
1986. The spiny dogfish (*Squalus acanthias*) in the Northeast Pacific and a history of its utilization. Can. Spec. Publ. Fish. Aquat. Sci. 88, 78 p.

Nammack, M. F., J. A. Musick, and J. A. Colvocoresses.
1985. Life history of spiny dogfish off the Northeastern United States. Trans. Am. Fish. Soc. 114:367–376.

Official Statistics of Norway.
1996. Fishery statistics 1961–1996. Graphic Communication Systems, Inc. (GCS) A/S, Oslo, Norway, 116 p.

Saunders, M. W., and G. A. McFarlane.
1993. Age and length at maturity of the female spiny dogfish, *Squalus acanthias*, in the Strait of Georgia, British Colombia, Canada. Environ. Biol. Fish. 38:49–57.

Templeman, W.
1944. The life-history of the spiny dogfish (*Squalus acanthias*) and the vitamin A values of dogfish liver oil. Nfld. Gov., Dep. Nat. Resour., Fish. Res. Bull. 15:1–100.

# Estimating live standard length of net-caught walleye pollock (*Theragra chalcogramma*) larvae using measurements in addition to standard length*

**Steven M. Porter**

**Annette L. Brown**

**Kevin M. Bailey**

Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
7600 Sand Point Way NE
Seattle, Washington 98115

E-mail address (for S. M. Porter): steve.porter@noaa.gov

Accurate measurement of larval fish lengths from ichthyoplankton samples is critical in the estimation of growth, birthdate and mortality. It is well known that larvae shrink in length 1) when caught in plankton nets (Theilacker, 1980; Hay, 1981), 2) between the time of collection and preservation, and 3) afterwards from the effects of preservation in chemicals or from freezing (Theilacker, 1980; Fowler and Smith, 1983; Yin and Blaxter, 1986). Shrinkage caused by the plankton net (the net can damage a larva's integument; Holliday and Blaxter, 1960) and by delays in preservation can be up to 40% of the initial live length (Hay, 1981). In some cases, measurements of larval body length may be unsuitable for growth estimation (Jennings, 1991). In other cases, algorithms have been developed to estimate live lengths from preserved larvae (Bailey 1982, Hjörleifsson and Klein-MacPhee, 1992; Fey, 1999) and from net-caught and preserved larvae (Theilacker, 1980; Theilacker and Porter, 1995; Fox, 1996). It has also been noted that constant shrinkage-correction factors should be applied cautiously (Pepin et al., 1998).

Shrinkage of the standard length of fish larvae is species-specific (Jennings, 1991; Fey, 1999), size-dependent (Fowler and Smith, 1983; Theilacker and Porter, 1995; Fey, 1999), solution-dependent (Hay, 1982; Tucker and Chester, 1984), and may also depend to some degree on the ambient temperature at the time of preservation (Hay, 1982). Other body measurements may shrink due to effects of collection and preservation. In fact, many corrections for larval shrinkage have been developed with respect to estimating morphometric-dependent condition factors (Theilacker, 1980; McGurk, 1985; Yin and Blaxter, 1986).

Measurement of otoliths has been proposed as a method to correct for shrinkage in the length of fish larvae (Leak, 1986; Radtke, 1989), but this method may not be generally applicable because of systematic variations in the relationship between fish size and otolith size (Neilson and Campana, 1990). In addition, this relationship can be nonlinear and unpredictable. Another approach to obtain shrinkage correction factors is by controlled experiments in which larval shrinkage is recorded from the time of death, or for the period of time larvae are in a plankton net, and again after preservation. The problem with this approach is that during most sampling at sea, the time larvae enter the net to the time of their preservation is unknown and can vary from several minutes to a half hour or more, depending on their depth of capture and the duration of the tow.

In our study, we used measurements of other body parts as ancillary data to estimate live lengths of preserved fish larvae caught at sea. We report on two common preservatives: ethanol and formalin. To avoid the problem of duration of time in the net, shrinkage correction equations were formulated by pooling data of known-length larvae that were treated for varying durations in simulated plankton tows in the laboratory and then preserved. Morphometric measurements were collected on preserved larvae that could be used in equations to obtain more accurate and precise estimates of live length. Rather than use alternative body part measurements as a substitute for length, we hypothesized that other body parts, nonshrinking or otherwise, could be used to correct for shrinkage, in spite of how long larvae had been in nets.

## Materials and methods

### Rearing protocol

Spawning walleye pollock (*Theragra chalcogramma*) were collected by trawl in the eastern Bering Sea by the NOAA ship *Miller Freeman* during April 1999. Fertilized eggs were transported to the Alaska Fisheries Science Center, Seattle, Washington, where they were incubated at 6°C in the dark in 4-L glass jars filled with 3 L of filtered seawater at a salinity of 33‰. Larvae were reared in 120-L circular, black fiberglass tanks (62 cm diameter, 43 cm deep) filled with 90-L filtered seawater, and maintained at a temperature of 6°C. Two replicate tanks with approximately 1000 eggs each were maintained. A 16:8 hour light:dark cycle was started at hatching and the amount of light at the surface of the water varied from 3.0 to 3.5 μ mol photon/$m^2$/s. For the first two weeks of feeding, prey consisted of rotifers, *Brachionus plicatilis*, at a concentration of approximately 10/mL, and wild zooplankton (a mixture of *Acartia* sp. copepod nauplii and copepodites, and gastropod and polychaete larvae; at approximately

* Contribution FOCI-0400 of the Fisheries Oceanography Coordinated Investigations (FOCI), NOAA, 7600 Sand Point Way NE, Seattle, WA 98115.

Manuscript accepted 19 March 2001.
Fish. Bull. 99:691–696 (2001).

1/mL) collected from local lagoons. After two weeks of feeding, *Artemia* sp. brine shrimp (at approximately 10/mL) were added as well. Further details on larval rearing are described in Porter and Theilacker.[1]

### Shrinkage experiments

Larval walleye pollock shrinkage was examined for three age groups of larvae (8, 15, and 33 days after hatching) with treatment with preservative only (no net treatment), with 5-minute net treatment and then treatment with preservative, and 15-minute net treatment and then treatment with preservative. The preservatives used were 5% buffered formalin in 33‰ seawater, and 95% ethanol buffered with 0.6 mM sodium carbonate, and 0.6 mM sodium bicarbonate. For each age group, 10 larvae were sampled for each treatment and preservative combination. A pipette was used to place a live larva in seawater on a microscope slide and the larva was quickly recorded on videotape with a Panasonic 5100 digital video camera mounted on a dissecting microscope. Next, the larva was either placed directly into preservative or into the net-treatment apparatus, after which it was preserved. The net-treatment apparatus consisted of a small net submerged in a tank of 6°C seawater. A submersible pump was used to circulate seawater through the net to simulate a plankton tow (Theilacker, 1980). Water flow through the net was adjusted so that a larva would come in contact with the mesh as it might during a tow at sea. One month was chosen as the length of time walleye pollock larvae would remain in preservative before they were recorded on videotape. The length of time for standard length to stabilize in preservative varies depending on the species of fish. For herring (*Clupea harengus*) larvae, standard length was stable after 30 days (Fox, 1996); for red sea bream (*Chrysophrys major*) larvae, most of the shrinkage occurred during the first 2–3 days in presevative (Rosenthal et al., 1978). Optimus version 5.0 image analysis software was used to measure standard length (SL), eye diameter, head length (tip of snout to pectoral fin), and body depth at the anus of both live and preserved larvae.

Otoliths were dissected from the preserved larvae, and maximum diameter of the sagittae were measured at 1000× magnification with an ocular micrometer. Some of the otoliths from larvae preserved in formalin had eroded edges and their diameters could not be measured reliably; therefore, otoliths from formalin-preserved larvae were not used in the shrinkage correction model.

Myomere width measurements taken from formalin preserved larvae were used in the shrinkage correction model. The width of five consecutive myomeres located between the two pigment bands on the tail were measured at 50× with an ocular micrometer and polarized light. For ethanol-preserved larvae, myomere measurements could not be made because the tissue had turned opaque during preservation.

### Data analysis

For each preservative, data from all larval ages and shrinkage treatments were pooled and backward stepwise regression (SPSS, Inc., 1998) was used to determine which variables produced the best model for estimating live SL. The data were pooled because the regression is used to estimate live SL of larvae caught at sea and because the amount of time larvae spend in the sampling net is unknown; therefore a general relationship was sought.

For each preservative, the growth rate of walleye pollock larvae between the first (8 days after hatching) and last sampling days (33 days after hatching) was determined by using linear regression. Only the first and last sampling days were used because there is a lag in growth of walleye pollock larvae that occurs for about a week during the transition from endogenous to exogenous feeding (Yamashita and Bailey, 1989). Dunnett's test (Zar, 1996) was used to compare growth rates calculated with live SL (which was considered to be the control) against growth rates calculated from the shrinkage correction models.

## Results

### Shrinkage with ethanol

Walleye pollock larvae shrank an average of 6% in 95% ethanol when all shrinkage treatments were pooled (Fig. 1A). Larvae used in the shrinkage experiments ranged in live SL from 5.41 to 9.25 mm. The best model to correct for shrinkage contained preserved SL, body depth, and otolith diameter, $r^2 = 0.90$ (Table 1, Fig. 1B). Results indicated that there was borderline multicollinearity in the data (the variance inflation factor for otolith diameter was 4.79), probably due to correlation between preserved SL and otolith diameter. The $r^2$ with only preserved SL was 0.82. The use of additional variables reduced the mean square error (MSE) by 44% (Table 1).

Residuals from the best model increased as the percentage of shrinkage increased, which suggests that the shrinkage correction model does not work well for larvae that shrink greatly. At largest values of shrinkage, 17% and 19%, the best model underestimated SL by 10%, but at 15% shrinkage the error was 3%.

### Shrinkage with formalin

For larvae preserved in 5% formalin, the overall mean shrinkage was 10%. Larvae ranged in live SL from 5.61 to 9.43 mm (Fig. 2A). When compared with the model using only preserved SL ($r^2$=0.81, MSE=0.112; Table 1), the model containing both preserved SL and body depth at the anus produced the best model for estimating live SL ($r^2$=0.88, MSE=0.073; Table 1, Fig. 2B).

[1] Porter, S. M., and G. H. Theilacker. 1996. Larval walleye pollock, *Theragra chalcogramma*, rearing techniques used at the Alaska Fisheries Science Center, Seattle, Washington. AFSC Processed Report 96-06, 26 p. U.S. Dep. Commerce, NOAA, NMFS, Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle, WA, 98115.

**Table 1**

Shrinkage correction equations used to estimate the live standard length (SL) of net-treated and preserved laboratory-reared walleye pollock (*Theragra chalcogramma*) larvae. Best model and model containing only preserved standard length (PSL) are shown. For 95% ethanol, results indicated that there was borderline multicollinearity in the data (see "Results" section). All variables were measured in millimeters. Body depth was measured at the anus. Maximum diameter of the sagitta otolith was used.

| 95% ethanol ($n$=91; SL range: 5.41 to 9.25 mm[1]) | $r^2$ | MSE |
|---|---|---|
| *Live SL* = 2.18 + 0.464(*PSL*) + 1.71(*body depth*) + 30.53(*otolith diameter*) | 0.90 | 0.073 |
| *Live SL* = 0.230 + 1.02(*PSL*) | 0.82 | 0.130 |
| 5% formalin ($n$=90; SL range: 5.61 to 9.43[1]) | | |
| *Live SL* = 2.52 + 0.509(*PSL*) + 3.18(*body depth*) | 0.88 | 0.073 |
| *Live SL* = 1.88 + 0.807(*PSL*) | 0.81 | 0.112 |

[1] Live standard length of larvae used to formulate regression equations.

**Table 2**

Growth rates of laboratory-reared walleye pollock (*Theragra chalcogramma*) larvae between the first and last sampling days determined by using live standard length (LSL), and standard length corrected for shrinkage (for shrinkage correction equations see Table 1).

| 95% ethanol ($n$=61) | Growth rate (mm/day) | % Error from LSL |
|---|---|---|
| Live standard length | 0.056 | |
| Standard length corrected with only preserved standard length | 0.048 | 14 |
| Standard length corrected with best model | 0.058 | 4 |
| 5% formalin ($n$=60) | | |
| Live standard length | 0.050 | |
| Standard length corrected with only preserved standard length | 0.043 | 14 |
| Standard length corrected with best model | 0.045 | 10 |

For 5% formalin, residuals for the best shrinkage correction model did not increase as the amount of shrinkage increased; therefore the model worked well for all shrinkage up to the maximum amount of 27%.

### Growth rates

Growth rates were calculated with live SL, corrected SL from the preserved SL only models, and corrected SL from the multivariate models. For each preservative there was no significant difference between the growth rate for live SL and growth rates calculated with the models (Dunnett's test, $P$>0.05). Although the growth rates were not significantly different, growth rates calculated from the multivariate models were more similar to growth rates calculated from live SL (Table 2); therefore additional morphometric variables improve growth rate estimates over using preserved SL alone.

## Discussion

Fish larvae shrink considerably when they are caught in nets and preserved. Shrinkage during collection is most likely caused by the plankton net, which can damage the larva's integument (Holliday and Blaxter, 1960). Shrinkage also varies between preservatives because of the differing ionic strengths of these solutions (Parker, 1963; Hay, 1982; Tucker and Chester, 1984). We found that shrinkage of walleye pollock larvae was greater in 5% formalin than in 95% ethanol—a finding similar to results from other species (Bailey, 1982; Fey, 1999). Often only preserved SL has been used to correct for shrinkage in length (Theilacker and Porter, 1995; Fox, 1996; Kristoffersen and Gro Vea Salvanes, 1998), although other measurements, such as otolith size, have been suggested for use to correct for larval fish shrinkage (Leak, 1986; Radtke, 1989,1990). By using measurements that are easily made and readily adapted to




**Figure 1**

Uncorrected (**A**) and corrected (**B**) standard length determined by using best shrinkage correction model, see Table 1) for laboratory-reared walleye pollock (*Theragra chalcogramma*) larvae subjected to various shrinkage treatments and preserved in 95% ethanol ($n$=91). The 95% confidence limits are shown for each predicted value, and the diagonal line shows a 1:1 ratio.

routine data collection, we have shown improved accuracy of shrinkage correction models by use of measurements in addition to preserved SL: for 5% formalin, 7% more of the variance was explained by adding body depth; and for 95% ethanol, 8% more of the variance was explained by including body depth and otolith diameter. For 95% ethanol, the shrinkage correction model formulated in our study may significantly underestimate SL when shrinkage is high, but because there were only two larvae in this category, results were inconclusive.

Shrinkage correction models are usually formulated by using fed, laboratory-reared larvae (Jennings, 1991; Theilacker and Porter, 1995; Fox, 1996), but individual larvae response to handling and preservation can vary significantly (Pepin et al., 1998). These correction models were developed to allow more accurate calculation of the growth rate of fish larvae in the sea and to apply laboratory-derived indices to the field. We have shown factors in addition to preserved SL that may improve the accuracy of live-length estimates for walleye pollock.




**Figure 2**

Uncorrected (**A**) and corrected (**B**) standard length determined by using best shrinkage correction model, see Table 1) for laboratory-reared walleye pollock (*Theragra chalcogramma*) larvae subjected to various shrinkage treatments and preserved in 5% formalin ($n$=90). The 95% confidence limits are shown for each predicted value, and the diagonal line shows a 1:1 ratio.

## Acknowledgments

We thank Debbie Blood for care of the pollock eggs aboard ship, and for bringing them back to Seattle. Kathy Mier, Susan Picquelle, and three anonymous reviewers commented on drafts of the manuscript and offered improvements.

## Literature cited

Bailey, K. M.
1982. The early life history of Pacific hake, *Merluccius productus*. Fish. Bull. 80:589–598.

Fey, D. P.
1999. Effects of preservation technique on the length of larval fish: methods of correcting estimates and their implication for studying growth rates. Arch. Fish. Mar. Res. 47: 17–29.

Fowler, G. M., and S. J. Smith.
1983. Length changes in silver hake (*Merluccius bilinearis*) larvae: effects of formalin, ethanol, and freezing. Can. J. Fish. Aquat. Sci. 40:866–870.

Fox, C. J.
1996. Length changes in herring (*Clupea harengus*) larvae: effects of capture and storage in formaldehyde and alcohol. J. Plankton Res. 18:483–493.

Hay, D. E.
1981. Effects of capture and fixation on gut contents and body size of Pacific herring larvae. Rapp. P-V. Reun. Cons. Int. Explor. Mer 178:395–400.
1982. Fixation shrinkage of herring larvae: effects of salinity, formalin concentration, and other factors. Can. J. Fish. Aquat. Sci. 39:1138–1143.

Hjörleifsson E., and G. Klein-MacPhee.
1992. Estimation of live standard length of winter flounder *Pleuronectes americanus* larvae from formalin-preserved, ethanol-preserved and frozen specimens. Mar. Ecol. Prog. Ser. 82:13–19.

Holliday, F. G. T., and J. H. S. Blaxter.
1960. The effects of salinity on the developing eggs and larvae of the herring. J. Mar. Biol. Assoc. U.K. 39:591–603.

Jennings, S.
1991. The effects of capture, net retention and preservation upon lengths of larval and juvenile bass, *Dicentrarchus labrax* (L.). J. Fish Biol. 38:349–357.

Kristoffersen, J. B., and A. Gro Vea Salvanes.
1998. Effects of formaldehyde and ethanol preservation on body and otoliths of *Maurolicus muelleri* and *Benthosema glaciale*. Sarsia 83:95–102.

Leak, J. C.
1986. The relationship of standard length and otolith diameter in larval bay anchovy, *Anchoa mitchilli* (Val.). A shrinkage estimator. J. Exp. Mar. Biol. Ecol. 95:167–172.

McGurk, M. D.
1985. Effects of net capture on the postpreservation morphometry, dry weight, and condition factor of Pacific herring larvae. Trans. Am. Fish. Soc. 114:348–355.

Neilson, J. D., and S. E. Campana.
1990. Comment on "Larval fish age, growth, and body shrinkage: information available from otoliths." Can. J. Fish. Aquat. Sci. 47: 2461–2463.

Parker, R. R.
1963. Effects of formalin on the length and weight of fishes. J. Fish. Res. Board Can. 20:1441–1455.

Pepin, P., J. F. Dower, and W. C. Leggett
1998. Changes in the probability density function of larval fish body length following preservation. Fish. Bull. 96:633–640.

Radtke, R. L.
1989. Larval fish age, growth, and body shrinkage: information available from otoliths. Can. J. Fish. Aquat. Sci. 46:1884–1894.
1990. Information storage capacity of otoliths: response to Nielson and Campana. Can. J. Fish. Aquat. Sci. 47:2463–2467.

Rosenthal, H., D. Kuhlmann, and O. Fukuhara
1978. Shrinkage of newly hatched larvae of the red sea bream (*Chrysophrys major* Temminck and Schlegel) preserved in formalin. Arch. Fischwiss. 29:59–63.

SPSS, Inc.
1998. SYSTAT 8.0: statistics. SPSS, Inc., Chicago, IL, 1086 p.

Theilacker, G. H.
1980. Changes in body measurements of larval northern anchovy, *Engraulis mordax*, and other fishes due to handling and preservation Fish. Bull. 78:685–692.

Theilacker, G. H., and S. M. Porter.
1995. Condition of larval walleye pollock, *Theragra chalcogramma*, in the western Gulf of Alaska assessed with histological and shrinkage indices. Fish. Bull. 93:333–344.

Tucker, J. W., Jr., and A. J. Chester.
1984. Effects of salinity, formalin concentration and buffer on quality of preservation of southern flounder (*Paralichthys lethostigma*) larvae. Copeia 1984:981–988.

Yamashita, Y., and K. M. Bailey.
1989. A laboratory study of the bioenergetics of larval walleye pollock, *Theragra chalcogramma*. Fish. Bull. 87:525–536.

Yin, M. C., and J. H. S. Blaxter.
1986. Morphological changes during growth and starvation of larval cod (*Gadus morhua* L.) and flounder (*Platichthys flesus* L.). J. Exp. Mar. Biol. Ecol. 104:215–228.

Zar, J. H.
1996. Biostatistical analysis, third ed. Prentice Hall, Upper Saddle River, NJ, 622 p.

# Preliminary study of albacore (*Thunnus alalunga*) stock differentiation inferred from microsatellite DNA analysis

**Motohiro Takagi**
**Tetsuro Okamura**
Department of Cultural Fisheries
Faculty of Agriculture
Kochi University, Monobe
Nankoku, Kochi 783-8502, Japan

**Seinen Chow**
National Research Institute of Far Seas Fisheries
5-7-1 Orido
Shimizu 424-8633, Japan

**Nobuhiko Taniguchi**
Department of Cultural Fisheries
Faculty of Agriculture
Kochi University, Monobe
Nankoku, Kochi 783-8502, Japan
Present address (for N. Taniguchi, contact author): Graduate School of Agriculture
Tohoku University
1-1 Tsutsumidori, Amamiyamachi
1-2 Sendai 981-8555, Japan
E-mail address (for N. Taniguchi, contact author): nobuhiko@bios.tohoku.ac.jp

The albacore (*Thunnus alalunga*) is a highly migratory large pelagic tuna, common from tropical to temperate areas of all oceans, including the Mediterranean Sea. Although albacore populations of each ocean or each hemisphere have been managed separately, relationships between albacore populations of northern and southern hemispheres within an ocean are controversial. Differences in morphometry, movement, and catch statistics of albacore between northern and southern hemispheres within Atlantic and Pacific Oceans have been reported (Kurogane and Hiyama, 1958; Ishii, 1965; Nakamura, 1969). Examining mtDNA variation, Chow and Ushiama (1995) detected very little genetic difference between samples from northern and southern hemispheres. They proposed that minor gene flow may have occurred between albacore populations of northern and southern hemispheres—enough to prevent genetic differentiation. However, it is also possible that insufficient time has elapsed since population subdivision for mtDNA genotypic rearrangement. Because all tuna species of the genus *Thunnus* are thought to be phylogenetically new (Chow and Kishino, 1995), use of highly variable genetic markers may be necessary to investigate genetic differentiation between stocks.

Recently, Takagi et al. (1999) isolated four microsatellite loci from Pacific northern bluefin tuna (*Thunnus thynnus orientalis*) and demonstrated successful cross-species amplification of homologous microsatellites in other tuna species. In our study, we used these microsatellite primers to evaluate genetic variation within and between albacore samples from the Atlantic and the Pacific Oceans and present genetic evidence of population structuring between and within ocean samples of albacore.

## Materials and methods

Five albacore samples were drawn from archival stock materials in the National Research Institute of Far Seas Fisheries laboratory. One sample each came from the Northwest Pacific (NW Pacific; Japan), Southwest Pacific (SW Pacific; Australia), Southeast Pacific (SE Pacific; Chile and Peru), and two came from the Northeast Atlantic (NE Atlantic; Biscay Bay) and the Southwest Atlantic (SW Atlantic; Brazil). These samples were derived from the same sample lots used by Chow and Ushiama (1995). Nucleotide sequences of the four primer sets, PCR amplification conditions needed to amplify the four microsatellite loci (*Ttho-1**, *-4**, *-6** and *-7**) developed for Pacific northern bluefin tuna (*T. thynnus orientalis*), and electrophoresis procedures can be found in Takagi et al. (1999). Differentiation of allele frequencies between and among samples was estimated by a fixation index ($F_{ST}$) with Arlequin version 1.1 (Schneider et al., 1997).

## Results

Alleles observed in each locus were as follows: 9 in *Ttho-1**, 29 in *Ttho-4**, 31 in *Ttho-6**, and 18 in *Ttho-7** (Table 1). All four sets of PCR primers successfully amplified scorable microsatellite loci for all samples. Observed heterozygosity ($H_0$) ranged from 0.391 to 0.914 at *Ttho-1**, from 0.688 to 0.886 at *Ttho-4**, from 0.548 to 0.884 at *Ttho-6**, and from 0.857 to 1 at *Ttho-7**. We found no substantial discrepancy between observed and expected number of genotypes for any locus.

All samples except NE Atlantic shared the most common alleles for all loci, whereas the NE Atlantic sample shared the most common alleles only within *Ttho-1**. The NE Atlantic sample also did not share the second most common allele with the other samples for all loci. $F_{ST}$ among all five

Manuscript accepted 11 April 2001.
Fish. Bull. 99:697–701 (2001).

**Table 1**

Allele frequency and genetic variability for four microsatelite loci surveyed for albacore used in this study.

| Locus | Allele (base pairs) | NW Pacific | SW Pacific | SE Pacific | SW Atlantic | NE Atlantic |
|---|---|---|---|---|---|---|
| Ttho-1* | 172 | 0.000 | 0.000 | 0.021 | 0.000 | 0.000 |
| | 174 | 0.000 | 0.016 | 0.000 | 0.000 | 0.000 |
| | 176 | 0.075 | 0.065 | 0.043 | 0.054 | 0.065 |
| | 178 | 0.000 | 0.097 | 0.096 | 0.000 | 0.032 |
| | 180 | 0.234 | 0.177 | 0.202 | 0.141 | 0.145 |
| | 182 | 0.521 | 0.452 | 0.436 | 0.685 | 0.516 |
| | 184 | 0.149 | 0.161 | 0.160 | 0.098 | 0.210 |
| | 186 | 0.000 | 0.032 | 0.043 | 0.022 | 0.032 |
| | 188 | 0.021 | 0.000 | 0.000 | 0.000 | 0.000 |
| No. of samples | | 47 | 31 | 47 | 46 | 31 |
| No. of allele | | 5 | 7 | 7 | 5 | 6 |
| Effective no. of alleles[1] | | 2.82 | 3.62 | 3.70 | 1.99 | 2.96 |
| Observed heterozygosity ($H_o$) | | 0.659 | 0.871 | 0.914 | 0.391 | 0.645 |
| Expected heterozygosity ($H_e$) | | 0.645 | 0.724 | 0.730 | 0.498 | 0.662 |
| Ttho-4* | 134 | 0.000 | 0.000 | 0.000 | 0.000 | 0.032 |
| | 136 | 0.000 | 0.016 | 0.011 | 0.000 | 0.096 |
| | 138 | 0.011 | 0.016 | 0.033 | 0.011 | 0.065 |
| | 140 | 0.043 | 0.031 | 0.067 | 0.011 | 0.452 |
| | 142 | 0.032 | 0.031 | 0.167 | 0.000 | 0.032 |
| | 144 | 0.085 | 0.094 | 0.100 | 0.102 | 0.065 |
| | 146 | 0.383 | 0.313 | 0.178 | 0.261 | 0.048 |
| | 148 | 0.032 | 0.078 | 0.089 | 0.080 | 0.097 |
| | 150 | 0.011 | 0.094 | 0.056 | 0.023 | 0.032 |
| | 152 | 0.053 | 0.047 | 0.067 | 0.102 | 0.048 |
| | 154 | 0.096 | 0.031 | 0.044 | 0.046 | 0.032 |
| | 156 | 0.011 | 0.000 | 0.011 | 0.068 | 0.000 |
| | 158 | 0.032 | 0.047 | 0.000 | 0.057 | 0.000 |
| | 160 | 0.043 | 0.047 | 0.044 | 0.046 | 0.000 |
| | 162 | 0.021 | 0.031 | 0.033 | 0.034 | 0.000 |
| | 164 | 0.011 | 0.078 | 0.044 | 0.023 | 0.000 |
| | 166 | 0.053 | 0.000 | 0.000 | 0.023 | 0.000 |
| | 168 | 0.021 | 0.031 | 0.011 | 0.011 | 0.000 |
| | 170 | 0.000 | 0.000 | 0.022 | 0.011 | 0.000 |
| | 174 | 0.000 | 0.000 | 0.011 | 0.023 | 0.000 |
| | 176 | 0.000 | 0.000 | 0.000 | 0.023 | 0.000 |
| | 178 | 0.000 | 0.000 | 0.011 | 0.023 | 0.000 |
| | 180 | 0.032 | 0.000 | 0.000 | 0.000 | 0.000 |
| | 182 | 0.011 | 0.000 | 0.000 | 0.000 | 0.000 |
| | 184 | 0.011 | 0.000 | 0.000 | 0.000 | 0.000 |
| | 186 | 0.000 | 0.016 | 0.000 | 0.000 | 0.000 |
| | 190 | 0.011 | 0.000 | 0.000 | 0.000 | 0.000 |
| | 202 | 0.000 | 0.000 | 0.000 | 0.011 | 0.000 |
| | 232 | 0.000 | 0.000 | 0.000 | 0.011 | 0.000 |
| No. of samples | | 47 | 32 | 45 | 44 | 31 |
| No. of allele | | 20 | 16 | 18 | 21 | 11 |
| Effective no. of alleles[1] | | 5.62 | 7.14 | 10.20 | 8.85 | 4.17 |
| Observed heterozygosity ($H_o$) | | 0.851 | 0.688 | 0.844 | 0.886 | 0.710 |
| Expected heterozygosity ($H_e$) | | 0.822 | 0.860 | 0.902 | 0.887 | 0.760 |
| Ttho-6* | 127 | 0.000 | 0.000 | 0.000 | 0.012 | 0.000 |
| | 129 | 0.000 | 0.000 | 0.000 | 0.012 | 0.000 |
| | 131 | 0.011 | 0.032 | 0.000 | 0.000 | 0.000 |
| | 133 | 0.011 | 0.000 | 0.000 | 0.000 | 0.000 |
| | 135 | 0.033 | 0.113 | 0.047 | 0.093 | 0.000 |

*continued*

**Table 1 (continued)**

| Locus | Allele (base pairs) | NW Pacific | SW Pacific | SE Pacific | SW Atlantic | NE Atlantic |
|---|---|---|---|---|---|---|
| Ttho-6* (continued) | 137 | 0.100 | 0.000 | 0.023 | 0.105 | 0.000 |
| | 139 | 0.056 | 0.016 | 0.093 | 0.047 | 0.000 |
| | 141 | 0.478 | 0.629 | 0.233 | 0.349 | 0.083 |
| | 143 | 0.067 | 0.048 | 0.081 | 0.105 | 0.017 |
| | 145 | 0.167 | 0.113 | 0.163 | 0.128 | 0.050 |
| | 147 | 0.067 | 0.048 | 0.093 | 0.058 | 0.300 |
| | 149 | 0.000 | 0.000 | 0.081 | 0.047 | 0.083 |
| | 151 | 0.011 | 0.000 | 0.023 | 0.012 | 0.017 |
| | 153 | 0.000 | 0.000 | 0.012 | 0.000 | 0.033 |
| | 155 | 0.000 | 0.000 | 0.035 | 0.000 | 0.050 |
| | 157 | 0.000 | 0.000 | 0.000 | 0.012 | 0.033 |
| | 159 | 0.000 | 0.000 | 0.000 | 0.023 | 0.050 |
| | 161 | 0.000 | 0.000 | 0.012 | 0.000 | 0.083 |
| | 163 | 0.000 | 0.000 | 0.023 | 0.000 | 0.000 |
| | 165 | 0.000 | 0.000 | 0.000 | 0.000 | 0.033 |
| | 167 | 0.000 | 0.000 | 0.023 | 0.000 | 0.050 |
| | 169 | 0.000 | 0.000 | 0.000 | 0.000 | 0.033 |
| | 171 | 0.000 | 0.000 | 0.012 | 0.000 | 0.017 |
| | 177 | 0.000 | 0.000 | 0.012 | 0.000 | 0.000 |
| | 183 | 0.000 | 0.000 | 0.000 | 0.000 | 0.017 |
| | 185 | 0.000 | 0.000 | 0.000 | 0.000 | 0.017 |
| | 187 | 0.000 | 0.000 | 0.012 | 0.000 | 0.000 |
| | 189 | 0.000 | 0.000 | 0.012 | 0.000 | 0.000 |
| | 191 | 0.000 | 0.000 | 0.012 | 0.000 | 0.000 |
| | 199 | 0.000 | 0.000 | 0.000 | 0.000 | 0.017 |
| | 201 | 0.000 | 0.000 | 0.000 | 0.000 | 0.017 |
| No. of samples | | 45 | 31 | 43 | 43 | 30 |
| No. of allele | | 10 | 7 | 19 | 13 | 19 |
| Effective no. of alleles[1] | | 3.58 | 2.34 | 8.47 | 5.65 | 7.87 |
| Observed heterozygosity ($H_o$) | | 0.689 | 0.548 | 0.884 | 0.884 | 0.867 |
| Expected heterozygosity ($H_e$) | | 0.721 | 0.573 | 0.882 | 0.823 | 0.873 |
| Ttho-7* | 182 | 0.000 | 0.031 | 0.000 | 0.000 | 0.000 |
| | 188 | 0.083 | 0.109 | 0.032 | 0.000 | 0.017 |
| | 190 | 0.024 | 0.000 | 0.021 | 0.010 | 0.033 |
| | 192 | 0.024 | 0.063 | 0.075 | 0.052 | 0.000 |
| | 194 | 0.321 | 0.203 | 0.287 | 0.250 | 0.217 |
| | 196 | 0.012 | 0.063 | 0.096 | 0.146 | 0.117 |
| | 198 | 0.024 | 0.078 | 0.053 | 0.125 | 0.317 |
| | 200 | 0.083 | 0.172 | 0.074 | 0.135 | 0.050 |
| | 202 | 0.012 | 0.031 | 0.053 | 0.042 | 0.083 |
| | 204 | 0.024 | 0.000 | 0.011 | 0.031 | 0.017 |
| | 206 | 0.012 | 0.016 | 0.043 | 0.000 | 0.017 |
| | 208 | 0.060 | 0.078 | 0.032 | 0.031 | 0.050 |
| | 210 | 0.131 | 0.063 | 0.075 | 0.042 | 0.017 |
| | 212 | 0.107 | 0.000 | 0.021 | 0.063 | 0.067 |
| | 214 | 0.024 | 0.031 | 0.053 | 0.031 | 0.000 |
| | 216 | 0.024 | 0.031 | 0.043 | 0.010 | 0.000 |
| | 218 | 0.024 | 0.031 | 0.032 | 0.010 | 0.000 |
| | 224 | 0.000 | 0.000 | 0.000 | 0.010 | 0.000 |
| No. of samples | | 42 | 32 | 47 | 48 | 30 |
| No. of allele | | 16 | 14 | 16 | 15 | 12 |
| Effective no. of allele | | 6.49 | 8.93 | 8.06 | 7.63 | 5.59 |
| Observed heterozygosity ($H_o$) | | 0.857 | 0.906 | 0.872 | 0.896 | 1.000 |
| Expected heterozygosity ($H_e$) | | 0.846 | 0.888 | 0.876 | 0.869 | 0.821 |

[1] Effective number of alleles was calculated with the equation $1/(1-H_e)$.

samples deviated considerably from zero ($F_{ST}$=0.018 to 0.070, $P$<0.001) for all loci. Significant departures of $F_{ST}$ from zero were observed for three loci (*Ttho-1**, *Ttho-4**, and *Ttho-6**) among all samples except the NE Atlantic sample ($F_{ST}$ =0.013 to 0.085, $P$<0.005) and among the three Pacific samples ($F_{ST}$=0.018 to 0.074, $P$<0.001). There were also significant differences in allele frequencies between some pairwise comparisons (Table 2). At *Ttho-1**, the SW Pacific sample showed significant difference from all other samples (all $P$<0.001), and there was also a significant difference between the SE Pacific and SW Atlantic samples ($P$=0.003). The NE Atlantic sample was significantly heterogeneous in comparison with all other samples in *Ttho-4** (all $P$<0.001). Difference between the NW and SE Pacific samples was also significant in this locus ($P$=0.001). In *Ttho-6** the NE Atlantic sample was again significantly heterogeneous in comparison with all other samples (all $P$<0.001). Furthermore, there were also significant differences between the NW and SE Pacific, between the SW and SE Pacific, and between the SW Pacific and SW Atlantic samples in this locus (all $P$<0.001). In *Ttho-7** the NE Atlantic sample showed significant differences from all three Pacific samples ($P$<0.001 for the NW and SW Pacific, $P$=0.003 for the SE Pacific) but not from the SW Atlantic sample ($P$=0.017).

**Table 2**

Pairwise comparison of $F_{ST}$ between five albacore samples.

| | NW Pacific | SW Pacific | SE Pacific | SW Atlantic |
|---|---|---|---|---|
| Ttho-1* | | | | |
| SW Pacific | 0.1401 | | | |
| SE Pacific | 0.005 | 0.099[1] | | |
| SW Atlantic | 0.018 | 0.249[1] | 0.0461 | |
| NE Atlantic | –0.004 | 0.154[1] | –0.003 | 0.023 |
| Ttho-4* | | | | |
| SW Pacific | 0.003 | | | |
| SE Pacific | 0.032[1] | 0.012 | | |
| SW Atlantic | 0.006 | –0.001 | 0.017 | |
| NE Atlantic | 0.151[1] | 0.131[1] | 0.094[1] | 0.131[1] |
| Ttho-6* | | | | |
| SW Pacific | 0.020 | | | |
| SE Pacific | 0.038[1] | 0.100[1] | | |
| SW Atlantic | 0.006 | 0.052[1] | 0.010 | |
| NE Atlantic | 0.135[1] | 0.208[1] | 0.0431 | 0.086[1] |
| Ttho-7* | | | | |
| SW Pacific | 0.012 | | | |
| SE Pacific | 0.006 | 0.003 | | |
| SW Atlantic | 0.020 | 0.006 | 0.001 | |
| NE Atlantic | 0.592[1] | 0.042[1] | 0.040[1] | 0.017 |

[1] Significant difference ($P$<0.005).

## Discussion

Because the number of alleles observed in microsatellite loci is usually large and the frequency of each allele may be low, a large sample size is necessary for satisfying subsequent statistic analyses. Ruzzante (1997) showed that a sample size of 50 <$n$< 100 individuals is generally satisfactory, although this size depends on allele number and frequency. Size of our samples ranged from 32 to 48, close to the lower margin of this threshold. Nevertheless, the distinct status of the Northeast Atlantic albacore sample from others is obvious. Although our study shared the same sample lots with Chow and Ushiama (1995), their mtDNA analysis revealed much less genetic differentiation between samples from the North and South Atlantic. MtDNA analysis is thought to be a more effective indicator for population subdivision than nuclear DNA (Nei and Li, 1979). But for albacore in the Atlantic, this is obviously not the case. No selection toward microsatellite alleles is obvious; therefore, differences in evolutionary rate between mtDNA and nDNA may explain differences in allele frequency distribution in the Atlantic albacore. Because mtDNA variation observed by Chow and Ushiama (1995) was much lower than the microsatellite DNA variation observed in our present study, there might have been insufficient elapsed time for unique mitochondrial genotypes to have arisen within the existing stocks. Thus, the rapidly evolving microsatellites appear to reflect albacore population subdivision.

MtDNA analyses have also failed to detect genetic difference between samples from northern and southern hemispheres within the Pacific (Chow and Ushiama, 1995). In contrast, present microsatellite analysis detected significant differences among three Pacific samples, clearly indicating that microsatellites appear to be more sensitive and powerful in detecting more subtle signals of genetic differentiation in albacore samples than mtDNA analysis. These results support several ecological and morphometric studies (Nakamura, 1969; Lewis, 1990) which assumed negligible migration of albacore across the equator in the Pacific. Separate North and South Pacific albacore stocks is a reasonable assumption because two major spawning grounds confined in the western to mid tropical Pacific are spatiotemporarily separated (Nishikawa et al., 1985). However, it is difficult to explain genetic differentiation between southwest and southeast Pacific albacore samples because no major spawning ground of albacore has been determined in the southeast Pacific. Microsatellite analysis of a sample from a different year class and a larger sample size is necessary to better define the observed genetic differences among the Pacific samples and the South Atlantic sample.

## Acknowledgments

We gratefully acknowledge R. D. Ward, Commonwealth Scientific and Industrial Research Organization, Australia, for reading the manuscript. This study was partially supported by the Fishery Agency of Japan.

## Literature cited

Chow, S., and H. Kishino.
1995. Phylogenetic relationships between tuna species of the genus *Thunnus* (Scombridae: Teleostei): Inconsistent implications from morphology, nuclear and mitochondrial genomes. J. Mol. Evol. 41:741–748.

Chow, S., and H. Ushiama.
1995. Global population structure of albacore (*Thunnus alalunga*) inferred by RFLP analysis of the mitochondrial ATPase gene. Mar. Biol. 123:39–45.

Ishii, T.
1965. Morphometric analysis of the Atlantic albacore populations mainly her eastern areas. Bull. Jap. Soc. Soc. Fish. 31:333–339.

Kurogane, K., and Y. Hiyama.
1958. Morphometric comparison of the albacore from the northwest, the equatorial and the southwest Pacific. Res. Oceanogr. Wks. Japan 4:64–75.

Lewis, A. D.
1990. South Pacific albacore stock structure: a review of available information. 3rd South Pacific Commission, Noumea, New Caledonia, p. 1–13.

Nakamura, H.
1969. Tuna distribution and migration. Fishing News Ltd, London, 76 p.

Nei, M., and W. H. Li.
1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. Proc. Natl. Acad. Sci. U.S.A. 76:5269–5273.

Nishikawa, Y., M. Honma, S. Ueyanagi, and S. Kikawa.
1985. Average distribution of larvae of oceanic species of scombrid fishes 1956–1981, vol. 12. National Research Institute of Far Seas Fisheries, Shimizu, 99 p.

Ruzzante D. E.
1997. A comparison of several measures of genetic distance and population structure with microsatellite data: bias and sampling variance. Can. J. Fish. Aquat. Sci. 55:1–14.

Schneider, S., J. M. Kueffer, D. Roessli, and L. Excoffier.
1997. Arlequin version 1.1. Univ. Geneva, Geneva, Switzerland.

Takagi, M., T. Okamura, S. Chow, and N. Taniguchi.
1999. PCR primers for microsatellite loci in tuna species of the genus *Thunnus* and its application for population genetic study. Fish. Sci. 65:571–576.

## Acknowledgment of reviewers

The editorial staff of *Fishery Bulletin* would like to thank the following referees for their time and efforts in providing reviews of the manuscripts published in 2000–2001. Their contributions have helped ensure the publication of quality science.

Dr. A. Acosta
Dr. Peter B. Adams
Dr. Juergen Alheit
Dr. Robert J. Allman
Dr. John Anderson
Dr. Todd W. Anderson
Dr. Michael P. Armstrong
Dr. Herbert M. Austin
Dr. Donald M. Baltz
Dr. William H. Bayliff
Dr. Terry D. Beacham
Dr. Patricia M. Biesiot
Dr. Asmund Bjorndal
Dr. Calvin L. Blood
Dr. James A. Bohnsack
Dr. A. J. Booth
Dr. Brian W. Bowen
Dr. Paul A. Breen
Dr. Richard W. Brill
Dr. Richard D. Brodeur
Dr. P. J. Bromley
Dr. Edward B. Brothers
Mr. Gerald Brothers
Dr. Richard Broughton
Dr. Bonnie L. Brown
Mr. Jay Burnett
Mr. Michael Burton
Dr. John Calambokidis
Dr. Steve Campana
Dr. Martin Castonguay
Dr. Jose J. Castro
Mr. Albert E. Caton
Dr. Patrice Cayre
Dr. Milani Y. Chaloupka
Dr. Robert W. Chapman
Dr. David M. Checkley Jr.
Dr. Yong Chen
Dr. V. C. Chong
Dr. Thomas A. Clarke
Dr. M. Comeau
Dr. Joseph Conti
Dr. Jean Cramer
Dr. Matthew A. Cronin
Dr. Willie Davidson
Dr. R. W. Davis
Dr. J. T. DeAlteris
Dr. Lawrence B. DeLancey
Dr. Jean Diebolt
Dr. Edward E. DeMartini
Dr. Michael L. Domeier
Dr. David Donald
Dr. Janet T. Duffy-Anderson
Dr. Jakov Dulcic
Dr. John S. Edmonds
Dr. Nicholas G. Elliot
Dr. Darryl Felder
Dr. Dagmar Fertl
Dr. Gary R. Fitzhugh
Mr. John Foster
Dr. Arild Folkvord
Dr. Clive Fox
Dr. James S. Franks
Dr. S. Fuwa
Dr. R. W. Gauldie
Dr. Tony Gharrett
Dr. Angel F. Gonzalez
Mr. C. Phillip Goodyear
Dr. E. P. Goubanov
Dr. John E. Graves
Dr. John Gunn
Dr. Manuel Haimovici
Dr. Peter Hanson
Dr. Patrick J. Harris
Ms. Dana-Lyn Hartley
Dr. Kyle J. Hartman
Ms. L. A. Harwood
Ms. Ruth Haas-Castro
Dr. Edward J. Heist
Dr. J.R.G. Hislop
Dr. Kim M. Holland
Dr. Edward D. Houde
Mr. C. J. Jackson
Dr. Thomas Jefferson
Dr. Simon Jennings
Dr. Naohisa Kanda
Dr. Jon B. Kristoffersen
Dr. Gordon H. Kruse
Ms. Kathy L. Lang
Dr. Robert R. Lauth
Dr. M. Kimberly Lowe
Dr. Molly E. Lutcavage
Dr. Charles D. Madenjian
Dr. Sherry C. Manickshand-Heileman
Dr. F. Marsac
Dr. Andrew Martin
Dr. T. Matsumoto
Dr. Kathleen R. Matthews
Dr. J. Mattila
Dr. John McDermott
Dr. A. J. McGregor
Ms. Anne L. McMillen-Jackson
Dr. Richard D. Methot
Dr. Jaime Mejuto
Dr. Thomas J. Miller
Dr. David A. Milton
Dr. Beatrice Morales-Nin
Dr. Paul Moran
Mr. Bruce C. Mundy
Dr. James Nance
Dr. Stephen J. Newman
Dr. David L. Nieland
Dr. T. Okutani
Dr. John Olney
Dr. James D. Parrish
Dr. C. A. Pattison
Dr. Pierre Pepin
Mr. Jose Antonio Perez-Comas
Dr. Mark S. Peterson
Dr. Ruth Phillips
Dr. N. V. C. Polunin
Dr. Allyn B. Powell
Dr. Imants G. Priede
Dr. Vladimir Radchenko
Mr. Boodhum Ramcharrum
Dr. Kimberly L. Raum-Suryan
Dr. Catalina P. Raya
Dr. Andrew J. Read
Dr. D. Reznick
Dr. Fritz Rohde
Dr. Pierre Richard
Dr. William J. Richards
Dr. Linda R. Richardson
Dr. T. A. Rien
Dr. Axayacatl Rocha-Olivares
Dr. Callum M. Roberts
Dr. Roger A. Rulifson
Dr. Thomas L. Rutecki
Dr. Gary T. Sakagawa
Dr. Mark W. Saunders
Dr. Jeffrey R. Schmid
Dr. Joanne Schultz
Dr. Gerald P. Scott
Dr. Nancy Shackell
Dr. Michael D. Scott
Dr. George Sedberry
Dr. Tom Shirley
Dr. Malcolm J. Smale
Mr. Joseph W. Smith
Dr. Peter Smouse
Ms. Melissa Snover
Dr. Susan M. Sogard
Dr. Trisha Spears
Dr. Linda L. Stehlik
Dr. Bradley G. Stevens
Dr. Jill S. Stevenson
Dr. Kevin D. E. Stokesbury
Dr. Kenneth Sulak
Dr. J. S. Sunada
Dr. D. P. Swain
Mr. G. N. Swinney
Dr. Stephen T. Szedlmayer
Dr. Steven Thomas
Dr. Simon R. Thorrold
Dr. O. I. Tomakhina
Dr. Fred M. Utter
Dr. Douglas S. Vaughan
Dr. J. R. Voight
Dr. Stephen J. Walsh
Dr. Yoshiro Watanabe
Dr. Mason T. Weinrich
Mr. Andrew J. Westgate
Dr. Jerry A. Wetherall
Dr. Gary Winans
Dr. Peter R. Witthames
Dr. Cheryl M. Woodley
Dr. Yongshun Xiao
Dr. Dirk Zeller

The index and/or table of contents has been removed and photographed separately within this volume year.

For roll film users, this information for the current volume year is at the beginning of the microfilm. For a prior year volume, this information is at the end of the microfilm.

For microfiche users, the index and/or contents is contained on a separate fiche.

# Erratum

*Fishery Bulletin* **99(3):399–409**

**Colura, Robert L., and Britt W. Bumguardner**

**Effect of the salt-box catch-bycatch separation procedure, as used by the Texas shrimp industry, on short-term survival of bycatch**

Erratum. Page 400, left column, last four lines.

The sentence "For each species, probit analysis was conducted on untransformed data to determine the time at which 50% of the test animals exposed to the test salinity died ($LE_{50}$) (Sokal and Rohlf, 1981)" should read

"For each species (48 hour), probit analysis was conducted on untransformed data to determine the time at which 50% of the test animals exposed to the test salinity died ($LE_{50}$) (Sokal and Rohlf, 1981)."

Superintendent of Documents **Publications** Order Form

*5178

☐ YES, please send me the following publications:

____ Subscriptions to *Fishery Bulletin*
for $45.00 per year ($56.25 foreign)

The total cost of my order is $ _____ . Prices include regular domestic postage and handling and are subject to change.

_____
(Company or Personal Name) (Please type or print)

_____
(Additional address/attention line)

_____
(Street address)

_____
(City, State, ZIP Code)

_____
(Daytime phone including area code)

_____
(Purchase Order No.)

Charge your order.
IT'S EASY!

VISA

MasterCard

**Please Choose Method of Payment:**

☐ Check Payable to the Superintendent of Documents
☐ GPO Deposit Account ☐☐☐☐☐☐–☐
☐ VISA or MasterCard Account
☐☐☐☐☐☐☐☐☐☐☐☐☐☐☐☐
☐☐☐☐ (Credit card expiration date)

**To fax your orders (202) 512-2250**

_____
(Authorizing Signature)

Mail To: Superintendent of Documents
P.O. Box 371954, Pittsburgh, PA 15250-7954

***Thank you for your order!***

This statement is required by the Act of August 12, 1970, Section 3685, Title 39, U.S. Code, showing ownership, management, and circulation of the *Fishery Bulletin*, publication number 366-370, and was filed on 14 September, 2001. The *Bulletin* is published quarterly (four issues annually) with an annual subscription price of $50.00 (sold by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402). The complete mailing address of the office of publication is NMFS Scientific Publications Office, NOAA, 7600 Sand Point Way NE, BIN C15700, Seattle, WA 98115. The complete mailing address of the headquarters of the publishing agency is National Marine Fisheries Service, NOAA, Department of Commerce, 1335 East-West Highway, Silver Spring, MD 20910. The name of the publisher is Willis Hobart and the managing editor is Sharyn Matriotti; their mailing address is: NMFS Scientific Publications Office, 7600 Sand Point Way NE, BIN 15700, Seattle, WA 98115. The owner is the U.S. Department of Commerce, 14th St. N.W., Washington, DC 20230; there are no bondholders, mortgages, or other security holders. The purpose, function, and nonprofit status of the organization (agency) and the exempt status for Federal income tax purposes has not changed during the preceding 12 months. The extent and nature of circulation is as follows: total number of copies (A) (average number of copies of each issue during the preceding 12 months) was 1918 and the actual number of copies of the single issue published nearest to the filing dates was 1906. Paid circulation (B) is handled by the U.S. Government Printing Office, Washington, DC 20402, and the total number printed for sales (mail subscriptions and individual sales) was 488 for both the average number of copies each issue during the preceding 12 months and 475 the actual number of copies of the single issue published nearest to the filing date (C). Free distribution (D) by mail; samples, complimentary and other free copies (average number of copies each issue during the preceding 12 months) was 1405 and the actual number of copies of the single issue published nearest to the filing date was 1418. Free distribution outside the mail (E) by carriers or other means was 0 for both average number of copies and actual number of copies. Total free distribution (F) was 0 for both average number of copies and actual number of copies of the single issue published nearest the filing date. The total distribution (G: sum of D and B) (average number of copies each issue during the preceding 12 months) was 1893 and the actual number of copies of the single issue published nearest to the filing date was 1881. There were 25 copies not distributed (H). The total (I: sum of G and H) is equal to the net press run figures shown in Item A: 1893 and 25 copies, respectively. I certify that the statements made by me above are correct and complete: (Signed) Willis Hobart, Publisher.

# Fishery Bulletin

## Guidelines for contributors

### Content of papers

#### Articles

Articles are reports of 10 to 30 pages (double spaced) that describe original research in one or a combination of the following fields of marine science: taxonomy, biology, genetics, mathematics (including modeling), statistics, engineering, economics, and ecology.

#### Notes

Notes are reports of 5 to 10 pages without an abstract that describe methods and results not supported by a large body of data. Although all contributions are subject to peer review, responsibility for the contents of articles and notes rests upon the authors and not upon the editor or the publisher. It is therefore important that authors consider the contents of their manuscripts carefully. Submission of an article is un-derstood to imply that the article is original and is not being considered for publication elsewhere. Manuscripts must be written in English. Authors whose native language is not English are strongly advised to have their manuscripts checked for fluency by English-speaking colleagues prior to submission.

### Preparation of papers

#### Text

**Title page** should include authors' full names and mailing addresses (street address required) and the senior author's telephone, fax number, e-mail address, as well as a list of key words to describe the contents of the manuscript. **Abstract** must be less than one typed page (double spaced) and must not contain any citations. It should state the main scope of the research but emphasize the author's conclusions and relevant findings. Because abstracts are circulated by abstracting agencies, it is important that they represent the research clearly and concisely. **General text** must be typed in double-spaced format. A brief introduction should state the broad significance of the paper; the remainder of the paper should be divided into the following sections: Materials and methods, Results, Discussion (or Conclusions), and Acknowledgments. Headings within each section must be short, reflect a logical sequence, and follow the rules of multiple subdivision (i.e. there can be no subdivision without at least two subheadings). The entire text should be intelligible to interdisciplinary readers; therefore, all acronyms and abbreviations should be written out and all lesser-known technical terms should be defined the first time they are mentioned. The scientific names of species must be written out the first time they are mentioned; subsequent mention of scientific names may be abbreviated. Follow *Scientific style and format: CBE manual for authors, editors, and publishers* (6th ed.) for editorial style and the most current issue of the *American Fisheries Society's common and scientific names of fishes from the United States and Canada* for fish nomenclature. Dates should be written as follows: 11 November 1991. Measurements should be expressed in metric units, e.g. metric tons (t). The numeral one (1) should be typed as a one, not as a lower-case el (l).

#### Footnotes

Use footnotes to add editorial comments regarding claims made in the text and to document unpublished works or works with local circulation. Footnotes should be numbered with Arabic numerals and inserted in 10-point font at the bottom of the first page on which they are cited. Footnotes should be formatted in the same manner as citations. If a manuscript is unpublished, in the process of review, or if the information provided in the footnote has been conveyed verbally, please state this information as "unpubl. data," "manuscript in review," and "personal commun.," respectively. Authors are advised wherever possible to avoid references to nonstandard literature (unpublished literature that is difficult to obtain, such as internal reports, processed reports, administrative reports, ICES council minutes, IWC minutes or working papers, any "research" or "working" documents, laboratory reports, contract reports, and manuscripts in review). If these references are used, please indicate whether they are available from NTIS (National Technical Information Service) or from some other public depository. Footnote format: author (last name, followed by first-name initials); year; title of report or manuscript; type of report and its administrative or serial number; name and address of agency or institution where the report is filed.

#### Literature cited

The literature cited section comprises works that have been published and those accepted for publication (works in press) in peer-reviewed journals and books. Follow the name and year system for citation format. In the text, write "Smith and Jones (1977) reported" but if the citation takes the form of parenthetical matter, write "(Smith and Jones, 1977)." In the literature cited section, list citations alphabetically by last name of senior author: For example, Alston, 1952; Mannly, 1988; Smith, 1932; Smith, 1947; Stalinsky and Jones, 1985. Abbreviations of journals should conform to the abbreviations given in the *Serial sources for the BIOSIS previews database*. Authors are responsible for the accuracy and completeness of all citations. Literature citation format: author (last name, followed by first-name initials); year; title of report or article; abbreviated title of the journal in which the article was published, volume number, page numbers. For books, please provide publisher, city, and state.

#### Tables

Tables should not be excessive in size and must be cited in numerical order in the text. Headings in tables should be short but ample enough to allow the table to be intelligible on its own. All unusual symbols must be explained in the table legend. Other incidental comments may be footnoted (use italic arabic numerals for footnote markers). Use asterisks only to indicate probability in statistical data. Place table legends on the same page as the table data. We accept tables saved in most spreadsheet software programs (e.g. Microsoft Excel). Please note the following:

- Use a comma in numbers of five digits or more (e.g. 13,000 but 3000).
- Use zeros before all decimal points for values less than one (e.g. 0.31).

#### Figures

Figures include line illustrations, computer-generated line graphs, and photographs (or slides). They must be cited in numerical order in the text. Line illustrations are best submitted as original drawings. Computer-generated line graphs should be printed on laser-quality paper. Photographs should be submitted on glossy paper with good contrast. All figures are to be labeled with senior author's name and the number of the figure (e.g. Smith, Fig. 4). Use Helvetica or Arial font to label anatomical parts (line drawings) or variables (graphs) within figures; use Times Roman bold font to label the different sections of a figure (e.g. **A**, **B**, **C**). Figure legends should explain all symbols and abbreviations seen within the figure and should be typed in double-spaced format on a separate page at the end of the manuscript. We advise authors to peruse a recent issue of *Fishery Bulletin* for standard formats. Please note the following:

- Capitalize the first letter of the first word of axis labels.
- Do not use overly large font sizes to label axes or parts within figures.
- Do not use boldface fonts within figures.
- Do not create outline rules around graphs.
- Do not use horizontal lines through graphs.
- Do not use large font sizes to label degrees of longitude and latitude on maps.
- Indicate direction of degrees longitude and latitude on maps (e.g. 170°E).
- Avoid placing labels on a vertical plane (except on *y* axis).
- Avoid odd (nonstandard) patterns to mark sections of bar graphs and pie charts.

#### Copyright law

*Fishery Bulletin*, a U.S. government publication, is not subject to copyright law. If an author wishes to reproduce any part of *Fishery Bulletin* in his or her work, he or she is obliged, however, to acknowledge the source of the extracted literature.

### Submission of papers

Send four printed copies (one original plus three copies)—clipped, *not stapled*—to the Scientific Editor, at the address shown below. Send photocopies of figures with initial submission of manuscript. Original figures will be requested later when the manuscript has been accepted for publication. Do not send your manuscript on diskette until requested to do so.

Dr. John V. Merriner
Scientific Editor, *Fishery Bulletin*
Center for Coastal Fisheries and Habitat Research, NOS
101 Pivers Island Road
Beaufort, NC 28516

Once the manuscript has been accepted for publication, you will be asked to submit a software copy of your manuscript. The software copy should be submitted in WordPerfect or Word format (in Word, save as Rich Text Format). Please note that we do not accept ASCII text files.

#### Reprints

Copies of published articles and notes are available free of charge to the senior author (50 copies) and to his or her laboratory (50 copies). Additional copies may be purchased in lots of 100 when the author receives page proofs.




1.0
1.1
1.25
1.4
1.6
1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0
4.5